

પત્રક-એ

હવામાન આગાહી આધારીત કૃષિ સલાહ બુલેટીન

(સમય તા : ૨૪.૦૮.૧૩ થી ૨૮.૦૮.૧૩)

પાછલા અઠવાડીયાનું હવામાન

હવામાન પરીબળો	કૃષિ આબોહવાક્ય વિસ્તાર દક્ષિણ સૌરાષ્ટ્ર
વરસાદ (મીભી)	0.00
મહત્વમાન તાપમાન (સે.ગે.)	31.3
લઘુત્વમાન તાપમાન (સે.ગે.)	24.5
ભેજ(%) (સવાર/બપોરબાદ)	89-66
પવનની ગતિ (કિભી/કલાક)	6.7

પત્રક-૬ : જિલ્લાવાર હવામાન આગાહી (તા. ૨૮.૦૮.૧૩ સુધી)

હવામાન પરીબળો	તારીખ	દક્ષિણ સૌરાષ્ટ્ર	
		જુનાગઢ	પોરબંદર
વરસાદ (મીમી)	૨૪.૦૮.૧૩	૦	૦
	૨૫.૦૮.૧૩	૦	૦
	૨૬.૦૮.૧૩	૦	૦
	૨૭.૦૮.૧૩	૦	૦
	૨૮.૦૮.૧૩	૦	૦
મહત્વમાન તાપમાન (સે.ગ્રે.)	૨૪.૦૮.૧૩	૩૦	૩૦
	૨૫.૦૮.૧૩	૩૦	૩૦
	૨૬.૦૮.૧૩	૩૧	૩૧
	૨૭.૦૮.૧૩	૩૧	૩૧
	૨૮.૦૮.૧૩	૩૧	૩૨
લઘુત્વમાન તાપમાન (સે.ગ્રે.)	૨૪.૦૮.૧૩	૨૪	૨૬
	૨૫.૦૮.૧૩	૨૪	૨૬
	૨૬.૦૮.૧૩	૨૪	૨૬
	૨૭.૦૮.૧૩	૨૪	૨૬
	૨૮.૦૮.૧૩	૨૫	૨૬
વાદળની સ્થિતિ (ઓક્ટો)	૨૪.૦૮.૧૩	મોટેભાગે વાદળણાયું	મોટેભાગે વાદળણાયું
	૨૫.૦૮.૧૩	મોટેભાગે વાદળણાયું	મોટેભાગે વાદળણાયું
	૨૬.૦૮.૧૩	અંશતઃ વાદળણાયું	અંશતઃ વાદળણાયું
	૨૭.૦૮.૧૩	અંશતઃ વાદળણાય	અંશતઃ વાદળણાય
	૨૮.૦૮.૧૩	અંશતઃ વાદળણાય	અંશતઃ વાદળણાય
મહત્વમાન ભેજ (%)	૨૪.૦૮.૧૩	૮૩	૮૨
	૨૫.૦૮.૧૩	૮૩	૮૩
	૨૬.૦૮.૧૩	૮૪	૮૪
	૨૭.૦૮.૧૩	૮૪	૮૪
	૨૮.૦૮.૧૩	૮૫	૮૫
લઘુત્વમાન ભેજ (%)	૨૪.૦૮.૧૩	૭૫	૭૬
	૨૫.૦૮.૧૩	૭૬	૭૬
	૨૬.૦૮.૧૩	૭૫	૭૫
	૨૭.૦૮.૧૩	૭૫	૭૪
	૨૮.૦૮.૧૩	૭૬	૭૬
પવનની ગતિ (કિમી/કલાક)	૨૪.૦૮.૧૩	૦૨૦	૦૧૯
	૨૫.૦૮.૧૩	૦૧૯	૦૧૭
	૨૬.૦૮.૧૩	૦૨૦	૦૧૯
	૨૭.૦૮.૧૩	૦૧૯	૦૧૭
	૨૮.૦૮.૧૩	૦૨૦	૦૧૭
પવનની દિશા (ડિગ્રી)	૨૪.૦૮.૧૩	૨૬૩	૨૬૩
	૨૫.૦૮.૧૩	૨૫૧	૨૫૦
	૨૬.૦૮.૧૩	૨૩૯	૨૩૫
	૨૭.૦૮.૧૩	૨૩૯	૨૩૬
	૨૮.૦૮.૧૩	૨૪૪	૨૪૦

પત્રક-ક

જિલ્લાવાર કૃષિ સલાહ

ખેત આબોહવાક્ય વિસ્તાર	પાકનું નામ	કૃષિ સલાહ
	મગફળી	<ul style="list-style-type: none"> ➢ પાનના ચાંચવાના નિયત્રણ માટે મીથાઈલ પેરાથીયોન ૧૦ મીલી અથવા ક્વીનાલ ફોસ ૨૦ મીલી અથવા કાબાંરીલ ૪૦ ગ્રામ અથવા એસીફેટ ૧૫ ગ્રામ ૧૦ લિટર પાણીમા ભેળવી છંટકાવ કરવો. ➢ મગફળી મા ઘૈણ ના નિયત્રણ માટે કલોરપાયરીફોસ દવા હેકટરે ૪ લિટર પ્રમાણે પિયત ના પાણી સાથે આપવી અથવા કલોરપાયરીફોસ દવા ૨૫ મીલી ૧૦ લિટર પાણી મા ઓગાળી આ પ્રવાહી મીશ્રણ ને છોડ ના મુળ પાસે અને જમીન મા ઉત્તરે તે રીતે પંપ ની નોઝલ કાઢી રેડવાથી નિયત્રણ મેળવી શકય છે ➢ પાક મા ટીક રોગ ના નિયત્રણ માટે મેન્કોઝેબ ૨૫ ગ્રામ અથવા ટેબુકોનાજોલ ૧૦ મીલી/ ૧૦ લિટર પાણીમા ભેળવી છંટકાવ કરવો. ➢ વધુ વરસાદ ના કારણે પાક પીળો પડી ગયેલ હોય તો ફેરસ સલ્ફેટ (હીરાકસી) ૧૦૦ ગ્રામ અને સાઇટ્રી ક એસીડ (લીમ્બુ ના ફૂલ) ૧૦ ગ્રામ ૧૦ લિટર પાણી મા ઓગાળી છંટકાવ કરવો..
દક્ષિણ સૌરાષ્ટ્ર	કપાસ	<ul style="list-style-type: none"> ❖ કપાસ મા ખૂણીયા ટપકાના રોગના નિયત્રણ માટે સ્યુડોમોનાસ ફલુરોસન્સ ૧૦ મીલી/૧૫ લિટર પાણી મા ઓગાળી વાવણી બાદ ૩૦, ૪૦, ૭૦, અને ૮૦ દીવસે છંટકાવ કરવો. ❖ કપાસ મા ચુસીયા પ્રકાર ની જીવાત ના નિયત્રણ માટે ડાઈમીથોયેટ ૧૫ મીલી/૧૫ લિટર પાણી અથવા મીથાઈલઓડેમેટોન ૩૦ મીલી/૧૫ લિટર પાણી અથવા ઈમીડાકલોપ્રાઇંડ ૫ મીલી/૧૫ લિટર પાણી નો છંટકાવ કરવો. ❖ વધુ વરસાદ ના કારણે પાક પીળો પડી ગયેલ હોય તોવ ૨ ટકા ઉરીયા ના દ્રાવણ નો છંટકાવ કરવો..
	તલ	<ul style="list-style-type: none"> ▪ પાક ના વાવેતર બાદ ૪૦ થી ૪૫ દીવસે પુરતી ખાતર હેકટર દીઠ ૧૨.૫ કીગ્રા (૨૭ કીગ્રા યુરીયા) મુજબ આપવું.
	દિંવેલા	<ul style="list-style-type: none"> ➢ પાક મા ઘોડીયા ઈયળ ના નીયંત્રણ માટે ડીડીવીપી દવા ૫ મીલી ૧૦ લિટર પાણી મા ઓગાળી છંટકાવ કરવો..
	ભીડો	કુખ અને ફળ કોરી ખાનાર ઈયળ નીયંત્રણ માટે લીમ્બોળી ના બીજ નો અંક ૫૦૦ મીલી અથવા લીમ્બોળી નુ તેલ ૫૦ મીલી ૧૦ લિટર પાણી મા ઓગાળી ૧૫ દીવસ ના અંતરે ૪

		છંટકાવ કરવા.
	મરચા	પાક મા સુકારા રોગ ના નીચત્રણ માટે ટ્રાઇકોડરમા વીરીડી ૪૦ ગ્રમ ૧૦ લિટર પાણી મા ઓગાળી છોડ ના થડ ની પાસે ડેચિંગ કરવુ
	પશુપાલન	<ul style="list-style-type: none"> ❖ ઉછરતા પશુ ઓમા બૃષેલાકાલ્ફવૂડ નુ રસી કરણ કરવુ. ❖ ઈતરડી ના નીચત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાજ દવા ૧૦ લિટર પાણીમા નાખી છંટકાવ કરવો. ❖ પુષ્ટ પશુ ને ૬ થી ૮ કિલો સારી ગુણવત્તા વાલો સુકો ચારો અને ૧૫ થી ૨૦ કિલો લીલો ચારો આપવો . ❖ ચોમાસા દરમ્યાન જાનવરોને માખીઓ અને મછરોના ઉપક્રમ થી બચાવવા જાનવરો ના રહેઠાણ ની જગ્યાએ ફીનાઈલ નો છંટકાવ કરવો

JUNAGADH

Phone: 0285-2670139

Date: 23-08-2013

(Period 24 to 28 August -2013)

Significant weather for the preceding week

Rain fall (mm)	000.0
Maximum temperature °C	30.5-31.5
Minimum temperature °C	24.3-25.0
RH % (Morning/Afternoon)	89-66
Wind speed (Kmph)	6.3-7.4
Cloud cover	5-7

Weather forecast for Junagadh Dist. until 08:30 hrs of 28- 08-2013

Parameters	Day-1 24-08-13	Day-2 25-08-13	Day-3 26-08-13	Day-4 27-08-13	Day-5 28-08-13
Rain fall (mm)	0	0	0	0	0
Maximum temperature °C	30	30	31	31	31
Minimum temperature °C	24	24	24	24	25
Cloud cover (Okta)	6	6	5	5	5
RH % (Morning/After noon)	83	83	84	84	85
Wind speed (Kmph)	020	019	020	019	020
Wind direction(Deg)	263	251	239	239	244

Agro meteorological Advisories

Crop	Stage	Agromet Advisories
Groundnut	Pegging	<ul style="list-style-type: none"> - For the control of leaf weevil, spray methyl parathion 10ml or quinalphos 20ml or carbaryl 40gm or acephate 15 gm in 10 litres of water. - For the control of white grub, apply Chlorpyriphos @ 4 litres/ha with irrigation water OR Mix Chlorpyriphos 25 ml in 10 liters of water , then drenching of this mixture should be done near the root zone of the crop. - For the control of tikka disease, spray Mancozeb @25gm/15 liter or tebuconazol 10 ml/10 litre of water. - Due to continues rain, if yellowing is found spray ferrous sulphate (Hiracasi) 100 gm and citric acid (limbu na phool) 10 gm/10 litres of water
Cotton	Branching/Squirting	<ul style="list-style-type: none"> - For the control of sucking pest, spray either Dimethoate @ 15 ml/ 15 liter water or Methyl-o-demeton @ 30 ml/ 15 liter water or Imidaclopride @ 5 ml/15 liter water. - For the control of angular leaf spot disease, foliar spray of <i>Pseudomonas fluorescens</i> @ 10 ml/ 15 liter water after 30, 50, 70, 90 days of sowing. - If yellowing is found spray 2 % solution of urea.
Sesame	Flowering	<ul style="list-style-type: none"> - For the control of leaf spot disease, two spraying of cabendenzim 0.02 % or Mancozeb 0.2 % for 25 gm/ 15 liter water should carried out.

Castor	Branching	- For the control of semi lopper pest, spray DDVP@5ml in 10 liter of water.
Ladies Finger	Flowering/ Fruiting	- For the control of Shoot and fruit borer, spray neem seed extract 500ml or neem seed oil 50 ml/10 liter of water at 15 days interval (total 4 spray).
Chili	Flowering/ Fruiting	- For the control of wilting disease, drenching of <i>Trichoderma viride</i> @40 gm/10 liter of water
Live stocks	-	<ul style="list-style-type: none"> ➤ Vaccination of Brucellosis calf wood to growing animals ➤ For the control of tick, spray Deltamathrin or Amitaz 20 ml/10 liter of water. ➤ Provide 6-8 kg of good quality of dry fodder with 15 to 20 kg green fodder to animal. ➤ Spray phenyl in animal shed to protect the animals against flies and mosquitoes

PORBANDAR

Phone : 0285-2670139

Date : 23-08-2013

(Period 24 to 28 August -2013)

Weather forecast for Porbandar Dist. until 08:30 hrs of 28- 08-2013

Parameters	Day-1 24-08-13	Day-2 25-08-13	Day-3 26-08-13	Day-4 27-08-13	Day-5 28-08-13
Rain fall (mm)	0	0	0	0	0
Maximum temperature °C	30	30	31	31	32
Minimum temperature °C	26	26	26	26	26
Cloud cover (Okta)	6	6	5	5	5
RH % (Morning/After noon)	82-76	83-76	84-75	84-74	85-76
Wind speed (Kmph)	019	017	019	017	017
Wind direction(Deg)	263	250	235	236	240

Agro meteorological Advisories

Crop	Stage	Agromet Advisories
Groundnut	Pegging	<ul style="list-style-type: none"> - For the control of leaf weevil, spray methyl parathion 10ml or quinalphos 20ml or carbaryl 40gm or acephate 15 gm in 10 litres of water. - For the control of white grub, apply Chlorpyriphos @ 4 litres/ha with irrigation water OR Mix Chlorpyriphos 25 ml in 10 liters of water , then drenching of this mixture should be done near the root zone of the crop. - For the control of tikka disease, spray Mancozeb @25gm/15 liter or tebuconazol 10 ml/10 litre of water. - Due to continues rain, if yellowing is found spray ferrous sulphate (Hiracasi) 100 gm and citric acid (limbu na phool) 10 gm/10 litres of water
Cotton	Branching/Squirting	<ul style="list-style-type: none"> - For the control of sucking pest, spray either Dimethoate @ 15 ml/ 15 liter water or Methyl-o-demeton @ 30 ml/ 15 liter water or Imidaclopride @ 5 ml/15 liter water. - For the control of angular leaf spot disease, foliar spray of <i>Pseudomonas fluorescens</i> @ 10 ml/ 15 liter water after 30, 50, 70, 90 days of sowing. - If yellowing is found spray 2 % solution of urea.
Sesame	Flowering	<ul style="list-style-type: none"> - For the control of leaf spot disease, two spraying of cabendenzim 0.02 % or Mancozeb 0.2 % for 25 gm/ 15 liter water should carried out.
Castor	Branching	<ul style="list-style-type: none"> - For the control of semi lopper pest, spray DDVP@5ml in 10 liter of water.
Ladies Finger	Flowering/Fruiting	<ul style="list-style-type: none"> - For the control of Shoot and fruit borer, spray neem seed extract 500ml or neem seed oil 50 ml/10 liter of water at 15 days interval (total 4 spray).

Chili	Flowering/ Fruiting	- For the control of wilting disease, drenching of <i>Trichoderma viride</i> @40 gm/10 liter of water
Live stocks	-	<ul style="list-style-type: none"> ➤ Vaccination of brucellosis calf wood to growing animals ➤ For the control of Tick, Spray delta mathrin or Amitaz 20 ml/10 liter of water. ➤ Provide 6-8 kg of good quality of dry fodder with 15 to 20 kg green fodder to animal. ➤ Spray phenyl in animal shed to protect the animals against flies and mosquitoes