

હવામાન આગાહી આધારીત કૃષિ સલાહ બુલેટીન

દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ

તા. ૦૧-૦૮-૨૦૧૮ થી તા. ૦૫-૦૮-૨૦૧૮

(ગ્રામીણ કૃષિ મૌસમ સેવા, જુનાગઢ કૃષિ યુનિવર્સિટી, જુ.કૃ.ય.)

હવામાન આગાહી આધારીત બુલેટીન : ૨૦૧૮-૧૯

પાછલા અઠવાડીયાનું જુનાગઢનું હવામાન

હવામાન પરીબળો	દક્ષિણ સૌરાષ્ટ્ર કૃષિ આબોહવાકીય વિસ્તાર
વરસાદ (મીમી)	૦૦૦.૦
મહત્વમાન તાપમાન (°સે.)	૩૧.૨
લઘુત્વમાન તાપમાન (°સે.)	૨૩.૬
ભેજ (%) (સવાર/બપોરબાદ)	૮૭ - ૭૪
પવનની ગતિ (કિમી/કલાક)	૦૫.૪

પત્રક-૬ : જુલ્લાવાર હવામાન આગાહી (તા.૦૫-૦૮-૨૦૧૮ સુધી)

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહીતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના

૧. જુનાગઢ:- જુલ્લામા આગામી તારીખ ૦૧/૦૮/૨૦૧૮ થી ૦૫/૦૮/૨૦૧૮ દરમ્યાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં અંશત વાદળછાયું થી વાદળછાયું રહેશે, આવતી કાલે હળવા વરસાદ ની સંભાવના રહેલી છે અને બાકીના દિવસોમાં વરસાદની કોઈ શક્યતાઓ રહેલી નથી અને પવનની ઝડપ ૧૨ થી ૧૯ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાન તાપમાન ૨૮ થી ૨૮ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્વમાન ૨૬ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

૨. પોરબંદર:- જુલ્લામા આગામી તારીખ ૦૧/૦૮/૨૦૧૮ થી ૦૫/૦૮/૨૦૧૮ દરમ્યાન આકાશ મુખ્યત્વે આગામી પાંચ દિવસોમાં અંશત વાદળછાયું થી વાદળછાયું રહેશે, આવતા બે દિવસ હળવા વરસાદ ની સંભાવના રહેલી છે અને બાકીના દિવસોમાં વરસાદની કોઈ શક્યતાઓ રહેલી નથી. પવનની ઝડપ ૧૭ થી ૨૩ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્વમાન તાપમાન ૨૮ થી ૨૮ ડિગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્વમાન ૨૫ ડિગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

જુનાગઢ

હવામાન પરીબળો / તારીખ	દિવસ-૧ ૦૧-૦૬-૨૦૧૮	દિવસ-૨ ૦૨-૦૬-૨૦૧૮	દિવસ-૩ ૦૩-૦૬-૨૦૧૮	દિવસ-૪ ૦૪-૦૬-૨૦૧૮	દિવસ-૫ ૦૫-૦૬-૨૦૧૮
વરસાદ (મી.મી)	૦૪	૦૦	૦૦	૦૦	૦૦
મહત્વમાન તાપમાન (સે.ગ્રે)	૨૯	૨૯	૨૯	૨૯	૨૯
લઘુત્વમાન તાપમાન (સે.ગ્રે)	૨૬	૨૬	૨૬	૨૬	૨૬
વાદળની સ્થિતિ	વાદળછાયું	વાદળછાયું	મોટેભાગે વાદળછાયું	મોટેભાગે વાદળછાયું	અંશત વાદળછાયું
સવારના ભેજના ટકા	૮૭	૮૫	૮૨	૮૨	૮૪
બપોર બાદ ભેજના ટકા	૭૫	૭૩	૬૫	૬૫	૬૫
પવનની ગતિ (કિમી/કલાક)	૦૧૭	૦૧૬	૦૧૬	૦૧૫	૦૧૨
પવનની દિશા (ડીગ્રી)	૨૪૩	૨૪૩	૨૪૬	૨૪૭	૨૪૮

પોરબંદર

વરસાદ (મી.મી)	૦૦	૦૩	૦૦	૦૦	૦૦
મહત્વમાન તાપમાન (સે.ગ્રે)	૨૯	૨૯	૨૯	૨૮	૨૮
લઘુત્વમાન તાપમાન (સે.ગ્રે)	૨૫	૨૫	૨૫	૨૫	૨૫
વાદળની સ્થિતિ	વાદળછાયું	વાદળછાયું	વાદળછાયું	મોટેભાગે વાદળછાયું	અંશત વાદળછાયું
સવારના ભેજના ટકા	૮૩	૮૩	૭૬	૭૭	૭૬
બપોર બાદ ભેજના ટકા	૭૫	૭૩	૬૭	૬૭	૬૫
પવનની ગતિ (કિમી/કલાક)	૦૨૩	૦૨૩	૦૨૧	૦૨૩	૦૧૭
પવનની દિશા (ડીગ્રી)	૨૪૩	૨૪૮	૨૪૬	૨૪૮	૨૪૪

દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાક્ય વિભાગ

પાકનું નામ	પાક અવસ્થા	રોગ / જીવાત / જાત	કૃષિ સલાહ
			<p>આવતી કાલે જુનાગઢ જીલ્લામાં હળવા વરસાદ ની સંભાવના રહેલી છે અને બાકીના દિવસોમાં વરસાદની કોઈ શક્યતાઓ રહેલી નથી જ્યારે પોરબંદર જીલ્લામા આવતા બે દિવસ હળવા વરસાદ ની સંભાવના રહેલી છે અને બાકીના દિવસોમાં વરસાદની કોઈ શક્યતાઓ રહેલી નથી.</p> <p>વરસાદની ખેડે જાણાય તો જમીનની પ્રતને ધ્યાને લઈ પાકને પીયત આપવું. (સુચા બેસવા, ડોડવા બંધાવા અને તેના</p>

			વિકાસની અવસ્થાએ પાણીની ખેંચ જણાય તો અવશ્ય પીયત આપવું.)
ચોમાસું મગફળી	ફૂલ / સુયા અવસ્થા	ઉંગસુક રોગ / ચુસીયા જીવાત	<ul style="list-style-type: none"> ❖ જો પાણીની અછત હોય તો મગફળી પાકમાં સુયા બેસવા વખતે અવશ્ય પીયત આપવું. ❖ મગફળી પાકને લોહ તત્વની ઉણપના કારણે પાન પીળા દેખાય તો ૧૦૦ ગ્રામ હીરાકશી તથા ૧૦ ગ્રામ લીંબુના ફૂલને ૧૦ લીટર પાણીમાં ઓગાળીને છંટકાવ કરવો. ❖ મગફળીમાં ચુસીયા, મોલોમશી, થ્રીપ્સ ના નિયંત્રણ માટે ફોસ્ફામીડોન ૩ મીલી દવા અથવા ડાયમીથોએટ ૧૫ મીલી દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. મોલોમશી સાથે દાળીયા જીવાત જોવા મળે તો દાવાનો છંટકાવ મુલતવી રાખવો. ❖ સતત ભેજવાળા અને વાદળછાયું વાતાવરણમાં પાકમાં પાનના ટપકાના રોગના આગોતરા નિયંત્રણ માટે હેકાકોનાઝોલ ૧૫ મીલી અથવા ટેબ્યુંકોનાઝોલ ૧૦ મીલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટકાવ કરવો. ❖ મગફળીના ઘૈણના નિયંત્રણ જમીનમાં પુરતો ભેજ હોય ત્યારે ફોરેટ ૧૦ જી ૨૫ કી.ગ્રા./હે. અથવા કલોરોપાયરીફોસ હેકટરે ૪ લીટર પીયત સાથે આપવી અથવા પીયતની સગવડ ન હોય તો નોઝલ કાઢી ૧૦ લીટર પાણીમાં ૨૫ મીલી દવા ભેળવી મગફળીના મુળ પાસે પડે અને જમીનમાં ઉત્તરે તે રીતે રેડવાથી નિયંત્રણ મળે છે.
કપાસ	ચાપવા અવસ્થા / ફૂલ અવસ્થા	ચુસીયા જીવાત	<ul style="list-style-type: none"> ❖ કપાસના પાકમાં થ્રીપ્સ અને લીલા તડતડીયાના નિયંત્રણ માટે ૩.૦ મીલી ફોસ્ફામીડોન અથવા ૧૫ મીલી ડાયમીથોએટ અથવા ૧૫ મીલી મીથાઈલ-ઓ-ડેમેટોન દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ કપાસનો પાક ત્રણ માસનો થાય ત્યારે પુર્ણ ખાતરનો હસ્તો હેકટરે ૧૬૬.૬ કી.ગ્રા. ચુરીયા અથવા ૪૦૦ કી.ગ્રા અમોનીયમ સલ્ફેટ આપવું.
સોયાબીન	વૃદ્ધી	ગુજ. સોયા.- ૧,૨,૩.	<ul style="list-style-type: none"> ❖ સોયાબીનનું વાવેતર કર્યા બાદ ૩૦ દિવસે રીલે અંતરપાક તરીકે બે હાર પછી એક હાર એરેંડાનું વાવેતર કરવું જેથી વધુ ઉત્પાદન મેળવી શકાય છે.
તલ	પાન કથીરી	પૂર્વ -૧, ગુજ.	<ul style="list-style-type: none"> ❖ પાકમાં કથીરીના નિયંત્રણ માટે ડાયકોફોલ ૨૦ મીલી અથવા

		તલ- ૧,૨.	દ્રાવ્ય ગંધક ૧૭ ગ્રામ ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો જરૂર જણાયતો બીજો છંટકાવ ૧૫ દિવસ બાદ કરવો.
એરંડા	વાવેતર / ઉગાવો	ગુ.એ.હા.- ૧ અને ગુ.એ.હા.- ૪, ૫, ૬, ૭	<ul style="list-style-type: none"> ❖ પાકમાં વાવેતર સમયે પાચામાં ડીએપી ૮૫.૨ ક્રીલો અને ૬૧.૪ ક્રીલો યુરીયા/હે આપવું. ❖ એરંડા વાવતા પહેલા બીયારણને થાયરમ, કેપ્ટન અને મેન્કોઝેબ દવા ૧ કી.ગ્રામ બીજ દીઠ ૩.૦૦ ગ્રામ મુજબ પટ આપવો. ❖ ખેત મજુરોની અછતની પરીસ્થીતીમાં પાક વાવ્યા બાદ નિંદામણ નાશક દવા ટ્રાઇફ્લુરાલીન ૧.૦ કી.ગ્રા. અથવા પેન્ડીમીથેલીન ૧.૦ કી.ગ્રા. સકીય તત્વ રૂપે છંટકાવ કરવો.
તુવેર	વાવેતર / ઉગાવો	ગુ.તુ.- ૧૦૦, બીડીએન-૨	<ul style="list-style-type: none"> ❖ તુવેર વાવણી માટે વધુ ઉત્પાદન આપતીવિશ્વાષપાત્ર સર્ટિફિકેડ બીયારણ જ ખરીદવું. ❖ તુવેરવાવતા પહેલા બીયારણને થાયરમ, કેપ્ટન અને મેન્કોઝેબ દવા ૧ કી.ગ્રામ બીજ દીઠ ૩.૦૦ ગ્રામ મુજબ પટ આપવો.
મગ	વૃદ્ધી	ચુસીયા જીવાત	<ul style="list-style-type: none"> ❖ પાકમાં ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે, ૧૦ મી.લી. ડાયમીથોયેટ અથવા મીથાઈલ-ઓ-ડેમેટોન દવા ૧૦ લીટર પાણીમાં ભેળવી જરૂરીયાત મુજબ છંટકાવ કરવો.
અડદ	વૃદ્ધી	ચુસીયા જીવાત	<ul style="list-style-type: none"> ❖ પાકમાં ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે, ૧૦ મી.લી. ડાયમીથોયેટ અથવા મીથાઈલ-ઓ-ડેમેટોન દવા ૧૦ લીટર પાણીમાં ભેળવી જરૂરીયાત મુજબ છંટકાવ કરવો.
શાકભાજી	વૃદ્ધી/કુલ/ ફળ	-	<ul style="list-style-type: none"> ❖ વેલાવાળા શાકભાજીના પાકમાં પાન કોરીયાનો ઉપદ્રવ ટાળવા માટે મુખ્ય પાકની ફરતે દિવેલા તથા ગલગોટાનું વાવેતર કરવાથી તેનું નિયંત્રણ કરી શકાય છે જો ઉપદ્રવ વધુ જોવા મળે તો ૧૦ મીલી મીથાઈલ ઓ ડેમેટોન (૨૫ ઈસી) અથવા ૧૦ મીલી ડાયમીથોયેટ (૩૦ ઈસી) ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો ત્યાર બાદ પછીથી બીજો છંટકાવ ૧૫ દિવસ પછી કરવો. ❖ શાકભાજીના પાકમાં ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીનું મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦મીલી ૧૦ લી. પાણીમાં ભેળવી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટામ્બાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦લી. અથવા દ્રાવ્યઝોક્સ ૪૦ ટકા ૨૦ મીલી/૧૦ લી.પાણીમાં મીશ્ર કરી છંટકાવ કરવો.

			<ul style="list-style-type: none"> ❖ મરચીના પાકમાં શ્રીપસના નિયંત્રણ માટે ૨૦મીલી પ્રોફેનોફ્રોસ અને લિગ્બોળીનું તેલ ૫૦ મીલી ૧૦ લીટર પાણી મિશ્રણ કરી છાંટવી. ❖ શાકભાજુનાપાકની રોપણી બાદ ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે દાણાદાર ફોરેટ ૧૦ જી છોડ દીઠ ૨ ગ્રામ મુજબ આપીને પીયત આપવું. ❖ મરચીના પાકમાં વાઈરસના કારણે પીળા થઈ કુકડાઈ ગયેલ છોડ ઉપાડી જમીનમાં દાટી નાશ કરવો અને ૫ મીલી ઈમીડાકલોપ્રાઇડ ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ રીંગાણ અને ટમેટાના ફળ અને અને ડોકાની ઈયળોનો ઉપદ્રવ જણાયતો તોનુકશાન પામેલ ફળ અને ડોકાને તોડીને જમીનમાં દાટી દેવા અને રીનાક્ષીપાયર દવા ૩મીલીદવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ રીંગાણ અને ટમેટાના પાકમા ફળમાખીના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા પગ્રામ દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ શાકભાજુ ને સમયસર ઉતારી ગ્રેડીંગ કરી બજારમાં મુકવા.
પશુપાલન	-	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો અને પશુઓના રહેઠાણ ચોખમા રાખવા. ❖ માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ પશુઓને યોગ્ય આહારમાંખનિજ્યુક્ત મિશ્રણ સાથે સંગૃહિત ચારા આપો. ચારા ને સલામત સ્થળે સંગૃહિત કરવો જોઈએ. ❖ તીવ્ર વરસાદ અને પૂરથી પ્રાણીઓને એકસાથે રાખવામાં આવે છે. ❖ ચારા માટે ચાફ કર્ટર વાપરો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાજ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરીક પાવડરથી બરાબર સાફ કરવા.

JUNAGADH

Phone: 0285-2670139

Date: - 31-08-2018

Significant weather for the preceding week

Rain fall (mm)	000.0
Maximum temperature °C	28.7 – 31.2
Minimum temperature °C	23.9 – 24.6
RH % (Morning/Afternoon)	92 – 64
Wind speed (Kmph)	05.0 – 05.9
Cloud cover	2 - 8

(Period from 01 to 05 September 2018)

Weather forecast for Junagadh Dist. until 08:30 hrs of 05-09-2018

Parameters	Day-1 01-09-18	Day-2 02-09-18	Day-3 03-09-18	Day-4 04-09-18	Day-5 05-09-18
Rain fall (mm)	04	00	00	00	00
Maximum temperature °C	29	29	29	28	28
Minimum temperature °C	26	26	26	26	26
Cloud cover (Okta)	8	8	7	6	3
RH I % (Morning)	87	85	82	82	84
RH II % (After noon)	75	73	65	65	65
Wind speed (Km/h)	017	019	016	015	012
Wind direction(Deg)	253	253	249	247	248

Agro meteorological Advisories

General Advice	Rain fall	There is forecasting of light rainfall tomorrow and no rainfall during next four days.
	Temperature	Maximum and minimum temperature likely to be in range of 28 - 29°C and 26°C, respectively.

	RH (%)	The Relative Humidity is likely to be in the range of 87 – 65 %.	
	Wind Direction and Wind Speed	Direction and speed of wind is likely to W and SW and 12 - 19 km/h, respectively.	
Crop	Variety/ Pest/Insect	Stage	Agro met Advisories
			<ul style="list-style-type: none"> ○ The sky will remain variable cloudy and forecasting of light rainfall tomorrow and no rainfall during next four days. ○ Apply irrigation to the crop at different growth stage due to the shortage of rainfall.
Kharif Groundnut	Stem rot / Sucking pest	Flowering / Pegging stage	<ul style="list-style-type: none"> ➢ Interculturing and hand weeding should be carried out in groundnut. ➢ Under limited water condition groundnut should be irrigated during pegging stage without fail. ➢ For the control of stem rot, furrow application of 2.5 kg/ha <i>Trichoderma viride</i>, 500 kg castor cake OR vermicomposting. ➢ In Ground nut, spray 0.1% FeSO₄ @ 100 gram with 10 gram citric acid solution to correct chlorosis. ➢ For the control of sucking pest (aphid, thrips jassid), spray 3 ml Phosphamidon OR 15 ml Dimethoate in 10 litre of water. For ladybird beetle is observed with aphid then postpone the spraying operation. ➢ For the control of rust and tikka, spray fungicides mencozeb 26 gm mix in 10 litre of water at after sowing of 35 days. ➢ Due to continuous humid & cloudy weather spray Tebuconazole 10 ml/10 lit of to prevent leaf spot infestation. ➢ For the control of white grub, soil application

			of phorate 25 kg/ha OR 4 liter/ha choloropayriphos apply with irrigation OR Drenching of 25 ml choloropayriphos in 10 liter of water near the base of the plant.
Cotton	Sucking pest	Square initiation / Flowering	<ul style="list-style-type: none"> ➤ For the control of Thrips and Jassid, spray 3 ml phosphamidon OR 15 ml dimethoate OR 15 ml methyl-O-demeton mix in 10 liter of water. ➤ When the cotton crop is three months, apply the supplementary fertilizer dose is 166.6 Urea kg/ha or 400 kg/ha Ammonium sulphate.
Soyabean	Gujarat Soya bean – 1, 2, 3.	Vegetative growth	<ul style="list-style-type: none"> ➤ 30 days after sowing of soybean relay intercropping with castor planting can be obtained more production. ➤ For the control of prodenia, spray 25 ml chloropyariphos OR 10 ml methomile OR 5 ml DDVP in 10 litre of water. OR use of integrated pest management like a pheromone trap, <i>verticillium locati</i> fungi etc.
Seasame	Purva-1, Guj. Til – 1, 2.	Vegetative growth / Flowering	<ul style="list-style-type: none"> ➤ Due to cloudy weather condition incidence of mite, for the control of spray dicophol 20 ml OR ethion 20 ml in 10 litre of water.
Castor	GAUCH-1 and GCH- 4,5,6,7	Sowing / germination	<ul style="list-style-type: none"> ➤ In unirrigated castor apply 65.2 kg DAP/ha & 61.4 kg urea/ha at the time of sowing. ➤ To give seed treatment with Thiram, Captan and Mencozeb @ 3 grams per kg of seeds before sowing. ➤ After sowing of castor before emergence of seed spray weedicides 1.0 kg/ha trifluralin OR 1.0 kg/ha pendimethalin.
Pigeon pea	GT- 100, BDN-2	Sowing/ germination	<ul style="list-style-type: none"> ➤ To procure good quality certified seeds of Pigeon pea. ➤ To give seed treatment with Thiram, Captan

			and Mencozeb @ 3 grams per kg of seeds before sowing.
Green gram	Sucking pest	Vegetative growth	➤ For the control of sucking pest, spray 10 ml Dimethoate OR Methyl-O-Demeton in 10 litre of water.
Black gram	Sucking pest	Vegetative growth	➤ For the control of sucking pest, spray 10 ml Dimethoate OR Methyl-O-Demeton in 10 litre of water.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<p>➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water.</p> <p>➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.</p> <p>➤ For the control of thrips in chilli, spray 20 ml profenophos in 10 litres of water.</p> <p>➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.</p> <p>➤ Rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of 5 mi Imidacloprid per 10 litre of water is advised for control the vector, when sky</p>

			<p>remains clear.</p> <ul style="list-style-type: none"> ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops constant monitoring of borers by using pheromone traps 4-6 per acre is advised, for the control of fruit borer, spray reenaxipayar @ 3.0ml in 10 lit of water. ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetemaphrid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ In severe rain and flood to keep untether the animals. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.

PORBANDAR

Phone: 0285-2670139

Date: - 31-08-2018

(Period from 01 to 05 September 2018)

Weather forecast for Porbandar Dist. until 08:30 hrs of 05-09-2018

Parameters	Day-1 01-09-18	Day-2 02-09-18	Day-3 03-09-18	Day-4 04-09-18	Day-5 05-09-18
Rain fall (mm)	00	03	00	00	00
Maximum temperature °C	29	29	29	28	28
Minimum temperature °C	25	25	25	25	25
Cloud cover (Okta)	8	8	8	6	3
RH I % (Morning)	83	83	76	77	79
RH II % (After noon)	75	73	67	67	65
Wind speed (Km/h)	023	023	021	023	017
Wind direction(Deg)	253	248	249	249	244

Agro meteorological Advisories

General Advice	Rain fall		There is forecasting of light rainfall next two days and no rainfall during 3 to 5 September.
	Temperature		Maximum and minimum temperature likely to be in range of 28 - 29°C and 25°C, respectively.
	RH (%)		The Relative Humidity is likely to be in the range of 83 – 65 %.
	Wind Direction and Wind Speed		Direction and speed of wind is likely to be W and SW and 17 – 23 km/h, respectively.
Crop	Variety/ Pest / Insect	Stage	Agro met Advisories
			• The sky will remain variable cloudy and forecasting of light rainfall next two days and no rainfall during 3 to 5 September.

			<ul style="list-style-type: none"> • Apply irrigation to the crop at different growth stage due to the shortage of rainfall.
Kharif Groundnut	Stem rot / Sucking pest	Flowering / Pegging	<ul style="list-style-type: none"> ➤ Under limited water condition groundnut should be irrigated during pegging stage without fail. ➤ For the control of stem rot, furrow application of 2.5 kg/ha <i>Trichoderma viridi</i>, 500 kg castor cake OR vermicomposting. ➤ In Ground nut, spray 0.1% FeSO₄ @ 100 gram with 10 gram citric acid solution to correct chlorosis. ➤ For the control of sucking pest (aphid, thrips jassid), spray 3 ml Phosphamidon OR 15 ml Dimethoate in 10 litre of water. For ladybird beetle is observed with aphid then postpone the spraying operation. ➤ For the control of rust and tikka, spray fungicides mencozeb 26 gm mix in 10 litre of water at after sowing of 35 days. ➤ Due to continuous humid & cloudy weather spray Tebuconazole 10 ml/10 lit of to prevent leaf spot infestation. ➤ For the control of white grub, soil application of phorate 25 kg/ha OR 4 liter/ha choloropayriphos apply with irrigation OR Drenching of 25 ml choloropayriphos in 10 liter of water near the base of the plant.
Cotton	Sucking pest	Square initiation /	<ul style="list-style-type: none"> ➤ For the control of Thrips and Jassid, spray 3 ml phosphamidon OR 15 ml dimethoate OR 15 ml

		Flowering	<p>methyl-O-demeton mix in 10 liter of water.</p> <ul style="list-style-type: none"> ➤ When the cotton crop is three months, apply the supplementary fertilizer dose is 166.6 Urea kg/ha or 400 kg/ha Ammonium sulphate.
Soyabean	Gujarat Soya bean – 1, 2, 3.	Vegetative growth	<ul style="list-style-type: none"> ➤ 30 days after sowing of soybean relay intercropping with castor planting can be obtained more production. ➤ For the control of prodenia, spray 25 ml chloropyariphos OR 10 ml methomile OR 5 ml DDVP in 10 litre of water. OR use of integrated pest management like a pheromone trap, <i>verticillium locati</i> fungi etc.
Seasame	Purva-1, Guj. Til – 1, 2.	Vegetative growth / Flowering	<ul style="list-style-type: none"> ➤ Due to cloudy weather condition incidence of mite, for the control of spray dicophol 20 ml OR ethion 20 ml in 10 litre of water.
Castor	GAUCH-1 and GCH- 4,5,6,7	Sowing / Germination	<ul style="list-style-type: none"> ➤ In unirrigated castor apply 65.2 kg DAP/ha & 61.4 kg urea/ha at the time of sowing. ➤ To give seed treatment with Thiram, Captan and Mencozeb @ 3 grams per kg of seeds before sowing. ➤ After sowing of castor before emergence of seed spray weedicides 1.0 kg/ha trifluralin OR 1.0 kg/ha pendimethalin.
Pigeon pea	GT- 100, BDN-2	Sowing	<ul style="list-style-type: none"> ➤ To procure good quality certified seeds of Pigeon pea. ➤ To give seed treatment with Thiram, Captan and Mencozeb @ 3 grams per kg of seeds before sowing.
Green gram	Sucking pest	Vegetative growth	<ul style="list-style-type: none"> ➤ For the control of sucking pest, spray 10 ml Dimethoate OR Methyl-O-Demeton in 10 litre of water.

Black gram	Sucking pest	Vegetative growth	<ul style="list-style-type: none"> ➤ For the control of sucking pest, spray 10 ml Dimethoate OR Methyl-O-Demeton in 10 litre of water.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water. ➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ For the control of thrips in chilli, spray 20 ml profenophos in 10 litres of water. ➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ Rouging of viralinfested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of 5 ml Imidacloprid per 10 litre of water is advised for control the vector, when sky remains clear. ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops constant monitoring of borers by using pheromone traps 4-6 per acre is advised, for the control of

			<p>fruit borer, spray reenaxipayar @ 3.0ml in 10 lit of water.</p> <ul style="list-style-type: none"> ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetemaphrid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food and keep the animals clean shed. ➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Give stored fodder with mineral mixture. Fodder should be stored at safe place. ➤ In severe rain and flood to keep untether animals. ➤ Feed silage & hay material along with concentrate feed. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.