

હવામાન આગાહી આધારીત કૃષિ સલાહ બુલેટીન  
દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ  
તા. ૦૨-૦૧-૨૦૧૬ થી તા. ૦૬-૦૧-૨૦૧૬

(ગ્રામીણ કૃષિ મૌસમ સેવા, જુનાગઢ કૃષિ યુનિવર્સીટી, જી.કૃ.યુ.)

હવામાન આગાહી આધારીત બુલેટીન : ૨૦૧૫

પાછલા અઠવાડીયાનું જુનાગઢનું હવામાન

હવામાન પરીબળો	દક્ષિણ સૌરાષ્ટ્ર કૃષિ આબોહવાકીય વિસ્તાર
વરસાદ (મીમી)	0.00
મહત્તમ તાપમાન (સે.ગે.)	36.4
લઘુતમ તાપમાન (સે.ગ્રે.)	14.7
ભેજ(%) (સવાર/બપોરબાદ)	75-34
પવનની ગતિ (કિમી/કલાક)	2.5

**પત્રક-બ : જિલ્લાવાર હવામાન આગાહી ( તા. ૦૬-૦૧-૧૬ સુધી )**

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહિતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જૂનાગઢ કૃષિ યુનિવર્સિટી, જૂનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના

**૧. જૂનાગઢ:-** જીલ્લામા આગામી તારીખ ૦૨/૦૧/૨૦૧૬ થી ૦૬/૦૧/૨૦૧૬ દરમ્યાન આકાશ મુખ્યત્વે ચોખ્ખુ રહેશે, વરસાદ પડવાની શક્યતા રહેશે નહિ અને પવનની ઝડપ ૧૩ થી ૧૯ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્તમ તાપમાન ૩૨ થી ૩૪ ડીગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્તમ તાપમાન ૧૬ થી ૨૦ ડીગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

**૨. પોરબંદર:-** જીલ્લામા આગામી તારીખ ૦૨/૦૧/૨૦૧૬ થી ૦૬/૦૧/૨૦૧૬ દરમ્યાન આકાશ મુખ્યત્વે ચોખ્ખુ રહેશે, વરસાદ પડવાની શક્યતા રહેશે નહિ અને પવનની ઝડપ ૧૧ થી ૧૮ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્તમ તાપમાન ૩૩ થી ૩૫ ડીગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્તમ તાપમાન ૧૪ થી ૧૫ ડીગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

**જૂનાગઢ**

હવામાન પરીબળો / તારીખ	દિવસ-૧ ૦૨.૦૧.૧૬	દિવસ-૨ ૦૩.૦૧.૧૬	દિવસ-૩ ૦૪.૦૧.૧૬	દિવસ-૪ ૦૫.૦૧.૧૬	દિવસ-૫ ૦૬.૦૧.૧૬
વરસાદ (મી.મી)	૦	૦	૦	૦	૦
મહત્તમ તાપમાન (સે.ગ્રે)	૩૪	૩૪	૩૩	૩૩	૩૨
લઘુત્તમ તાપમાન (સે.ગ્રે)	૨૦	૧૯	૧૭	૧૬	૧૬
વાદળની સ્થિતિ	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	મોટેભાગે ચોખ્ખું
સવારના ભેજના ટકા	૪૬	૪૯	૫૪	૫૩	૫૪
બપોર બાદ ભેજના ટકા	૪૧	૩૯	૧૩	૨૦	૧૭
પવનની ગતિ (કિમી/કલાક)	૦૧૫	૦૧૬	૦૧૩	૦૨૦	૦૧૭
પવનની દિશા (ડીગ્રી)	૨૬	૨૬	૪૯	૧૧	૬૦

**પોરબંદર**

વરસાદ (મી.મી)	૦	૦	૦	૦	૦
મહત્તમ તાપમાન (સે.ગ્રે)	૩૫	૩૫	૩૪	૩૪	૩૩
લઘુત્તમ તાપમાન (સે.ગ્રે)	૧૫	૧૫	૧૪	૧૪	૧૫
વાદળની સ્થિતિ	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	ચોખ્ખું	મોટેભાગે ચોખ્ખું
સવારના ભેજના ટકા	૪૨	૪૮	૫૪	૫૧	૫૩
બપોર બાદ ભેજના ટકા	૩૭	૩૭	૪૨	૪૦	૩૫
પવનની ગતિ (કિમી/કલાક)	૦૧૪	૦૧૭	૦૧૧	૦૧૮	૦૧૭
પવનની દિશા (ડીગ્રી)	૨૬	૨૫	૧૬	૧૨	૨૪

## દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકિય વિભાગ

પાકનું નામ	રોગ/જીવાત/ જાત	કૃષિ સલાહ
ઘઉં	લોક-૧, જી. ડબલ્યુ-૩૬૬, જી. ડબલ્યુ-૧૯૦, જી. ડબલ્યુ-૪૯૬	<ul style="list-style-type: none"> <li>❖ પાકને મુકુટમૂળ અવસ્થાએ ૧૩૦ કિલો યુરીયા/હેક. અને પિયત આપવું.</li> <li>❖ વાવેતર બાદ ૩૦ થી ૩૫ દિવસે નિદામણ નિયંત્રણ માટે ૨૦ મીલી ૨,૪ ડી. અથવા મેટસલ્ફ્યુરન મિથાઈલ ૦.૮ ગ્રામ ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો.</li> </ul>
ધાણા	ગુ. ધાણા- ૧ & ૨	<ul style="list-style-type: none"> <li>❖ ધાણામા આવતી થ્રીપ્સ, મોલોમસી ના નિયંત્રણ માટે ડાયમિથોએટ ૧૦ મીલી અથવા ટ્રાયઝોફોસ ૧૦ મીલી પ્રવાહીને ૧૦ લીટર પાણીમાં મિશ્રણ કરી ૧૫ દિવસના અંતરે છંટકાવ કરવા અથવા કુગજન્ય જૈવિક જંતુનાશક દવા “વર્ટીસીલીયમ લેકાની” અને “બીવેરીયા બાઝીયાના” નામની કુગનો પાવડર ૧૦ લીટર પાણીમાં ૪૦ ગ્રામ પ્રમાણે ભેળવી છંટકાવ કરવાથી અસરકારક પરીણામ મળે છે.</li> </ul>
લસણ	ગુ. લસણ-૩ & ૪	<ul style="list-style-type: none"> <li>❖ ઉગસકુનોરોગ સફેદ ફૂગના કારણે હોવાથી ટેબયકોનાઝોલ ૧૦ મીલી દવાને ૧૦ લીટર પાણીમાં મિશ્રણ કરીને ડ્રેનર્યાંગ કરવું.</li> <li>❖ લસણમાં થ્રીપ્સના નિયંત્રણ માટે ફોસ્ફામીડોન ૩ મીલી અથવા ડાયમિથોએટ ૧૦ મીલી અથવા મીથાઈલ-ઓ-ડીમેટોન ૧૦ મીલી ના પ્રવાહીને ૧૦ લીટર પાણીમાં મિશ્રણ કરી ૧૦ થી ૧૨ દિવસના અંતરે છંટકાવ કરવા.</li> </ul>
જીરું	ગુ. જીરું- ૨ & ૪	<ul style="list-style-type: none"> <li>❖ જીરૂના પાકમાં થ્રીપ્સ અને મોલોમસીના નિયંત્રણ માટે થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્રણ કરી છાંટવી. તેમજ ચરમીનો રોગ ના થાય તે માટે ૨૫ ગ્રામ ડાયથેન એમ-૪૫ ૧૦ લીટર પાણીમાં મિશ્રણ કરી ૧૦ દિવસના અંતરે છંટકાવ કરવા.</li> </ul>
ચણા	જી.સી.-૨ & ૩, ICCC-૪	<ul style="list-style-type: none"> <li>❖ ચણાના પાકમાં લીલી અને લશ્કરી ઈયળના નિયંત્રણ માટે ૬ ગ્રામ એમાકેટીન બેન્ઝોએટ અથવા ૨૦ મીલી ક્વીનાલફોસ ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવાથી સારૂ પરીણામ મળે છે.</li> </ul>
ડુંગળી	નાસીક-૫૩, તળાજા- ૨૬, જૂનાગઢ લોકલ	<ul style="list-style-type: none"> <li>❖ ડુંગળીમાં જાંબલી રંગના ધાબાના રોગના નિયંત્રણ માટે ૨૫ ગ્રામ મેન્કોઝેબ અથવા ૧૫ મીલી ક્વિનાલફોસ ૧૦ લીટર પાણીમાં મિશ્ર કરી ૧૦ દિવસના અંતરે છંટકાવ કરવો.</li> <li>❖ ડુંગળીમાં થ્રીપ્સના નિયંત્રણ માટે ફોસ્ફામીડોન ૩ મીલી અથવા ડાયમિથોએટ ૧૦ મીલી અથવા મીથાઈલ-ઓ-ડીમેટોન ૧૦ મીલી ના પ્રવાહીને ૧૦ લીટર પાણીમાં મિશ્રણ કરી ૧૦ થી ૧૨ દિવસના</li> </ul>

		અંતરે છંટકાવ કરવા.
કપાસ	-	❖ કપાસની ગુલાબી ઈયળના નિયંત્રણ માટે જે કપાસના પાકમાં ગુલાબી ઈયળનો ઉપદ્રવ હોય તયાં બીજો ફાલ લેવો નહિ અને કપાસની વીણી કરીને કપાસ કાઢીને સળગાવી દેવો.
તુવેર	-	❖ પાકમાં શિંગો કોરી ખાનાર ઈયળના નિયંત્રણ માટે ૪૫ એસ.એલ સ્પિનોસાડ ૩ મીલી અથવા ક્લોરાન્દ્રાનિલીપોલ ૩ મીલી અથવા ફ્લુબેન્ડીયામાઈડ ૨ મીલી દવા ને ૧૦ લીટર પાણીમાં ભેળવી બે છંટકાવ કરવા.
દિવેલા	-	❖ એરંડાના પાકમાં ઘોડીયા ઈયળના નિયંત્રણ માટે ૨૦ મીલી ક્વિનાલફોસ અથવા ૫ મીલી ડી.ડી.વી.પી. અથવા ૪૦ ગ્રામ કાર્બારીલ દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો.
શાકભાજી	-	❖ ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે લીબોળીનુ તેલ ૫૦ મીલી અથવા લીબોળીનુ મીજનુ ૫ ટકા અર્કનુ દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં ભેળવી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટામ્રાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦લી. અથવા દ્રાયઝોફોસ ૪૦ ટકા ૨૦ મીલી/ ૧૦ લી. પાણીમાં મીશ્ર કરી છંટકાવ કરવો. ❖ ફળ અને ડોકાની ઈયળનો ઉપદ્રવ વધુ હોયતો નુક્શાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં ડાટી દેવા અથવા ૩ મીલી રીનાક્ષીપાયર દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ પાકમાં કથીરીના નિયંત્રણ માટે પ્રોપરગાઈટ ૧૦ મીલી/૧૦ લીટર પાણી મિશ્રણ કરી છાંટવી.
પશુ પાલન	-	❖ પશુઓને યોગ્ય આહાર આપવો. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાઝ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ પશુઓને ખરવા-મોવાસા રોગની રસીનો પ્રથમ ડોઝ ૪ માસની ઉમરે, બીજો ડોઝ ૬ માસની ઉમરે અને ત્યાર બાદ દર ૬ માસે આપવો. ❖ શિયાળા ત્રૂતુમાં અતિશય ઠંડા સીધા પવનથી રક્ષણ માટે પશુઓને શેડમાં બાંધવા અને કંતાન કોથળાનો ઉપયોગ કરવો.

## JUNAGADH

Phone: 0285-2670139

Date:- 01-01-2016

(Period from 02 to 06 Jan. -2016)

### Significant weather for the preceding week

Rain fall (mm)	000.0
Maximum temperature °C	35.8-37.2
Minimum temperature °C	14.1-14.5
RH % (Morning/Afternoon)	75-34
Wind speed (Kmph)	2.1-3.1
Cloud cover	0-2

### Weather forecast for Junagadh Dist. until 08:30 hrs of 06-01-2016

Parameters	Day-1 02-01-16	Day-2 03-01-16	Day-3 04-01-16	Day-4 05-01-16	Day-5 06-01-16
Rain fall (mm)	0	0	0	0	0
Maximum temperature °C	34	34	33	33	32
Minimum temperature °C	20	19	17	16	16
Cloud cover (Okta)	0	0	0	0	1
RH I % (Morning)	46	49	54	53	54
RH II % (After noon)	41	39	13	20	17
Wind speed (Kmph)	015	016	013	020	017
Wind direction(Deg)	26	26	49	11	60

### Agro meteorological Advisories

<b>General Advice</b>	<b>Rain fall</b> <b>Temperature</b> <b>RH (%)</b> <b>Wind Direction and Speed</b>	<p>There is no forecasting of rainfall during next five days.</p> <p>Maximum and minimum temperature likely to be in range of 32-34°C and 16-20 °C, respectively.</p> <p>The Relative Humidity is likely to be in the range of 54-39 %.</p> <p>Direction and speed of wind is likely to be NE and 13-20 km/h, respectively.</p>	
Crop	Stage	Insect/Pest/ Variety	Agromet Advisories
<b>Wheat</b>	<b>CRI</b>	<b>GW-366, Lok-1, GW-190</b>	<ul style="list-style-type: none"> <li>➤ Apply 130kg Urea as top dressing per ha. and give irrigation to the crop.</li> <li>➤ At 30-35 days after sowing spray 20 ml 2-4-D ethyle ester or Metasulfuron 0.8 gm/10 liters of water to control broad leafy weeds in wheat crop.</li> </ul>
<b>Coriander</b>	<b>Vegetative</b>	<b>Guj. Cor. -1 &amp; 2</b> <b>Thrips, Aphid</b>	<ul style="list-style-type: none"> <li>➤ Due to cold weather condition for the control of thrips &amp; aphid, spray of Dimethoate 10 ml or Trizophos 10 ml in 10 liter of water at 15 days interval.</li> </ul>

<b>Garlic</b>	<b>Vegetative</b>	<b>Guj. Garlic-3 &amp;4</b>	➤ Yellowing of leaves due to stem (clove) rot, Drenching of Tebuconazole 10 ml/10 litre of water.
		<b>Thrips</b>	➤ For the control of thrips, spray Phosphamidon 3.0 ml or Dimethoate 10 ml or Methay-O-Demeton 10 ml in 10 liter of water at 10-12 days interval.
<b>Cumin</b>	<b>Sowing</b>	<b>Guj. Cumin-2 &amp;4</b>	➤ For the control of aphid and thrips, spray Thiomethoxam 5 gm/ 10 liters of water. ➤ Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.
<b>Chickpea</b>	<b>Flowering</b>	<b>GC-2 &amp;3, ICC-4</b>	➤ For the control of Prodenia ane Heliothis at ETL, spray 20 ml quinalphos in 10 litre of water.
<b>Onion</b>	<b>Sowing for seedling/ Transplanting</b>	<b>Nasik-53, JND Local, Talaja Red, Pusa Red</b>	➤ For the control of purple blotch, spray 25 gm Mencozeb and 15 ml hexaconazole in 10 litres of water at 10-15 days interval. ➤ For the control of thrips, spray Phosphamidon 3.0 ml or Dimethoate 10 ml or Methay-O-Demeton 10 ml in 10 liter of water at 10-12 days interval.
<b>Cotton</b>	<b>Ball formation</b>	<b>Bg-II GCH-6, 8, Mallika, Ajit, Vikram-5, Uttam, etc...</b>	➤ Under present weather condition heavy attack of <b>pink boll worm</b> observed in irrigated cotton, therefore farmers are advice not to take second flush of flowering and uproot the crop after picking.
<b>Pigeon pea</b>	<b>Pod formation</b>	<b>BDN-2, GTH-1</b> <b>Pod borer</b>	➤ For the control of pod borer, spray spinosad 5 ml OR clorantirinilipol 3 ml OR flubendimide 2 ml in 10 liter of water.
<b>Castor</b>	<b>Capsule formation</b>		➤ Apply sulphur 20 kg/ha (Gypsum 154 kg/ha) to obtain greater production and economic returns. ➤ For the control of capsule borer, spray quinalphos 20 ml OR 5 ml D.D.V.P OR 40 gm Carbaryl in 10 litre of water.
<b>Vegetable crops</b>	<b>Flowering/ fruiting</b>	<b>Thrips and White Fly</b>	➤ For the control of thrips and white fly, spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray).
		<b>Shoot and fruit borer</b>	➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ Collect damaged fruits and buried deep in soil and spray 3 ml renaxypyre in 10 litre of water.
<b>Livestock</b>	-	-	➤ Do vaccination in animal for the prevention of FMD and other diseases.

			<ul style="list-style-type: none"><li>➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water.</li><li>➤ Protection of animal due to the cold winter season cattle transfer to the shed.</li></ul>
--	--	--	--

## PORBANDAR

Phone : 0285-2670139

Date : 01-01-2016

(Period from 02 to 06 Jan. -2016)

**Weather forecast for Porbandar Dist. until 08:30 hrs of 06-01-2016**

Parameters	Day-1 02-01-16	Day-2 03-01-16	Day-3 04-01-16	Day-4 05-01-16	Day-5 06-01-16
Rain fall (mm)	0	0	0	0	0
Maximum temperature °C	35	35	34	34	33
Minimum temperature °C	15	15	14	14	15
Cloud cover (Okta)	0	0	0	0	1
RH I % (Morning)	42	48	54	51	53
RH II % (After noon)	37	37	42	40	35
Wind speed (Kmph)	014	017	011	018	017
Wind direction(Deg)	26	25	16	12	24

### Agro meteorological Advisories

<b>General Advice</b>	<b>Rain fall</b> <b>Temperature</b> <b>RH (%)</b> <b>Wind Direction and Speed</b>	<p>There is no forecasting of rainfall during next five days.</p> <p>Maximum and minimum temperature likely to be in range of 33-35°C and 14-15 °C, respectively.</p> <p>The Relative Humidity is likely to be in the range of 54-35 %.</p> <p>Direction and speed of wind is likely to be NNE and 11-18 km/h, respectively.</p>	
Crop	Stage	Insect/Pest/ Variety	Agromet Advisories
<b>Wheat</b>	<b>CRI</b>	<b>GW-366, Lok-1, GW-190</b>	<ul style="list-style-type: none"> <li>➤ Apply 130kg Urea as top dressing per ha. and give irrigation to the crop.</li> <li>➤ At 30-35 days after sowing spray 20 ml 2-4-D ethyle easter or Metasulfuron 0.8 gm/10 liters of water to control broad leafy weeds in wheat crop.</li> </ul>
<b>Coriander</b>	<b>Vegetative</b>	<b>Guj. Cor. -1 &amp; 2</b> <b>Thrips, Aphid</b>	<ul style="list-style-type: none"> <li>➤ Due to cold weather condition for the control of thrips &amp; aphid, spray of Dimethoate 10 ml or Trizophos 10 ml in 10 liter of water at 15 days interval.</li> </ul>
<b>Garlic</b>	<b>Vegetative</b>	<b>Guj. Garlic-3 &amp; 4</b> <b>Thrips</b>	<ul style="list-style-type: none"> <li>➤ Yellowing of leaves due to stem (clove) rot, Drenching of Tebuconazole 10 ml/10 litre of water.</li> <li>➤ For the control of thrips, spray Phosphamidon 3.0 ml or Dimethoate 10 ml or Methay-O-Demeton 10 ml in 10 liter of water at 10-12 days interval.</li> </ul>



<b>Cumin</b>	<b>Sowing</b>	<b>Guj. Cumin-2 &amp;4</b>	<ul style="list-style-type: none"> <li>➤ For the control of aphid and thrips, spray Thiomethoxam 5 gm/ 10 liters of water.</li> <li>➤ Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</li> </ul>
<b>Chickpea</b>	<b>Flowering</b>	<b>GC-2 &amp;3, ICC-4</b>	<ul style="list-style-type: none"> <li>➤ For the control of Prodenia ane Heliothis at ETL, spray 20 ml quinalphos in 10 litre of water.</li> </ul>
<b>Onion</b>	<b>Sowing for seedling/ Transplanting</b>	<b>Nasik-53, JND Local, Talaja Red, Pusa Red</b>	<ul style="list-style-type: none"> <li>➤ For the control of purple blotch, spray 25 gm Mencozeb and 15 ml hexaconazole in 10 litres of water at 10-15 days interval.</li> <li>➤ For the control of thrips, spray Phosphamidon 3.0 ml or Dimethoate 10 ml or Methay-O-Demeton 10 ml in 10 liter of water at 10-12 days interval.</li> </ul>
<b>Cotton</b>	<b>Ball formation</b>	<b>Bg-II GCH-6, 8, Mallika, Ajit, Vikram-5, Uttam, etc...</b>	<ul style="list-style-type: none"> <li>➤ Under present weather condition heavy attack of <b>pink boll worm</b> observed in irrigated cotton, therefore farmers are advice not to take second flush of flowering and uproot the crop after picking.</li> </ul>
<b>Pigeon pea</b>	<b>Pod formation</b>	<b>BDN-2, GTH-1</b> <b>Pod borer</b>	<ul style="list-style-type: none"> <li>➤ For the control of pod borer, spray spinosad 5 ml OR clorantirinilipol 3 ml OR flubendimide 2 ml in 10 liter of water.</li> </ul>
<b>Castor</b>	<b>Capsule formation</b>		<ul style="list-style-type: none"> <li>➤ Apply sulphur 20 kg/ha (Gypsum 154 kg/ha) to obtain greater production and economic returns.</li> <li>➤ For the control of capsule borer, spray quinalphos 20 ml OR 5 ml D.D.V.P OR 40 gm Carbaryl in 10 litre of water.</li> </ul>
<b>Vegetable crops</b>	<b>Flowering/ fruiting</b>	<b>Thrips and White Fly</b>	<ul style="list-style-type: none"> <li>➤ For the control of thrips and white fly, spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray).</li> </ul>
		<b>Shoot and fruit borer</b>	<ul style="list-style-type: none"> <li>➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.</li> <li>➤ Collect damaged fruits and buried deep in soil and spray 3 ml renaxypyre in 10 litre of water.</li> </ul>
<b>Livestock</b>	-	-	<ul style="list-style-type: none"> <li>➤ Do vaccination in animal for the prevention of FMD and other diseases.</li> <li>➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water.</li> <li>➤ Protection of animal due to the cold winter season cattle transfer to the shed.</li> </ul>

