

હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીયવિભાગ

તા. ૦૩-૦૬-૨૦૧૭ થી તા. ૦૭-૦૬-૨૦૧૭

(ગ્રામીણ કૃષિ મૌસમ સેવા, જુનાગઢ કૃષિ યુનિવર્સિટી, જી.કૃ.યુ.)

હવામાન આગાહી આધારીત બુલેટીન : ૨૦૧૬-૧૭

પાછલા અઠવાડીયાનું જુનાગઢનું હવામાન

હવામાન પરીબળો	દક્ષિણ સૌરાષ્ટ્ર કૃષિ આબોહવાકીય વિસ્તાર
વરસાદ (મીમી)	૦૦૦.૦
મહત્તમ તાપમાન (°સે.)	૩૯.૯
લઘુત્તમ તાપમાન (°સે.)	૨૬.૦
ભેજ(%) (સવાર/બપોરબાદ)	૮૦ - ૪૬
પવનની ગતિ (કિમી/કલાક)	૧૩.૮

પત્રક-બ : જુલલાવાર હવામાન આગાહી (તા.૦૭-૦૬-૨૦૧૭ સુધી)

ભારત સરકારશ્રીના હવામાન વિભાગ તરફથી મળેલ હવામાન આગાહીની માહિતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા યોજના, જુનાગઢ કૃષિ યુનિવર્સિટી, જુનાગઢ તરફથી જણાવવામા આવે છે કે દક્ષિણ સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના

૧.જુનાગઢ:- જુલલામા આગામી તારીખ ૦૩/૦૬/૨૦૧૭ થી ૦૭/૦૬/૨૦૧૭ દરમ્યાન આકાશમુખ્યત્વે મોટેભાગે વાદળછાયુ રહેશે, આગામી પાંચ દિવસદરમ્યાન હળવો વરસાદ પડવાની શક્યતા છે. પવનની ઝડપ ૦૬ થી ૧૫ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્તમ તાપમાન ૩૯ થી ૪૦ ડીગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્તમ તાપમાન ૨૫ થી ૨૬ ડીગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

૨.પોરબંદર:- જુલલામા આગામી તારીખ ૦૩/૦૬/૨૦૧૭ થી ૦૭/૦૬/૨૦૧૭ દરમ્યાન આકાશમુખ્યત્વે મોટેભાગે વાદળછાયુ રહેશે, આગામી પાંચ દિવસદરમ્યાન વરસાદ પડવાની શક્યતા નથી. પવનની ઝડપ ૦૪ થી ૧૪ કીમી/કલાક રહેવાની શક્યતા રહેશે. આ સમયગાળામાં મહત્તમ તાપમાન ૩૫ ડીગ્રી સેન્ટીગ્રેડ રહેશે તેમજ લઘુત્તમ તાપમાન ૨૮ થી ૨૯ ડીગ્રી સેન્ટીગ્રેડ રહેવાની સંભાવના છે.

જુનાગઢ

હવામાન પરીબળો / તારીખ	દિવસ-૧ ૦૩-૦૬-૨૦૧૭	દિવસ-૨ ૦૪-૦૬-૨૦૧૭	દિવસ-૩ ૦૫-૦૬-૨૦૧૭	દિવસ-૪ ૦૬-૦૬-૨૦૧૭	દિવસ-૫ ૦૭-૦૬-૨૦૧૭
વરસાદ (મી.મી)	૦	૦	૦	૦	૩
મહત્તમ તાપમાન (સે.ગ્રે)	૩૯	૪૦	૩૯	૩૯	૩૯
લઘુત્તમ તાપમાન (સે.ગ્રે)	૨૫	૨૫	૨૬	૨૬	૨૬
વાદળની સ્થિતિ	મોટેભાગે વાદળછાયુ	મોટેભાગે વાદળછાયુ	અંશત વાદળછાયુ	મોટેભાગે વાદળછાયુ	મોટેભાગે વાદળછાયુ
સવારના ભેજના ટકા	૮૪	૮૦	૮૨	૮૧	૭૯
બપોર બાદ ભેજના ટકા	૭૩	૬૯	૬૯	૭૦	૬૯
પવનની ગતિ (કિમી/કલાક)	૦૦૬	૦૦૯	૦૧૨	૦૧૧	૦૧૫
પવનની દિશા (ડીગ્રી)	૧૯૮	૨૨૯	૨૩૫	૨૩૦	૨૨૮

પોરબંદર

વરસાદ (મી.મી)	૦	૦	૦	૦	૦
મહત્તમ તાપમાન (સે.ગ્રે)	૩૫	૩૫	૩૫	૩૫	૩૫
લઘુત્તમ તાપમાન (સે.ગ્રે)	૨૮	૨૮	૨૯	૨૯	૨૯
વાદળની સ્થિતિ	મોટેભાગે વાદળછાયુ	મોટેભાગે વાદળછાયુ	અંશત વાદળછાયુ	મોટેભાગે ચોખ્ખુ	અંશત વાદળછાયુ
સવારના ભેજના ટકા	૮૫	૮૧	૮૩	૮૨	૮૧
બપોર બાદ ભેજના ટકા	૭૩	૭૧	૭૦	૭૧	૭૧
પવનની ગતિ (કિમી/કલાક)	૦૦૪	૦૦૯	૦૧૨	૦૧૧	૦૧૪
પવનની દિશા (ડીગ્રી)	૨૦૦	૨૩૦	૨૩૫	૨૨૮	૨૨૬

દક્ષિણસૌરાષ્ટ્ર ખેત આબોહવાકિયવિભાગ

પાકનું નામ	પાક અવસ્થા	રોગ/જીવાત/ જાત	કૃષિ સલાહ
-	-	-	<ul style="list-style-type: none"> ➤ ઉનાળુ પાકોની કાપણી થઈ ગયા બાદ આખા ખેતરનું અવલોકન કરી જમીનના નમુના લેવા અને પૃથ્થકરણ કરવા માટે લેબોરેટરીમાં મોકલી આપવા. ➤ અગામી ચોમાસુ પાકોના વાવેતર માટે ખેત આયોજન કરવું તે મુજબ બીયારણ, ખાતર અને દવાનો પુરવઠો મેળવી લેવો. ➤ કોમ્પોસ્ટ, દેશી ખાતર અને જીપ્સમ ભલામણ મુજબ યાસમાં

			<p>ભરવા.</p> <ul style="list-style-type: none"> ➤ જમીનમાં ઉંડી ખેડ કરવી જેથી રોગ અને કીટકોની જુદી જુદી અવસ્થાએ સુર્યતાપ સામે ખુલ્લી થતા નાશ પામે. ➤ પિયતની સગવડ હોય તો ખરીફ મગફળીનું આગોતરું વાવેતર કરવું. વાવેતર માટે અર્ધવેલડી જાત- જીજી.૨૦ અથવા ઉભડી જાતો જીજી.૨ જીજી.૫ જીજી.૭ ટી.જી. ૨૬ ની પસંદગી કરવી.
ઉનાળુતલ	પરીપક્વ	ગુજરાત તલ-૨, ગુજરાત તલ-૩	❖ તલનો પાક પરીપક્વ થાય એટલે સમયસર કાપણી કરવી.
ઉનાળુમગફળી	પરીપક્વ	-	❖ મગફળીનો પાક પરીપક્વ થાય એટલે પીયત બંધ કરવું.ડોડવા પરીપક્વ થયે પીયત આપવાથી મગફળીનો ઉગાવો થવાથી ઉત્પાદન અને ગુણવત્તા ઘટે છેઅને સમયસર કાપણી કરવી.
ઉનાળુ મગ	પરીપક્વ	ગુજરાત મગ-૪, કે-૮૫૧	❖ મગના પાકની પરીપક્વ શીંગોની વીણી કરવી.
ઉનાળુ અડદ	પરીપક્વ	ગુજરાત અડદ- ૧, ટી - ૯	❖ અડદના પાકની પરીપક્વ શીંગોની વીણી કરવી.
આંબો	ફળ	-	<ul style="list-style-type: none"> ❖ આંબાની નાની કલમો ને ૭ થી ૮ દિવસે પીયત આપવું. અને પુખ્ત વય ના ઝાડને મહીને એકાદ પીયત આપવું. ❖ શાખ પડે એટલે પરીપક્વ કેરી ઉતારી ગ્રેડીંગ કરી બજારમાં મોકલવી.
શાકભાજી	વૃધ્ધી/ફુલ/ફળ	-	<ul style="list-style-type: none"> ❖ વેલાવાળા શાકભાજીના પાકમાં પાન કોરીયાનો ઉપદ્રવ ટાળવા માટે મુખ્ય પાકની ફરતે દિવેલા તથા ગલગોટાનું વાવેતર કરવાથી તેનું નિયંત્રણ કરી શકાય છે જો ઉપદ્રવ વધુ જોવા મળે તો ૧૦ મીલી મીથાઈલ ઓ ડેમેટોન (૨૫ ઈંસી) અથવા ૧૦ મીલી ડાયમીથોએટ (૩૦ ઈંસી) ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો ત્યાર બાદ પછીથી બીજો છંટકાવ ૧૫ દિવસ પછી કરવો. ❖ શાકભાજીના પાકમાં ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીનું મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦મીલી ૧૦ લી. પાણીમાં ભેળવી છંટકાવ કરવો. જો વધુ ઉપદ્રવ

			<p>હોય તો એસીટામાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦લી. અથવા દ્રાચઝોફોસ ૪૦ ટકા ૨૦ મીલી/ ૧૦ લી.પાણીમાં મીશ્ર કરી છંટકાવ કરવો.</p> <ul style="list-style-type: none"> ❖ મરચીના પાકમાં થ્રીપ્સના નિયંત્રણ માટે ૨૦મીલી પ્રોફેનોફોસ અને લિમ્બોળીનુ તેલ ૫૦ મીલી ૧૦ લીટર પાણી મિશ્રણ કરી છાંટવી. ❖ શાકભાજીનાપાકની રોપણી બાદ ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે દાણાદાર ફોરેટ ૧૦ જી છોડ દીઠ ૨ ગ્રામ મુજબ આપીને પીયત આપવું. ❖ મરચીના પાકમાં વાઈરસના કારણે પીળા થઈ કુકડાઈ ગયેલ છોડ ઉપાડી જમીનમાં દાટી નાશ કરવો અને ૫ મીલી ઈમીડાક્લોપ્રાઈડ ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ રીંગણ અને ટમેટાના ફળ અને અને ડોકાની ઈયળોનો ઉપદ્રવ જણાયતો તોનુકશાન પામેલ ફળ અને ડોકાને તોડીને જમીનમા દાટી દેવા અને રીનાક્ષીપાયર દવા ૩મીલીદવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ રીંગણ અને ટમેટાના પાકમા ફળમાખીના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા ૫ગ્રામ દવા ૧૦ લીટર પાણીમાં ભેળવી છંટકાવ કરવો. ❖ શાકભાજી ને સમયસર ઉતારી ગ્રેડીંગ કરી બજારમાં મુકવા. ❖ વધારે ગરમીના દિવસોમાં શાકભાજીનાપાકનેટુંકા ગાળે પીયત આપવું.
પશુપાલન	-	-	<ul style="list-style-type: none"> ❖ પશુઓને યોગ્ય આહાર આપવો અને પશુઓના રહેઠાણ ચોખ્ખા રાખવાઅને માખી મચ્છરથી રક્ષણ માટે ફીનાઈલ છાટવું. ❖ તાપમાન વધારે રહેવાના કારણે ઢોરને દિવસના સમયે છાયડામા રાખવા અને મિનરલ મીક્ષર્યુકતપીવાના સ્વચ્છ પાણીની વ્યવસ્થા કરવી. ❖ પશુઓમાં ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાઝ ૨ મીલી/ ૧ લીટર પાણીમાં નાખીને છાટવી. ❖ દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા

			બોરીક પાવડરથી બરાબર સાફ કરવા. ❖ પશુઓમા કાળીયા તાવના નિયંત્રણ માટે સમયસર રસીકરણ કરાવવું.
--	--	--	--

JUNAGADH

Phone: 0285-2670139

Date:- 02-06-2017

(Period from 3 to 7 June 2017)

Significant weather for the preceding week

Rain fall (mm)	000.0
Maximum temperature °C	36.5 - 39.9
Minimum temperature °C	26.0 – 26.4
RH % (Morning/Afternoon)	84 – 38
Wind speed (Kmph)	11.9 - 16.7
Cloud cover	0 - 4

Weather forecast for Junagadh Dist. until 08:30 hrs of 07-06-2017

Parameters	Day-1 03-06-17	Day-2 04-06-17	Day-3 05-06-17	Day-4 06-06-17	Day-5 07-06-17
Rain fall (mm)	0	0	0	0	3
Maximum temperature °C	39	40	39	39	39
Minimum temperature °C	25	25	26	26	26
Cloud cover (Okta)	6	5	3	2	5
RH I % (Morning)	84	80	82	81	79
RH II % (After noon)	73	69	69	70	69
Wind speed (Km/h)	006	009	012	011	015
Wind direction(Deg)	198	229	235	230	228

Agro meteorological Advisories

General Advice	Rain fall Temperature RH (%) Wind Direction and Wind Speed	<p>There is forecasting of light rainfall during next five days.</p> <p>Maximum and minimum temperature likely to be in range of 39-40°C and 25-26°C, respectively.</p> <p>The Relative Humidity is likely to be in the range of 84-69 %.</p> <p>Direction and speed of wind is likely to be SW, S and SSW and 06-15 km/h, respectively.</p>
-----------------------	---	--

Crop	Variety/ pest/insect	Stage	Agro met Advisories
			<ul style="list-style-type: none"> ➤ After harvesting of summer crops, take the soil samples and send to the soil testing laboratory for the soil analysis. ➤ Deep ploughing of fields after harvesting of summer crops is advised so that eggs and pupating stages of insects and weed seeds are destroyed. ➤ Procure good quality of seeds, Fertilizer and pesticide for <i>kharif</i> crops in time. ➤ If irrigation facility available than pre-sowing operation should be carried out of Kharif ground nut select semi-spread GG.20 or erect GG.2, GG.5, GG.7 and TG-26 varieties.
Summer Green gram	-	Maturity	➤ Harvesting operation should be carried out at maturity stage.
Summer Black gram	-	Maturity	➤ Harvesting operation should be carried out at maturity stage.
Summer Groundnut	-	Maturity	➤ Harvesting operation should be carried out at maturity stage.
Summer sesame	-	Maturity	➤ Harvesting operation should be carried out at maturity stage.
Mango	-	Fruit formation	➤ Apply irrigation regularly and send mango fruit to the market often grading.
Vegetable crops	-	Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water. ➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of

			<p>water at 15 days interval.</p> <ul style="list-style-type: none"> ➤ For the control of thrips in chilli, spray 20 ml profenophos in 10 litres of water. ➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. ➤ In present dry weather condition, spread plastic or fodder as mulch in vegetable fields for conserving moisture. ➤ Rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of 5 ml Imidacloprid per 10 litre of water is advised for control the vector, when sky remains clear. ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops, for the control of fruit borer, spray reenaxipayar @ 3.0ml in 10 lit of water. ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetemaprid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market often grading.
Livestock	-	-	<ul style="list-style-type: none"> ➤ To provide adequate food and keep the animals clean shed. Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day.

			<ul style="list-style-type: none"> ➤ To maintain optimum temperature between 18° to 29° C in poultry shed for better egg production. ➤ Apply layer of grass or gunny bags on the roof of poultry shed and sprinkle water on it ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ For prevention of Anthrax disease in animals do vaccination with anthrax vaccine. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.
--	--	--	---

PORBANDAR

Phone: 0285-2670139

Date:-02-06-2017

(Period from 03 to 07 June-2017)

Weather forecast for Porbandar Dist. until 08:30 hrs of 07-06-2017

Parameters	Day-1 03-06-17	Day-2 04-06-17	Day-3 05-06-17	Day-4 06-06-17	Day-5 07-06-17
Rain fall (mm)	0	0	0	0	0
Maximum temperature °C	35	35	35	35	35
Minimum temperature °C	28	28	29	29	29
Cloud cover (Okta)	6	5	3	1	4
RH I % (Morning)	85	81	83	82	81
RH II % (After noon)	73	71	70	71	71
Wind speed (Km/h)	004	009	012	011	014
Wind direction(Deg)	200	230	235	228	226

Agro meteorological Advisories

General Advice	Rain fall Temperature RH (%) Wind Direction and Wind Speed	<p>There is no forecasting of rainfall during next five days.</p> <p>Maximum and minimum temperature likely to be in range of 35°C and 28-29°C, respectively.</p> <p>The Relative Humidity is likely to be in the range of 85-70%.</p> <p>Direction and speed of wind is likely to be SW and SSW and 08-20 km/h, respectively.</p>
-----------------------	---	--

Crop	Variety/ pest/insect	Stage	Agro met Advisories
			<ul style="list-style-type: none"> ➤ After harvesting of summer crops, take the soil samples and send to the soil testing laboratory for the soil analysis. ➤ Deep ploughing of fields after harvesting of summer crops is advised so that eggs and pupating stages of insects and weed seeds are destroyed. ➤ Procure good quality of seeds, Fertilizer and pesticide for <i>kharif</i> crops in time. ➤ If irrigation facility available than pre-sowing operation should be carried out of <i>Kharif</i> ground nut select semi-spread GG.20 or erect GG.2, GG.5, GG.7 and TG-26 varieties.
Summer Green gram	-	Maturity	➤ Harvesting operation should be carried out at maturity stage.
Summer Black gram	-	Maturity	➤ Harvesting operation should be carried out at maturity stage.
Summer Groundnut	-	Maturity	➤ Harvesting operation should be carried out at maturity stage.
Summer sesame	-	Maturity	➤ Harvesting operation should be carried out at maturity stage.
Mango	-	Fruit formation	➤ Apply irrigation regularly and send mango fruit to the market often grading.
Vegetable crops	-	Flowering/ Fruiting	<ul style="list-style-type: none"> ➤ For the control of leaf webber in fruit leafy vegetable, sowing of trap crop (castor and galgota) around the vegetable crop. For the heavy incidence spray 10 ml methyl O demeton (25 EC) Or 10 ml diamethoate (30 EC) in 10 liter of water. ➤ For the control of thrips and white fly, spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit

			<p>of water at 15 days interval.</p> <ul style="list-style-type: none"> ➤ For the control of thrips in chilli, spray 20 ml profenophosin in 10 litres of water. ➤ For the control of fruit and shoot borer, spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP @ 5.0 gm / 10 lit of water at 15 days interval. ➤ In present dry weather condition, spread plastic or fodder as mulch in vegetable fields for conserving moisture. ➤ Rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of 5 ml imidacloprid per 10 litre of water is advised for control of the vector, when sky remains clear. ➤ Apply regular irrigation at 6 to 8 days to the crop at the flowering and fruiting stage. ➤ In brinjal and tomato crops, for the control of fruit borer, spray renoxipayar @ 3.0ml in 10 lit of water. ➤ In brinjal and tomato crops, for the control of fruit fly, spray acetamiprid 5.0 ml in 10 lit of water. ➤ Keep vegetable crop weed free, apply irrigation regularly and send vegetable fruit to the market after grading.
--	--	--	--

<p>Livestock</p>	<p>-</p>	<p>-</p>	<ul style="list-style-type: none"> ➤ To provide adequate food and keep the animals clean shed. Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day. ➤ To maintain optimum temperature between 18° to 29° C in poultry shed for better egg production. ➤ Apply layer of grass or gunny bags on the roof of poultry shed and sprinkle water on it ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ For prevention of Anthrax disease in animals do vaccination with anthrax vaccine. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.
-------------------------	----------	----------	--