



ગ્રામીણ કૃષિ મૌસમ સેવા
હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ
તા.01-02-2017 થી તા.05-02-2017



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કૃ.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન.)

કૃષિહવામાન બુલેટીન નં. **88/2017**

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી)	0.0
મહત્તમ તાપમાન (સે.ગ્રે)	25.3-34.0
લઘુત્તમ તાપમાન (સે.ગ્રે)	10.6-12.4
ભેજ(%) સવાર	59-90
ભેજ(%) બપોરબાદ	40-50
પવનની ગતિ (કિમી/કલાક)	5.0-5.1

હવામાન આગાહી: પત્રક-ક

ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.01-02-2017 થી 05-02-2017

કૃષિ સલાહ:

સામાન્ય	<p>ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના રાજકોટ</p> <p>રાજકોટ: જિલ્લામાં આગામી તા.01-02-2017 થી તા.05-02-2017 દરમિયાન હવામાન સુકું અને મહદ અન્શેયોખ્મુ રહેશે. પવનની દિશા ઈશાન, પૂર્વ અને વાયવ્યની રહેવાની અને પવનની ઝડપ 10 થી 21 કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન 31-33 ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન 12-14 ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</p>																														
	<table border="1"><thead><tr><th>તારીખ</th><th>દિવસ-1</th><th>દિવસ-2</th><th>દિવસ-3</th><th>દિવસ-4</th><th>દિવસ-5</th></tr><tr><th>પેશમીટર</th><th>01-02-17</th><th>02-02-17</th><th>03-02-17</th><th>04-02-17</th><th>05-02-17</th></tr></thead><tbody><tr><td>વરસાદ(મી.મી)</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>મહત્તમ તાપમાન (સે.ગ્રે)</td><td>32</td><td>32</td><td>33</td><td>33</td><td>31</td></tr><tr><td>લઘુત્તમ તાપમાન (સે.ગ્રે)</td><td>13</td><td>13</td><td>14</td><td>13</td><td>12</td></tr></tbody></table>	તારીખ	દિવસ-1	દિવસ-2	દિવસ-3	દિવસ-4	દિવસ-5	પેશમીટર	01-02-17	02-02-17	03-02-17	04-02-17	05-02-17	વરસાદ(મી.મી)	0	0	0	0	0	મહત્તમ તાપમાન (સે.ગ્રે)	32	32	33	33	31	લઘુત્તમ તાપમાન (સે.ગ્રે)	13	13	14	13	12
તારીખ	દિવસ-1	દિવસ-2	દિવસ-3	દિવસ-4	દિવસ-5																										
પેશમીટર	01-02-17	02-02-17	03-02-17	04-02-17	05-02-17																										
વરસાદ(મી.મી)	0	0	0	0	0																										
મહત્તમ તાપમાન (સે.ગ્રે)	32	32	33	33	31																										
લઘુત્તમ તાપમાન (સે.ગ્રે)	13	13	14	13	12																										

વાદળનીસ્થિતિ(ઓકટા)	0	0	0	2	2
મહતમ ભેજ (%) સવાર	60	39	34	38	80
લઘુતમ ભેજ (%) બપોરબાદ	26	20	19	20	30
પવનની ગતિ (મી/કલાક)	13	16	10	12	21
પવનની દિશા (ડીગ્રી)	80	31	50	305	310

જામનગર: જીલ્લામાં આગામી તા.01-02-2017 થી તા.05-02-2017 દરમ્યાન હવામાન સુકું અને મહદ અન્શે યોખ્મુ રહેશે. પવનની દિશા અગ્નિ, ઈશાન અને વાયવ્યની રહેવાની અને પવનની ઝડપ **10 થી 25 કીમી/કલાક** રહેવાની શક્યતા છે. આ સમયગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન **27-28** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન **13-13** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-1	દિવસ-2	દિવસ-3	દિવસ-4	દિવસ-5
પેરામીટર	01-02-17	02-02-17	03-02-17	04-02-17	05-02-17
વરસાદ(મી.મી)	0	0	0	0	0
મહતમ તાપમાન (સે.ગ્રે)	27	28	28	27	27
લઘુતમ તાપમાન (સે.ગ્રે)	12	12	13	13	12
વાદળની સ્થિતિ(ઓકટા)	0	0	0	2	2
મહતમ ભેજ (%) સવાર	69	41	34	50	76
લઘુતમ ભેજ (%) બપોરબાદ	40	24	21	23	42
પવનની ગતિ (કિમી/કલાક)	14	15	10	14	25
પવનની દિશા (ડીગ્રી)	169	33	54	297	304

અમરેલી: જીલ્લામાં આગામી તા. 01-02-2017 થી તા.05-02-2017 દરમ્યાન હવામાન સુકું અને મહદ અન્શે યોખ્મુ રહેશે. પવનની દિશા અગ્નિ, પૂર્વ અને વાયવ્યની રહેવાની અને પવનની ઝડપ **10 થી 17 કીમી/કલાક** રહેવાની શક્યતા છે. આ સમયગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન **33-34** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન **12-14** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-1	દિવસ-2	દિવસ-3	દિવસ-4	દિવસ-5
પેરામીટર	01-02-17	02-02-17	03-02-17	04-02-17	05-02-17
વરસાદ(મી.મી)	0	0	0	0	0
મહતમ તાપમાન (સે.ગ્રે)	33	33	34	34	33
લઘુતમ તાપમાન (સે.ગ્રે)	13	13	14	13	12
વાદળની સ્થિતિ(ઓકટા)	0	0	0	2	2
મહતમ ભેજ (%) સવાર	56	39	36	38	83

લઘુતમ ભેજ (%) બપોરબાદ	24	22	22	24	30
પવનની ગતિ કિમી/કલાક	13	16	10	10	17
પવનની દિશા (ડીગ્રી)	114	84	81	319	325



સુરેન્દ્રનગર: જીલ્લામાં આગામી તા.01-05-2017 થી તા. 05-02-2017 દરમ્યાન હવામાન સુકું અને યોખ્ખુ રહેશે. પવનની દિશા પૂર્વ, ઈશાન અને વાયવ્યની રહેવાની અને પવનની ઝડપ 10 થી 16 કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન 32-33 ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન 12-13 ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.




તારીખ પેરામીટર	દિવસ-1 01-02-17	દિવસ-2 02-02-17	દિવસ-3 03-02-17	દિવસ-4 04-02-17	દિવસ-5 05-02-17
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમ તાપમાન (સે.ગ્રે)	32	32	33	33	33
લઘુત્તમ તાપમાન (સે.ગ્રે)	12	12	13	13	12
વાદળની સ્થિતિ(ઓકટા)	0	0	0	1	1
મહત્તમ ભેજ (%) સવાર	49	34	31	34	77
લઘુત્તમ ભેજ (%) બપોરબાદ	25	17	16	16	22
પવનની ગતિ કિમી/કલાક	12	14	10	10	16
પવનની દિશા (ડીગ્રી)	73	66	63	272	303




5 ફેબ્રુઆરી સુધી હવામાન સુકું અને મહદ અન્શે યોખ્ખુ રહેશે. પવનની દિશા ઈશાન, પૂર્વ અને વાયવ્યની રહેવાની અને પવનની ઝડપ 10 થી 20 કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન 32-33 ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન 12-14 ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે. તેમજ તા. 5 ફેબ્રુઆરીના રોજ જાકળ આવવાની શક્યતા છે.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
ઉનાળુ મગફળી	વાવેતર	ટીએજી-૨૪, જીજી-૨, તીજી-૩૭-એ, ટીપીજી-૪૧	આ પખવાડીયામાં તાપમાન ૨૨-૨૫° સે. થાય એટલે ઉનાળુ મગફળીનું વાવેતર કરી દેવું. વાવેતર સમયે મગફળીના બીયાને ૧.૨૫ ગ્રામ ટેબ્યુકોનાઝોલ દવાનો ૧ કિ.ગ્રા. દાણા પ્રમાણે પટ આપવો. સીંગલ સુપર ફોસ્ફેટ ૩૧૨ કિ.ગ્રા., એમોનીયમ સલ્ફેટ ૧૨૫ કિ.ગ્રા. અને એરંડીનો ખોળ ૨૫૦ કિ.ગ્રા./ હેક્ટર અથવા

			ડી.એ.પી. ૧૦૮ કિ.ગ્રા. અને જીપ્સમ ૨૫૦ કિ.ગ્રા./ હેક્ટર અને એરંડીનો ખોળ ૨૫૦ કિ.ગ્રા./હેક્ટર પાયામાં આપવો.
ઉનાળુ તલ	વાવેતર	ગુજરાત તલ-૨ ગુજરાત તલ-૩	આ પખવાડીયામાં તાપમાન ૨૮-૩૦° સે. થાય એટલે ઉનાળુ તલનું વાવેતર કરી દેવું. બિયારણને ૨-૩ ગ્રામ/કિલો મુજબ થાયરમ-કાર્બોક્સીન, મેન્કોઝેબ અથવા કાર્બેન્હાઝીમનો પટ આપવો. -પાયામાં સીંગલ સુપર ફોસ્ફેટ ૨૫ કિ.ગ્રા./વિઘે અને યુરીયા ૮ કિ.ગ્રા./વિઘે આપવું. અથવા - ડી.એ.પી.૯ કિ.ગ્રા. અને જીપ્સમ ૮૦ કિ.ગ્રા./વિઘે અને યુરીયા ૮ કિ.ગ્રા./વિઘે આપવું. તેમજ તલના પાકમાં સરકારી ગ્રેડ-૫ વાળું સૂક્ષ્મતત્વોનું મિશ્રણ ૩.૨ કિ.ગ્રા./વિઘે અને એરંડીનો ખોળ ૮૦ કિ.ગ્રા./વિઘે જમીનમાં આપવો.
ટમેટા	ફેર રોપણી	પુસા હાઈબ્રીડ, પુસા બી, જુનાગઢ રૂબી, ગુજરાત ટમેટા-૧	વાવેતર અંતર ૬૦-૯૦ × ૪૫-૬૦ સે.મી. રાખવું તથા યુરીયા ૧૬૩, સીંગલ સુપર ફોસ્ફેટ ૬૨૫ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૧૬૭ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
રીંગણ	ફેર રોપણી	જુનાગઢ લંબગોળ, રીંગણ જુનાગઢ લાંબા, ગુજરાત રીંગણ-૨	વાવેતર અંતર ૯૦ × ૬૦ સે.મી. રાખવું તથા યુરીયા ૧૦૮, સીંગલ સુપર ફોસ્ફેટ ૩૧૨ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૮૩ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
ભીંડા	વાવેતર	ગુજરાત હાઈબ્રીડ ભીંડા-૧, ગુજરાત ભીંડા -૨	વાવેતર અંતર ૬૦ × ૪૫ સે.મી. રાખવું તથા યુરીયા ૧૦૮, સીંગલ સુપર ફોસ્ફેટ ૩૧૨ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૮૩ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
તુરીય	વાવેતર	પુસા નસદાર, જયપુરી	વાવેતર અંતર ૧.૫ × ૦.૫ મીટર રાખવું તથા યુરીયા ૨૭, સીંગલ સુપર ફોસ્ફેટ ૭૮ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૨૧ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
દૂધી	વાવેતર	પુસા સમર પ્રોલીફીક લોંગ,	વાવેતર અંતર ૨ × ૦.૫ મીટર રાખવું તથા યુરીયા

		જુનાગઢ ટેન્ડર લોગ	૧૦૮, સીંગલ સુપર ફોસ્ફેટ ૩૧૨ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૮૩ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
ગલકા	વાવેતર	પુસા ચીકની	વાવેતર અંતર ૧.૫ × ૦.૫ મીટર રાખવું તથા યુરીયા ૨૭, સીંગલ સુપર ફોસ્ફેટ ૭૮ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૨૧ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
કારેલા	વાવેતર	પ્રિયાપુસા, દો મોસમી	વાવેતર અંતર ૧ × ૧ મીટર રાખવું તથા યુરીયા ૬૫, સીંગલ સુપર ફોસ્ફેટ ૩૭૫ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૧૦૦ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
ગુવાર	વાવેતર	પુસા સદાબહાર, પુસા નવબહાર, આઈસી-૧૧૩૮૮	વાવેતર અંતર ૬૦ × ૩૦ સે.મી. રાખવું તથા યુરીયા ૫૫, સીંગલ સુપર ફોસ્ફેટ ૨૩૫ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૬૩ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
કાકડી	વાવેતર	જાપાનીઝ લાંબી લીલી, સ્ટ્રેઇટ-૮, પુસા સંયોગ	વાવેતર અંતર ૧ × ૧ મીટર રાખવું તથા યુરીયા ૫૫, સીંગલ સુપર ફોસ્ફેટ ૧૫૬ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૪૨ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
તરબુચ	વાવેતર	સેગર બેબી, અરકા જ્યોતી	વાવેતર અંતર ૨ × ૧ મીટર રાખવું તથા યુરીયા ૧૦૮, સીંગલ સુપર ફોસ્ફેટ ૩૧૨ કિ.ગ્રા., અને મ્યુરેટ ઓફ પોટાશ ૮૩ કિ.ગ્રા./ હેક્ટર પાયામાં આપવો.
લસણ	ગાંજો બન્યાવો	શ્રીપ્સ, પાનની ટોચનો  સુકારો 	શ્રીપ્સના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને પ્રોફેનોફોસ ૨૦ મિલી અથવા લેમ્બડા સાયહેલોથ્રીન ૧૫ મિલી દવાને ૧૦ લીટર પાણીમાં મિશ્ર લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. પાનની ટોચનો સુકારાના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામ ૧૦ લીટર પાણીમાં કરીને ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.
ચણા	કુલ/પોપટા	લીલી અને લશ્કરી ઈયળ	ચણાના પાકમાં લીલી અને લશ્કરી ઈયળના

	બન્ધાવા/ દાણા ભરાવા		નિયંત્રણ માટે પક્ષીઓને બેસવા માટે 'T' આકારના બેલાખડા મુકવા. જો વધુ ઉપદ્રવ જણાય તો એમામેકટીન બેન્ઝોએટ ૬.૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.
ઘઉં	દુધિયા દાણા ભરાવા	-	સમયસર પિયત આપવું.
જીરું	કુલ અને દાણા બન્ધાવા	થ્રીપ્સ અને મોલો ચરમી અને રાખોડી  સુકારો 	થ્રીપ્સ અને મોલોના ના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો. તેમજ ચરમીનો ઉપદ્રવ ન થાય તે માટે ડાયથેન એમ ૪૫, ૨૭ ગ્રામ અને હેક્ઝાકોનાઝોલ ૧૫ મિલી દવા વારાફરતી ૧૦ લીટર પાણીમાં મિશ્ર કરીને ૧૫ દિવસના અન્તરે છંટકાવ કરવો.  ઉભા પાકમાં સુકારાના નિયંત્રણ માટે પિયત આપવાનું બંધ કરવું.
ધાણા	દાણા બન્ધાવા	થ્રીપ્સ	થ્રીપ્સ અને મોલોના ના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.
ડુંગળી	વૃદ્ધિ	જાંબલી ધાબા	ડુંગળીના પાકમાં જાંબલી ધાબાના રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ

			૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામ ૧૦ લીટર પાણીમાં કરીને ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.
રીંગણી અને ટમેટી	કુલ/ફળ	સફેદ માખી ડુંખ અને ફળ કોરી ખાનાર ઈયળ → કથીરી →	<ul style="list-style-type: none"> ➤ લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ ફળ અને ડોકાની ઈયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને રીનાક્ષીપાયર દવા ૩ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ રીંગણી અને ટમેટીના પાકમાં પાન કથીરીના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
કોબી અને ફલાવર	વૃદ્ધિ/દડા	હિરાફૂદુ	કોબી અને ફલાવરમાં હિરાફૂદાણા નિયંત્રણ માટે સ્પીનોસાઈ ૩ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો
મરચી	કુલ/ફળ/વિણિ	પાનની કુકડ-થ્રીપ્સ→  કાલવ્રણ → 	<p>પ્રોફેનોફોસ ૨૦ મી.લી.અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>કાલવ્રણ રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.</p>

પશુ/મરઘા

પશુ/મરઘા



નાના વાછરડાઓને રાત્રે ઠંડીથી બચવા પશુ ઘરમાં રખવા. પ્રજીવથી થતા બૃસેલ્લા અને અન્ય રોગોની ચકાસણી કરાવવી. ખરવા મોવાસાના રોગનું રસીકરણ કરાવવું. ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મી.લી. ૧ લીટર પાણીમાં નાખીને છાંટવી. વિટામીન એ તથા ડી પુરકો આપવા. કરમિયાની દવા આપવી તથા ઉછરતા પશુઓમાં બૃસેલ્લા કાફવુંડનું રસીકરણ કરાવવું. મરઘા ઘરના પસ્ટિચિમ દિશામાં કંતાનના પડદા લગાવવા.

જુક્યુ/ટેકનિકલ ઓફિસર.(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૧૭,
તરઘડિયા(રાજકોટ), તારીખ: 31/01/2017.

પ્રતિ,

તંત્રીશ્રી/ડાયરેક્ટરશ્રી,

રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનંતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર
સુકી ખેતી સંશોધન કેન્દ્ર
જુનાગઢ કૃષિ યુનિવર્સિટી
તરઘડીયા



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-88 for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 1-5 February- 2017)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	0.0
Maximum temperature (°C)	25.3-34.0
Minimum temperature (°C)	10.6-12.4
Relative Humidity (%) I	59-90
Relative Humidity (%) II	40-50
Wind speed (Kmph)	5.0-5.1



Weather forecast until 08-30 hrs of 05/02/2017




(1) RAJKOT DISTRICT:




	01/02/17	02/02/17	03/02/17	04/02/17	05/02/17
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	32	32	33	33	31
Min.Temp.(°C)	13	13	14	13	12
Total CC (octa)	0	0	0	2	2
RH-I (%)	60	39	34	38	80
RH-II (%)	26	20	19	20	30
Wind Speed (kmph)	13	16	10	12	21
Wind Direction(deg.)	80	31	50	305	310

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: Dry and mainly clear weather forecasted for next five days.• Temperature: Maximum and minimum temperature likely to be in range of 31-33°C and 12-14°C, respectively.• Relative Humidity: The RH at morning and noon is likely to be in the range of 38-80 % and 19-30 %, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be E, NE, NW and 10-21 km/h, respectively.		
Crops	Stage	Insect/Pest/Variety	Advisory
Summer Groundnut	Sowing	TAG-24,TG-37-A, TPG-41,GG-2, GJG-31	Carry out sowing of summer groundnut as temperature increases to 22-25° C and give seed treatment of Tebuconazole 1.25 gm/1 kg seed. Apply Single Super Phosphate 312 kg, Ammonium sulphate 125 kg and Castor cake 250

			kg/ha OR DAP 108 kg, Gypsum 250 kg and castor cake 250 kg/ha as basal.
Summer Sesame	Preparation for sowing	G.Til-2, G.Til-3	Procure good quality seeds of sesame and carry out sowing of summer SESAME as minimum and maximum temperature rises up to 19 °C and 30°C, respectively and give seed treatment of vitavax (Carboxin 37.5% + Thiram 37.5%) or mencozeb or carbendazim 2-3 gm/1 kg seed. Apply Single Super Phosphate 156 kg and 55 kg urea/ha, OR DAP 54 kg and 33 kg urea/ha as basal.
Brinjal	Transplan-ting	Junagadh oblong, Junagadh long, Guj. Bringle-2	Keep planting distance 90 X 60 cm ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Lady finger	Sowing	Guj. Hy.-1 & 2	Keep planting distance 60 X 45 cm ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Ridge gourd	Sowing	Pusa nasdar, Jaipuri,	Keep planting distance 1.5 X 0.5 m ² and apply urea 27, single super phosphate 78 and murret of potash 21 kg/ha at sowing time.
Bottle gourd	Sowing	Junagadh tender long, Pusa summer prolific long	Keep planting distance 2 X 0.5 m ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Sponge guard	Sowing	Pusa chikni	Keep planting distance 1.5 X 0.5 m ² and apply urea 27, single super phosphate 78 and murret of potash 21 kg/ha at sowing time.
Bitter guard	Sowing	Priya, Pusa domosmi	Keep planting distance 1.0 X 1.0 m ² and apply urea 65, single super phosphate 375 and murret of potash 100 kg/ha at sowing time.
Cluster bean	Sowing	Pusa sadabahar, Pusa navbahar, IC-11388	Keep planting distance 60 X 30 cm ² and apply urea 55, single super phosphate 235 and murret of potash 63kg/ha at sowing time.
Water melon	Sowing	Sugar baby, Arka jyoti	Keep planting distance 2.0 X 1.0 m ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Garlic	Girthing	<p>Thrips</p>  <p>Alternaria blight/Purple Blotch</p> 	<p>In present weather conditions attack of thrips has been reported. Make alternative sprays of imidaciprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval.</p> <p>The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop.</p> <p>Make two spray of 27 gm Mencozeb or thiophanae methyl 12 gm/10 liters of water at 10 days interval.</p>

Chick pea	Pod formation/development	Monitor the crop regularly for prodenia and heliothes infestation	“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation cross ETL spray 3 ml SPINOSAD in 10 litres of water.
Wheat	Grain filling	-	Apply irrigation without stress.
Cumin	Grain formation/development	Aphid and thrips Blight and Powery mildew 	Under present weather condition attack of thrips and aphids has been reported spray imidacloprid 5 ml /10 liters of water. Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation. 
Coriander	Grain formation/development	Aphid, Thrips and Jassids	Spray neem seed extract 5% and Thiomethoxam 5 gm/ 10 liters of water. Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent powdery mildew infestation.
Onion	Growth	Thrips Purple blotch → 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Spray Mencozeb 25 gm and hexaconazole 15 ml in 10 liters of water at 10 days interval.
Brinjal & Tomato	Flowering/fruiting	Sucking pest → & Fruit borer	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.

Cabbage and cauliflower	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.
Chili	Flowering & fruiting	Thrips, →  Wilting → Anthracnose → 	In present weather conditions severe attack of thrips has been reported. Spray 3 ml spinosad in 10 litres of water. Drenching of Trichoderma viridi 4 gm in 1 liter of water. Spray Mencozeb 27 gm in 10 liters of water
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose. Put curtains on west side of poultry house.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-88 for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
 (Period 1-5 February- 2017)



Weather forecast until 08-30 hrs of 05/02/2017





(2) JAMNAGAR DISTRICT:




	01/02/17	02/02/17	03/02/17	04/02/17	05/02/17
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	27	28	28	27	27
Min.Temp.(°C)	12	12	13	13	12
Total CC (octa)	0	0	0	2	2
RH-I (%)	69	41	34	50	76
RH-II (%)	40	24	21	23	42
Wind Speed (kmph)	14	15	10	14	25
Wind Direction(deg.)	169	33	54	297	304

Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry and mainly clear weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 27-28°C and 12-13°C, respectively. • Relative Humidity: The RH at morning and noon is likely to be in the range of 34-76 % and 21-42 %, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be S, NE, NW and 10-25 km/h, respectively. 		
Crops	Stage	Insect/Pest/Variety	Advisory
Summer Groundnut	Sowing	TAG-24,TG-37-A, TPG-41,GG-2, GJG-31	Carry out sowing of summer groundnut as temperature increases to 22-25° C and give seed treatment of Tebuconazole 1.25 gm/1 kg seed. Apply Single Super Phosphate 312 kg, Ammonium sulphate 125 kg and Castor cake 250 kg/ha OR DAP 108 kg, Gypsum 250 kg and castor cake 250 kg/ha as basal.
Summer Sesame	Preparation for sowing	G.Til-2, G.Til-3	Procure good quality seeds of sesame and carry out sowing of summer SESAME as minimum and maximum temperature rises up to 19 °C and 30°C, respectively and give seed treatment of vitavax (Carboxin 37.5% + Thiram 37.5%) or mencozeb or carbendazim 2-3 gm/1 kg seed. Apply Single Super Phosphate 156 kg and 55 kg urea/ha, OR

			DAP 54 kg and 33 kg urea/ha as basal.
Brinjal	Transplan-ting	Junagadh oblong, Junagadh long, Guj. Bringle-2	Keep planting distance 90 X 60 cm ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Lady finger	Sowing	Guj. Hy.-1 & 2	Keep planting distance 60 X 45 cm ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Ridge gourd	Sowing	Pusa nasdar, Jaipuri,	Keep planting distance 1.5 X 0.5 m ² and apply urea 27, single super phosphate 78 and murret of potash 21 kg/ha at sowing time.
Bottle gourd	Sowing	Junagadh tender long, Pusa summer prolific long	Keep planting distance 2 X 0.5 m ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Sponge guard	Sowing	Pusa chikni	Keep planting distance 1.5 X 0.5 m ² and apply urea 27, single super phosphate 78 and murret of potash 21 kg/ha at sowing time.
Bitter guard	Sowing	Priya, Pusa domosmi	Keep planting distance 1.0 X 1.0 m ² and apply urea 65, single super phosphate 375 and murret of potash 100 kg/ha at sowing time.
Cluster bean	Sowing	Pusa sadabahar, Pusa navbahar, IC-11388	Keep planting distance 60 X 30 cm ² and apply urea 55, single super phosphate 235 and murret of potash 63kg/ha at sowing time.
Water melon	Sowing	Sugar baby, Arka jyoti	Keep planting distance 2.0 X 1.0 m ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Garlic	Girthing	Thrips  Alternaria blight/Purple Blotch 	In present weather conditions attack of thrips has been reported. Make alternative sprays of imidaciprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval. The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop. Make two spray of 27 gm Mencozeb or thiophanae methyl 12 gm/10 liters of water at 10 days interval.
Chick pea	Pod formation/ development	Monitor the crop regularly for prodenia and	“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation cross ETL spray 3 ml SPINOSAD in

		heliiothes infestation	10 litres of water.
Wheat	Grain filling	-	Apply irrigation without stress.
Cumin	Grain formation/development	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray imidacloprid 5 ml/10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  
Coriander	Grain formation/development	Aphid, Thrips and Jassids	<p>Spray neem seed extract 5% and Thiomethoxam 5 gm/ 10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent powdery mildew infestation.</p>
Onion	Growth	<p>Thrips</p> <p>Purple blotch →</p> 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Spray Mencozeb 25 gm and hexaconazole 15 ml in 10 liters of water at 10 days interval.</p>
Brinjal & Tomato	Flowering/fruiting	<p>Sucking pest →</p> <p>& Fruit borer</p>	<p>To control sucking pest spray neem seed extract 5% or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Cabbage and cauliflower	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.

<p>Chili</p>	<p>Flowering & fruiting</p>	<p>Thrips, →  Wilting → Anthracnose → </p>	<p>In present weather conditions severe attack of thrips has been reported. Spray 3 ml spinosad in 10 litres of water.</p> <p>Drenching of Trichoderma viridi 4 gm in 1 liter of water.</p> <p>Spray Mencozeb 27 gm in 10 liters of water</p>
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose. Put curtains on west side of poultry house.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-88 for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
 (Period 1-5 February-2017)



Weather forecast until 08-30 hrs of 05/02/2017




(3) AMRELI DISTRICT:




	01/02/17	02/02/17	03/02/17	04/02/17	05/02/17
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	33	33	34	34	33
Min.Temp.(°C)	13	13	14	13	12
Total CC (octa)	0	0	0	2	2
RH-I (%)	56	39	36	38	83
RH-II (%)	24	22	22	24	30
Wind Speed (kmph)	13	16	10	10	17
Wind Direction(deg.)	114	84	81	319	325

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry and mainly clear weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 33-34°C and 12-14°C, respectively. • Relative Humidity: The RH at morning and noon is likely to be in the range of 36-83 % and 22-30 %, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be SE, NE, NW and 10-17 km/h, respectively. 		
Crops	Stage	Insect/Pest/Variety	Advisory
Summer Groundnut	Sowing	TAG-24,TG-37-A, TPG-41,GG-2, GJG-31	Carry out sowing of summer groundnut as temperature increases to 22-25° C and give seed treatment of Tebuconazole 1.25 gm/1 kg seed. Apply Single Super Phosphate 312 kg, Ammonium sulphate 125 kg and Castor cake 250 kg/ha OR DAP 108 kg, Gypsum 250 kg and castor cake 250 kg/ha as basal.
Summer Sesame	Preparation for sowing	G.Til-2, G.Til-3	Procure good quality seeds of sesame and carry out sowing of summer SESAME as minimum and maximum temperature rises up to 19 °C and 30°C, respectively and give seed treatment of vitavax (Carboxin 37.5% + Thiram 37.5%) or mencozeb or carbendazim 2-3 gm/1 kg seed. Apply Single Super Phosphate 156 kg and 55 kg urea/ha, OR

			DAP 54 kg and 33 kg urea/ha as basal.
Brinjal	Transplan-ting	Junagadh oblong, Junagadh long, Guj. Bringle-2	Keep planting distance 90 X 60 cm ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Lady finger	Sowing	Guj. Hy.-1 & 2	Keep planting distance 60 X 45 cm ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Ridge gourd	Sowing	Pusa nasdar, Jaipuri,	Keep planting distance 1.5 X 0.5 m ² and apply urea 27, single super phosphate 78 and murret of potash 21 kg/ha at sowing time.
Bottle gourd	Sowing	Junagadh tender long, Pusa summer prolific long	Keep planting distance 2 X 0.5 m ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Sponge guard	Sowing	Pusa chikni	Keep planting distance 1.5 X 0.5 m ² and apply urea 27, single super phosphate 78 and murret of potash 21 kg/ha at sowing time.
Bitter guard	Sowing	Priya, Pusa domosmi	Keep planting distance 1.0 X 1.0 m ² and apply urea 65, single super phosphate 375 and murret of potash 100 kg/ha at sowing time.
Cluster bean	Sowing	Pusa sadabahar, Pusa navbahar, IC-11388	Keep planting distance 60 X 30 cm ² and apply urea 55, single super phosphate 235 and murret of potash 63kg/ha at sowing time.
Water melon	Sowing	Sugar baby, Arka jyoti	Keep planting distance 2.0 X 1.0 m ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Garlic	Girthing	<p>Thrips</p>  <p>Alternaria blight/Purple Blotch</p> 	<p>In present weather conditions attack of thrips has been reported. Make alternative sprays of imidaciliprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval.</p> <p>The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop. Make two spray of 27 gm Mencozeb or thiophanae methyl 12 gm/10 liters of water at 10 days interval.</p>
Chick pea	Pod formation/ development	Monitor the crop regularly for prodenia and heliothes infestation	“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation cross ETL spray 3 ml SPINOSAD in 10 litres of water.
Wheat	Grain filling	-	Apply irrigation without stress.

Cumin	Grain formation/development	Aphid and thrips Blight and Powery mildew 	<p>Under present weather condition attack of thrips and aphids has been reported spray imidacloprid 5 ml/10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p> 
Coriander	Grain formation/development	Aphid, Thrips and Jassids	<p>Spray neem seed extract 5% and Thiomethoxam 5 gm/ 10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent powdery mildew infestation.</p>
Onion	Growth	Thrips Purple blotch → 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Spray Mencozeb 25 gm and hexaconazole 15 ml in 10 liters of water at 10 days interval.</p>
Brinjal & Tomato	Flowering/fruiting	Sucking pest → & Fruit borer	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Cabbage and cauliflower	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.
Chili	Flowering & fruiting	Thrips, →	In present weather conditions severe attack of thrips has been reported. Spray 3 ml spinosad in 10 litres of water.

		 <p>Wilting →</p>  <p>Anthracnose →</p>	<p>Drenching of <i>Trichoderma viridi</i> 4 gm in 1 liter of water.</p> <p>Spray Mencozeb 27 gm in 10 liters of water</p>
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose. Put curtains on west side of poultry house.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:94277 28523



Agro met Advisory Service bulletin No.-88 for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 1-5 February-2017)



Weather forecast until 08-30 hrs of 05/02/2017




(4) SURENDRANAGAR DISTRICT:




	01/02/17	02/02/17	03/02/17	04/02/17	05/02/17
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	32	32	33	33	33
Min.Temp.(°C)	12	12	13	13	12
Total CC (octa)	0	0	0	1	1
RH-I (%)	49	34	31	34	77
RH-II (%)	25	17	16	16	22
Wind Speed (kmph)	12	14	10	10	16
Wind Direction(deg.)	73	66	63	272	303

Agro meteorological advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry and clear weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 32-33°C and 12-13°C, respectively. • Relative Humidity: The RH at morning and noon is likely to be in the range of 31-77 % and 16-25 %, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be SE, NE, NW and 10-16 km/h, respectively. 		
Crops	Stage	Insect/Pest/Variety	Advisory
Summer Groundnut	Sowing	TAG-24, TG-37-A, TPG-41, GG-2, GJG-31	Carry out sowing of summer groundnut as temperature increases to 22-25° C and give seed treatment of Tebuconazole 1.25 gm/1 kg seed. Apply Single Super Phosphate 312 kg, Ammonium sulphate 125 kg and Castor cake 250 kg/ha OR DAP 108 kg, Gypsum 250 kg and castor cake 250 kg/ha as basal.
Summer Sesame	Preparation for sowing	G.Til-2, G.Til-3	Procure good quality seeds of sesame and carry out sowing of summer SESAME as minimum and maximum temperature rises up to 19 °C and 30°C, respectively and give seed treatment of vitavax (Carboxin 37.5% + Thiram 37.5%) or mencozeb or carbendazim 2-3 gm/1 kg seed. Apply Single Super Phosphate 156 kg and 55 kg urea/ha, OR

			DAP 54 kg and 33 kg urea/ha as basal.
Brinjal	Transplan-ting	Junagadh oblong, Junagadh long, Guj. Bringle-2	Keep planting distance 90 X 60 cm ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Lady finger	Sowing	Guj. Hy.-1 & 2	Keep planting distance 60 X 45 cm ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Ridge gourd	Sowing	Pusa nasdar, Jaipuri,	Keep planting distance 1.5 X 0.5 m ² and apply urea 27, single super phosphate 78 and murret of potash 21 kg/ha at sowing time.
Bottle gourd	Sowing	Junagadh tender long, Pusa summer prolific long	Keep planting distance 2 X 0.5 m ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Sponge guard	Sowing	Pusa chikni	Keep planting distance 1.5 X 0.5 m ² and apply urea 27, single super phosphate 78 and murret of potash 21 kg/ha at sowing time.
Bitter guard	Sowing	Priya, Pusa domosmi	Keep planting distance 1.0 X 1.0 m ² and apply urea 65, single super phosphate 375 and murret of potash 100 kg/ha at sowing time.
Cluster bean	Sowing	Pusa sadabahar, Pusa navbahar, IC-11388	Keep planting distance 60 X 30 cm ² and apply urea 55, single super phosphate 235 and murret of potash 63kg/ha at sowing time.
Water melon	Sowing	Sugar baby, Arka jyoti	Keep planting distance 2.0 X 1.0 m ² and apply urea 108, single super phosphate 312 and murret of potash 83 kg/ha at sowing time.
Garlic	Girthing	Thrips  Alternaria blight/Purple Blotch 	In present weather conditions attack of thrips has been reported. Make alternative sprays of imidaciliprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval. The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop. Make two spray of 27 gm Mencozeb or thiophanae methyl 12 gm/10 liters of water at 10 days interval.
Chick pea	Pod formation/ development	Monitor the crop regularly for prodenia and heliothes infstation	“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation cross ETL spray 3 ml SPINOSAD in 10 litres of water.

Wheat	Grain filling	-	Apply irrigation without stress.
Cumin	Grain formation/development	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray imidacloprid 5 ml/10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p> 
Coriander	Grain formation/development	Aphid, Thrips and Jassids	<p>Spray neem seed extract 5% and Thiomethoxam 5 gm/ 10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent powdery mildew infestation.</p>
Onion	Growth	<p>Thrips</p> <p>Purple blotch →</p> 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Spray Mencozeb 25 gm and hexaconazole 15 ml in 10 liters of water at 10 days interval.</p>
Brinjal & Tomato	Flowering/fruiting	<p>Sucking pest →</p> <p>& Fruit borer</p>	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Cabbage and cauliflower	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.
Chili	Flowering & fruiting	Thrips, →	In present weather conditions severe attack of thrips has been reported. Spray 3 ml spinosad in

		 <p>Wilting →</p> <p>Anthracnose →</p> 	<p>10 litres of water.</p> <p>Drenching of <i>Trichoderma viridi</i> 4 gm in 1 liter of water.</p> <p>Spray Mencozeb 27 gm in 10 liters of water</p>
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose. Put curtains on west side of poultry house.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.