



ગ્રામીણ કૃષિ મૌસમ સેવા

હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન

ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ

તા.03-09-201૯ થી તા.09-09-201૯



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કૃ.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિહવામાન બુલેટીન નં. ૨૭/૨૦૧૯

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી)	27.8
મહત્તમ તાપમાન (સે.ગ્રે)	35.2-37.3
લઘુત્તમ તાપમાન (સે.ગ્રે)	23.4-26.2
ભેજ(%) સવાર	82-87
ભેજ(%) બપોરબાદ	47-53
પવનની ગતિ (કિમી/કલાક)	6.3-13.5

હવામાન આગાહી: પત્રક-ક

ઉત્તરસૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.03-09-201૯ થી 09-09-201૯

કૃષિ સલાહ:

સામાન્ય	<p>➤ ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના</p> <p>રાજકોટ: જુલ્લામાં આગામી તા.03-09-201૯ થી તા.09-09-201૯ દરમિયાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. 03 થી 0૫-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૨૦ થી ૨૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૬-૩૮ ડીગ્રી સેલ્શિયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૭-૨૭ ડીગ્રી સેલ્શિયસ જેટલું રહેવાની સંભાવના છે.</p>					
	તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
	પેરામીટર	૦૩-૦૭-૧૯	૦૪-૦૭-૧૯	૦૫-૦૭-૧૯	૦૬-૦૭-૧૯	૦૭-૦૭-૧૯
	વરસાદ(મી.મી)	5	4	5	0	0
	મહત્તમતાપમાન (સે.ગ્રે)	38	37	36	36	36

લઘુતમતાપમાન (સે.ગ્રે)	27	27	27	27	27
વાદળનીસ્થિતિ(ઓકટા)	3	6	8	7	6
મહતમભેજ(%) સવાર	72	69	76	72	72
લઘુતમભેજ (%) બપોરબાદ	43	40	60	46	48
પવનનીગતિ(કિમી/કલાક)	20	25	20	22	25
પવનનીદિશા(ડીગ્રી)	263	249	236	235	235

મોરબી: જીલ્લામાં આગામી તા.૦૩-૦૭-૨૦૧૯ થી તા.૦૭-૦૭-૨૦૧૯ દરમ્યાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૦૩ થી ૦૭-જુલાઈ ના રોજ હળવો થી મધ્યમ વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૨ થી ૨૪ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન ૩૬-૩૮ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૭-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૩-૦૭-૧૯	૦૪-૦૭-૧૯	૦૫-૦૭-૧૯	૦૬-૦૭-૧૯	૦૭-૦૭-૧૯
વરસાદ(મી.મી)	5	20	4	10	11
મહતમતાપમાન (સે.ગ્રે)	38	37	36	36	36
લઘુતમતાપમાન (સે.ગ્રે)	27	27	27	27	27
વાદળનીસ્થિતિ(ઓકટા)	6	8	8	8	7
મહતમભેજ(%) સવાર	74	91	86	83	80
લઘુતમભેજ (%) બપોરબાદ	47	47	70	61	52
પવનનીગતિ(કિમી/કલાક)	22	24	12	18	17
પવનનીદિશા(ડીગ્રી)	259	235	226	219	222

જામનગર: જીલ્લામાં આગામી તા.૦૩-૦૭-૨૦૧૯ થી તા.૦૭-૦૭-૨૦૧૯ દરમ્યાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૦૫ અને ૦૭-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૭ થી ૨૮ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન ૩૬-૩૭ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૭-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૩-૦૭-૧૯	૦૪-૦૭-૧૯	૦૫-૦૭-૧૯	૦૬-૦૭-૧૯	૦૭-૦૭-૧૯
વરસાદ(મી.મી)	0	0	3	0	4
મહતમતાપમાન (સે.ગ્રે)	37	36	36	37	37
લઘુતમતાપમાન (સે.ગ્રે)	27	27	27	27	27
વાદળનીસ્થિતિ(ઓકટા)	4	6	7	7	7

મહત્તમભેજ(%) સવાર	78	77	81	79	78
લઘુત્તમભેજ (%) બપોરબાદ	62	59	68	63	64
પવનનીગતિ(કિમી/કલાક)	17	22	24	27	28
પવનનીદિશા(ડીગ્રી)	261	252	241	237	236

દેવભૂમિ દ્વારકા: જીલ્લામાં આગામી તા.૦૩-૦૭-૨૦૧૯ થી તા.૦૭-૦૭-૨૦૧૯ દરમિયાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૦૩ થી ૦૫ અને ૦૭-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૪ થી ૨૯ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૨-૩૩ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૮-૨૯ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૩-૦૭-૧૯	૦૪-૦૭-૧૯	૦૫-૦૭-૧૯	૦૬-૦૭-૧૯	૦૭-૦૭-૧૯
વરસાદ(મી.મી)	૩	૩	૨	૦	૩
મહત્તમતાપમાન (સે.ગ્રે)	૩૩	૩૨	૩૨	૩૨	૩૨
લઘુત્તમતાપમાન (સે.ગ્રે)	૨૯	૨૯	૨૯	૨૮	૨૮
વાદળનીસ્થિતિ(ઓકટા)	૫	૬	૭	૭	૭
મહત્તમભેજ(%) સવાર	૭૮	૭૭	૮૦	૮૦	૭૮
લઘુત્તમભેજ (%) બપોરબાદ	૬૨	૫૬	૬૨	૬૧	૬૩
પવનનીગતિ(કિમી/કલાક)	૧૪	૨૧	૨૫	૨૮	૨૯
પવનનીદિશા(ડીગ્રી)	૨૬૨	૨૫૫	૨૪૨	૨૩૭	૨૩૬

અમરેલી: જીલ્લામાં આગામી તા.૦૩-૦૭-૨૦૧૯ થી તા.૦૭-૦૭-૨૦૧૯ દરમિયાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૦૩-જુલાઈ ના રોજ અતિ ભારે અને ૦૪ થી ૦૭-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૫ થી ૨૪ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૫-૩૬ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૮-૨૮ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૩-૦૭-૧૯	૦૪-૦૭-૧૯	૦૫-૦૭-૧૯	૦૬-૦૭-૧૯	૦૭-૦૭-૧૯
વરસાદ(મી.મી)	૭૦	૧૦	૪	૭	૯
મહત્તમતાપમાન (સે.ગ્રે)	૩૬	૩૫	૩૫	૩૫	૩૬
લઘુત્તમતાપમાન (સે.ગ્રે)	૨૮	૨૮	૨૮	૨૮	૨૮
વાદળનીસ્થિતિ(ઓકટા)	૭	૮	૮	૬	૬
મહત્તમભેજ(%) સવાર	૭૬	૭૫	૮૧	૭૯	૭૮

લઘુતમભેજ(%) બપોરબાદ	52	50	68	55	55
પવનનીગતિ(કિમી/કલાક)	15	22	17	20	24
પવનનીદિશા(ડીગ્રી)	261	252	240	238	238



સુરેન્દ્રનગર: જીલ્લામાં આગામી તા.૦૩-૦૭-૨૦૧૯ થી તા.૦૭-૦૭-૨૦૧૯ દરમ્યાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૦૩ અને ૦૫-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૯ થી ૩૧ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૬-૩૮ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૭-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.



તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૩-૦૭-૧૯	૦૪-૦૭-૧૯	૦૫-૦૭-૧૯	૦૬-૦૭-૧૯	૦૭-૦૭-૧૯
વરસાદ(મી.મી)	5	7	5	0	0
મહત્તમતાપમાન (સે.ગ્રે)	38	37	36	36	36
લઘુત્તમતાપમાન (સે.ગ્રે)	27	27	27	27	27
વાદળનીસ્થિતિ(ઓકટા)	3	6	8	7	6
મહત્તમભેજ(%) સવાર	69	68	75	70	70
લઘુત્તમભેજ(%) બપોરબાદ	41	40	57	44	48
પવનનીગતિ(કિમી/કલાક)	19	31	19	22	23
પવનનીદિશા(ડીગ્રી)	268	250	232	231	230

- ૦૩ થી ૦૭-જુલાઈ -૨૦૧૯ હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. રાજકોટ જીલ્લામાં તા. ૦૩ થી ૦૫-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે., મોરબી જીલ્લામાં તા. ૦૩ થી ૦૭-જુલાઈ ના રોજ હળવો થી મધ્યમ વરસાદ થવાની શક્યતા છે., જામનગર જીલ્લામાં તા. ૦૫ અને ૦૭-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે., દેવભૂમિ દ્વારકા તા. ૦૩ થી ૦૫ અને ૦૭-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે., અમરેલી જીલ્લામાં તા. ૦૩-જુલાઈ ના રોજ અતિ ભારે અને ૦૪ થી ૦૭-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે અને સુરેન્દ્રનગર જીલ્લામાં તા. ૦૩ અને ૦૫-જુલાઈ ના રોજ હળવો વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની મોટાભાગે રહેવાની અને પવનની ઝડપ ૧૨ થી ૩૧ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૫-૩૮ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૭-૨૮ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.
- અનિયમિત વરસાદનું જોખમ ઓછું કરવા કપાસના પાકમાં તલ, મગફળી, સોયાબીન, કઠોળ જેવા પાકોનું આંતર પાક તરીકે વાવેતરનું આયોજન કરવું.
- જે વિસ્તારમાં વાવણી લાયક વરસાદ થયેલ હોય ત્યાં પસંદગીના પાકનું સમયસર વાવેતર પૂર્ણ કરવું.

	<ul style="list-style-type: none"> સમયસર વાવણી કાર્ય પૂરું કરવું. વાવેતર સાથે અથવા પછી અનુકૂળતા મુજબ સમાર અથવા બેલી/કર્યુ ચલાવવું જેથી જમીનમાં ભેજ જળવાય રહે અને ઉગાવો સારો થાય. રાજકોટ, અમરેલી, જામનગર અને સુરેન્દ્રનગર જીલ્લામાં આગળ-પાછળ છુટ્ટો છવાયો વરસાદ થયેલ હોય પાકની અવસ્થા મુજબ ખાલા પુરવા અને પારવણી કાર્ય કરવું. વધારાના વરસાદી પાણીના નિકાલની વ્યવસ્થા કરવી અને તેનો ચેકડેમ, તળાવ, ખેતતલાવડીમાં સંગ્રહ કરવો. 		
મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
મગફળી	વાવેતર / ઉગાવો	અર્ધ વેલડી જીજી-૨૦, જીજી-૨૨ ઉભડી -જીજી-૨, જીજી-૩૨, ટીએજી-૨૪, ટીએલજી-૪૫, ટીપીજી-૪૧, તીજી-૩૭-એ	<p>વાવણી લાયક વરસાદ થયેલ હોય ત્યાં વાવેતર કરવું.</p> <p>વાવેતર સમયે સિંગલ સુપર ફોસ્ફેટ ૧૫૬ કી.ગ્રા./હે., એમોનીયમ સલ્ફેટ ૬૨ કી.ગ્રા./હે., અને એરંડીનો ખોળ ૫૦૦ કી.ગ્રા./હે. અથવા ડી.એ.પી. ૫૪ કી.ગ્રા. અને જીપ્સમ ૫૦૦ કી.ગ્રા./હે. સાથે એરંડીનો ખોળ ૫૦૦ કી.ગ્રા./હે. પાયામાં આપવો.</p> <p>તેમજ ૨.૫ કી.ગ્રા./હે ટ્રાયકોડર્મા ફૂગનો પાઉડર ૫૦૦ કી.ગ્રા./હે એરંડીનાં ખોળ સાથે મિશ્ર કરી પાયામાં વાવણી સમયે આપવો.</p> <p>વાવતી વખતે બિયારણને થાયરમ અથવા મેન્કોઝેબ અથવા વાયટાવેક્ષ દવાનો ૨-૩ ગ્રામ/કી. બીજને પટ આપવો.</p> <p>મગફળીમાં સફેદ મુંડાનાં નિયંત્રણ માટે ક્લોરોપાયરીફોસ ૨૦ % ઈસી દવાનો ૨૫ મિલી ૧ કી.ગ્રા. બીજને વાવેતર પહેલા પટ આપવો.</p>
કપાસ	વાવેતર / ઉગાવો	કપાસ બોલગાર્ડ-૨ ની પ્રમાણિત જાતો જેવી કે વહેલી પાકતી જાતો : ગુજરાત શંકર-૮, એટીએમ, ડિસ્ટ કલાસ, ૭૫૭૬, વી-૩૩૩, જાદુ, જય, સરજુ, સુરક્ષા, બળવાન, ભક્તિ, ભદ્રા, આરસીએચ-૭૭૯	<p>વાવણી લાયક વરસાદ થયે સમયસર વાવેતર કરવું.</p> <p>કપાસના પાકમાં ઈયળોમાં બીટી જીન સામે પ્રતિકારક શક્તિ ના આવે તે માટે બિયારણના પેકેટ સાથે આપવામાં આવતું નોનબીટી (રેફ્યુઝી) બીજનું વાવેતર અવશ્ય કરવું.</p> <ul style="list-style-type: none"> અનિયમિત વરસાદનું જોખમ ઓછું કરવા કપાસના પાકમાં તલ, મગફળી, સોયાબીન, કઠોળ જેવા પાકોનું આંતર પાક તરીકે વાવેતરનું આયોજન કરવું. વાવણી પૂર્ણ થયેલ હોય ત્યાં સમયસર ખાલા પુરવા.

		મધ્યમ મોડી પાકતી જાતો : ગુજરાત શંકર-૬, એગ્રીટોપ-૪૪૪, એગ્રીટોપ-૭૭૭ , સોલાર-૭૬, મલ્લિકા,આરસીએ ચ-૬૫૯, અજીત- ૧૫૫	
તલ	વાવેતર / ઉગાવો	ગુજરાત તલ- ૨, ગુજરાત તલ- ૩, ગુજરાત તલ- ૫ વાવણી પૂર્ણ થયેલ હોય ત્યાં સમયસર પારવાણી કરવી.	
બાજરી	વાવેતર / ઉગાવો	જીએચબી-૫૫૮ જીએચબી-૫૩૮, જીએચબી-૫૭૭, જીએચબી-૭૧૯, જીએચબી-૭૩૨ અને જીએચબી-૭૪૪ વાવણી પૂર્ણ થયેલ હોય ત્યાં સમયસર પારવાણી કરવી.	
કઠોળ પાકો	વાવેતર / ઉગાવો	ચોમાસું કઠોળ પાકો માટે મગની ગુજરાત મગ-૪ , કે-૮૫૧, ગુજરાત આણંદ મગ-૫ અને અડદની ગુજરાત અડદ -૧, ૨ ચોળા માટે - ગુજરાત ચોળા -૧,૨,૪, પૂસા ફાલ્ગુની, તુવેર માટે બીડીએન-૨, ગુજરાત તુવેર -૧૦૦, ગુજરાત તુવેર હાઈબ્રીડ-૧, પૈકી પસંદગીની જાતોનું સારી ગુણવત્તાનું પ્રમાણિત બીજ સમયસર મેળવી લેવું. વાવણી પૂર્ણ થયેલ હોય ત્યાં સમયસર પારવાણી કરવી.	
ભીંડા	ફૂલ/ફળ	સફેમાખી, પાન કથીરી	થ્રીપ્સના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૪ મિલી અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. જો પાન કથીરીનો ઉપદ્રવ હોય તો છથીઓન ૩૦ મિલી પ્રતિ પંપ છંટકાવ કરવો. ભીંડાની વીણી બાદ પ્રતિ એકરે ૫-૭ કી.ગ્રા. યુરિઆ ખાતર આપવું.
શાકભાજી	ઘડું ઉછેર/રોપણી વાવેતર	મરચી,ટમેટી, રીંગણ, અને કોબીઝના ઘડું તૈયાર કરવા (ગ્રીન શેડ નેટમાં અને નેટની ઊંચાઈ ૬.૫ ફૂટ રાખવી) ગાદી ક્યારા બનાવી તેમાં બીજનું વાવેતર કરવું. જો ઘડું તૈયાર હોય તો મરચી,ટમેટી, રીંગણ, અને કોબીઝની વરસાદ થયે રોપણી કરવી. તુરિયા, કારેલા, દુધી, ગલકા,પરવળ, કાકડી, ગુવાર, ભીંડો, ચોળી, તુવેર, મૂળા,તાંદળજો, પાલકનું વાવેતર કરવું.	

તુરીયા	ફૂલ/ફળ	થ્રીપ્સ/ સફેદ માખી	ઉનાળુ શાકભાજીમાં સફેદમાખીના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ઇમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૪ મિલી અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
દૂધી			જો પાન કથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છાન્કાવ કરવો.
ગલકા			વેલાવાળા શાકભાજીમાં તળછારાના નિયંત્રણ માટે મેન્કોઝેબ ૪૦ ગ્રામ પ્રતિ પંપમાં મિશ્ર કરી છંટકાવ કરવો.
કારેલા			વેલાવાળા શાકભાજી પાકોમાં હળવા અને ટુંકા અંતરે પિયત આપી યોગ્ય ભેજ જાળવવો જેથી પરાગનયન પ્રક્રિયા સારી રીતે થવાથી ઉત્પાદન સારું મળે.
ગુવાર			
કાકડી		તળછારો	
રીંગણી અને ટમેટી	ફૂલ/ફળ	સફેદ માખી	<ul style="list-style-type: none"> ▶ લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો ટ્રાયઝોફોસ ૪૦ ઇ.સી. ૨૦ મી.લી. અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ▶ ફળ અને ડોકાની ઈયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને ક્લોરાનટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઇસી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ▶ રીંગણી અને ટમેટીના પાકમાં પાન કથીરીના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
મરચી	ફૂલ/ફળ	પાનની કુક્ક- થ્રીપ્સ→	<p>દ્રાયઝોફોસ ૪૦ ઇ.સી. ૨૦ મી.લી. અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અથવા પ્રોફેનોફોસ ૨૦ મી.લી.અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>વાયરસના કારણે પીળા થઈ કુક્કાય ગયેલ છોડ ઉપાડી તેનો જમીનમાં દાટી નાશ કરવો. ત્યાં સફેદ માખીથી ફેલાતો હોવાથી તેના નિયંત્રણ માટે હોસ્ટાથીઓન દવા ૨૦ મી.લી./૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>
		 <p>પાનની કુક્ક-</p> 	

		કથીરી	જો પાન કથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છાન્કાવ કરવો.
લીંબુ	ફળનો વિકાસ	પાન કથીરી  સાયલા અને થ્રીપ્સ	ઊંચા તાપમાનને કારણે લીંબુના બગીચામાં પાન કથીરીનો ઉપદ્રવ વધુ હોય તો તેના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી ૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. તેના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૫ મીલી અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. વધુ ઉપદ્રવ હોય તો દાયઝોફોસ ૪૦ ટકા ૨૦/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
		બદામી ટપકાં	બદામી ટપકાના નિયંત્રણ માટે કોપર ઓક્સીક્લોરાઈડ ૪૦ ગ્રામ અને ૧.૫ ગ્રામ સ્ટ્રેપ્ટોસાયક્લીનના બે છંટકાવ ૧૫ દિવસના ગળે કરવા.
		પાનકોરીયુ	પાનકોરીયુના નિયંત્રણ માટે ડીડીવીપી ૧૦ મી.લી. ૧૦ લિટર પાણીમાં મિશ્ર કરીને છાંટવી.
પશુ/ મરઘા 			ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મિ.લિ. એક લિટર પાણીમાં નાખીને છાંટવી. વિટામીન એ તથા ડી પુરકો આપવા. કરમિયાની દવા, ટી.બી. તથા જે.ડી ના રોગની પરખ કરાવવી. વરસાદના સમયે પશુઓને સુરક્ષીત જગ્યાએ રાખવા. ચોમાસા દરમિયાન જાનવરોને રહેઠાણની જગ્યાની વ્યવસ્થિત સફાઈ તેમજ માખીઓના ઉપદ્રવથી બચવા ફિનાઈલ છાંટવું. દુધાળા પ્રાણીઓને નિયમિત ૧ કિલ્લો ફીડ + ૫૦ ગ્રામ ખનીજ તત્વોનું મિશ્રણ પ્રતિ ૨ લિટર દુધની ઉપજના પ્રમાણમાં આપવું. તેમજ લીલું ઘાસ + સૂકું ઘાસ + ખનીજ તત્વો ખોરાકમાં આપવા. ખુલ્લામાં રાખેલ ઘાસસારો વરસાદથી બગડે નહિ તે માટે જરૂરી પગલા ભરવા.

જુકૃયુ/ટેકનિકલ ઓફિસર.(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૧૯
તરઘડિયા(રાજકોટ), તારીખ: ૦૨/૦૭/૨૦૧૯.

પ્રતિ,
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર
સુકી ખેતીસંશોધન કેન્દ્ર
જુનાગઢ કૃષિ નિવર્સીટી
તરઘડીયા



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :942756433



Agro met Advisory Service bulletin No.27-for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 03 to 07-July - 2019)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	27.8
Maximum temperature (°C)	35.2-37.3
Minimum temperature (°C)	23.4-26.2
Relative Humidity (%) I	82-87
Relative Humidity (%) II	47-53
Wind speed (Kmph)	6.3-13.5

Weather forecast until 08-30 hrs.of 07/07/2019

(1) RAJKOT DISTRICT:







	03/07/19	04/07/19	05/07/19	06/07/19	07/07/19
Rainfall (mm)	5	4	5	0	0
Max.Temp.(°C)	38	37	36	36	36
Min.Temp.(°C)	27	27	27	27	27
Total CC (octa)	3	6	8	7	6
RH-I (%)	72	69	76	72	72
RH-II (%)	43	40	60	46	48
Wind Speed (kmph)	20	25	20	22	25
Wind Direction(deg.)	263	249	236	235	235

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: hot, humid and generally cloudy to cloudy weatherforecasted for next five days. During 03 to 05-July light rainfall forecasted.• Temperature: Maximum and minimum temperature likely to be in range of 36-38°C and 27-27°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 69-76% &40-60, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 20-25 km/h, respectively.• Complete sowing of kharif crops in time where sufficient rainfall received.• Carry out Planking or harrowing after sowing to conserve moisture for better
-----------------------	---

	<p>germination, where sowing is over.</p> <ul style="list-style-type: none"> • Due to sporadic and isolated rainfall in Rajkot, Amreli, Jamnagar and Surendranagar districts sowing, gap filling and thinning operations are going on simultaneously. • Make provision to drain excess water from field, harvest and store it in reservoir. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Kharif groundnut	Germination/ sowing	Semi spreading, GG-20, GG-22 Bunch, GG- 2,GJG-31, GJG- 32, GG-5, TAG-24, TG-37 A, TPG-41, TLG-45,	Carry out sowing in time where sufficient rainfall received. Apply 156 kg/ha SSP, 62 kg/ha Ammonium sulphate and 500 kg/ha Castor cake as basal dose. Seed treatment with Dithane M-45/Captan/vitavax @ 2 grams per kg of seeds before sowing is advised. As preventive measure for white grub infestation in groundnut crop treat the seed with Chlorpyrifos 20 % EC OR Quinalfos 25 % EC @ 25 ml/kg seed at the time of sowing.
Cotton	Germination/ sowing	Early: Bg-II GCH-8, GTHH-49, ATM, Jay, First Class, 7576, V-333, Neo, Beejdhan-2, Sarju, Atom, Bhakti,Bhadra Medium Late: Bg-II GCH-6, RCH-659, Agritop-777, Agritop-444, Solar-76, Swapna, Mallika, Ajit-155	Procure good quality of seeds for kharif . Carry out sowing in time where sufficient rainfall received. Farmers are kindly advised to sow non-BT cotton seeds (Refuse) given with cotton BT-II seeds to prevent resistance in ball worms against BT-II technology. Apply 20 kg nitrogen/ha as basal for rainfed cotton. Adopt inter crop like sesame, groundnut, soybean, pulses and maize in cotton crop for risk management.
Sesame	Germination/ sowing	GT-2 or GT-3 variety.	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pearl millet	Germination/ sowing	GHB-558,577,538,719, 732,744	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pulses	Germination/ sowing	Farmers growing pulses are advised to select variety GM-4, K-851 & GAM-5for green gram, Guj. Adud-1 for black gram, Gujarat cowpea-1,2,4, PusaFalguni for cowpea , BDN-2, GT-100, GTH-1 for pigeon pea	

		and Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.	
Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
	Sowing	Farmers are advised for sowing of Cucurbits in rainy season. It is most suitable time to sow recommended hybrids/ varieties. Some popular hybrids are Bottle Gourd-Junagadh long tender , Pusa Naveen, PusaSamridhi Bitter Gourd- PusaVisesh, Pusa-2 mosumi, Pumpkin-PusaVisvash , PusaVikash Ridge Gourd-G.J.H-1 , PusaChikni etc. Faremrs are also advised for Sowing of ladyfinger, cluster bean, kidney bean, cowpea, tur, amranthus etc.	
Lady finger	Flowering/ fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract orneem seed oil 50 ml with withacetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised. After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.
Ridge gourd, Bottle gourd, Sponge guard, Bitter guard& cucurbitaceous crops	Flowering & fruiting	Thrips& white flies/ Mites Downy mildew	Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation as dry conditions may lead to poor pollination and thus drop in yield of the crop. In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract orneem seed oil 50 ml withacetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites. For the control of downy mildew spray mencozeb 40g/pump.
Cluster bean	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval

Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus→  Mites	<p>In present weather conditions attack of mites has been reported.To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water.In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advisedfor control the vector, when sky remains clear.</p> <p>Constant monitoring of the chilli crop against attack of mites is advised If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water</p>
Lemon	Fruit development	Mite  Psylla and thrips  Citrus canker Leaf minor	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p> <p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water</p> <p>Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.</p>
Livestock & poultry 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes.</p> <p>Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :942756433



Agro met Advisory Service bulletin No.-27 for the Morbi district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 03 to 07-July- 2019)

Weather forecast until 08-30 hrs. of 07/07/2019

(2) MORBI DISTRICT:







	03/07/19	04/07/19	05/07/19	06/07/19	07/07/19
Rainfall (mm)	5	20	4	10	11
Max.Temp.(°C)	38	37	36	36	36
Min.Temp.(°C)	27	27	27	27	27
Total CC (octa)	6	8	8	8	7
RH-I (%)	74	91	86	83	80
RH-II (%)	47	47	70	61	52
Wind Speed (kmph)	22	24	12	18	17
Wind Direction(deg.)	259	235	226	219	222

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: hot, humid and generally cloudy to cloudy weatherforecasted for next five days. During 03 to 07-July light to moderate rainfall forecasted.• Temperature: Maximum and minimum temperature likely to be in range of 36-38°C and 27-27°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 74-91% & 47-70, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 12-24 km/h, respectively.• Complete sowing of kharif crops in time where sufficient rainfall received.• Carry out Planking or harrowing after sowing to conserve moisture for better germination, where sowing is over.• Make provision to drain excess water from field, harvest and store it in reservoir.• Farmer are also advised to adopt cotton-sesame, cotton-groundnut and cotton-
-----------------------	--

maize inter cropping system as risk management.			
Crops	Stage	Insect/Pest/ Variety	Advisory
Kharif groundnut	Germination/On set of monsoon	Semi spreading, GG-20, GG-22 Bunch, GG- 2,GJG-31, GJG- 32, GG-5, TAG-24, TG-37 A, TPG-41, TLG-45,	Carry out sowing in time where sufficient rainfall received. Apply 156 kg/ha SSP, 62 kg/ha Ammonium sulphate and 500 kg/ha Castor cake as basal dose. Seed treatment with Dithane M-45/Captan/vitavax @ 2 grams per kg of seeds before sowing is advised. As preventive measure for white grub infestation in groundnut crop treat the seed with Chlorpyrifos 20 % EC OR Quinalfos 25 % EC @ 25 ml/kg seed at the time of sowing.
Cotton	Germination/ sowing	Early: Bg-II GCH- 8, GTHH-49, ATM, Jay, First Class, 7576, V- 333, Neo, Beejdhan-2, Sarju, Atom, Bhakti,Bhadra Medium Late: Bg- II GCH-6, RCH- 659, Agritop-777, Agritop-444, Solar-76, Swapna, Mallika, Ajit-155	Procure good quality of seeds for kharif . Carry out sowing in time where sufficient rainfall received. Farmers are kindly advised to sow non-BT cotton seeds (Refuse) given with cotton BT-II seeds to prevent resistance in ball worms against BT-II technology. Apply 20 kg nitrogen/ha as basal for rainfed cotton. Adopt inter crop like sesame, groundnut, soybean, pulses and maize in cotton crop for risk management.
Sesame	Germination/ sowing	GT-2 or GT-3 variety.	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pearl millet	Germination/ sowing	GHB- 558,577,538,719, 732,744	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pulses	Germination/ sowing	Farmers growing pulses are advised to select variety GM-4, K-851 & GAM-5for green gram, Guj. Adud-1 for black gram, Gujarat cowpea- 1,2,4, PusaFalguni for cowpea , BDN-2, GT-100, GTH-1 for pigeon pea and Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.	

Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
	Sowing	Farmers are advised for sowing of Cucurbits in rainy season. It is most suitable time to sow recommended hybrids/ varieties. Some popular hybrids are Bottle Gourd-Junagadh long tender , Pusa Naveen, PusaSamridhi Bitter Gourd- PusaVisesh, Pusa-2 mosumi, Pumpkin-PusaVisvash , PusaVikash Ridge Gourd-G.J.H-1 , PusaChikni etc. Farmers are also advised for Sowing of ladyfinger, cluster bean, kidney bean, cowpea, tur, amranthus.	
Lady finger	Flowering/ fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised. After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.
Ridge gourd, Bottle gourd, Sponge guard, Bitter guard & cucurbitaceous crops	Flowering & fruiting	Thrips & white flies/ Mites Downy mildew	Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation as dry conditions may lead to poor pollination and thus drop in yield of the crop. In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites. For the control of downy mildew spray mencozeb 40g/pump.
Cluster bean	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval

Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  Mites	<p>In present weather conditions attack of mites has been reported.To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water.In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advisedfor control the vector, when sky remains clear.</p> <p>Constant monitoring of the chilli crop against attack of mites is advised If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water</p>
Lemon	Fruit development	Mite  Psylla and thrips  Citrus canker Leaf minor	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p> <p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.</p>
Livestock & poultry 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of light rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-27 for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 03 to 07-July - 2019)

Weather forecast until 08-30 hrs of 07/07/2019

(3) JAMNAGAR DISTRICT:

	03/07/19	04/07/19	05/07/19	06/07/19	07/07/19
Rainfall (mm)	0	0	3	0	4
Max. Temp.(°C)	37	36	36	37	37
Min. Temp.(°C)	27	27	27	27	27
Total CC (octa)	4	6	7	7	7
RH-I (%)	78	77	81	79	78
RH-II (%)	62	59	68	63	64
Wind Speed (kmph)	17	22	24	27	28
Wind Direction(deg.)	261	252	241	237	236







Agro meteorological Advisories:-

General Advice

- **Rain fall:** hot, humid and partly cloudy to generally cloudy weather forecasted for next five days. During 05 and 07-July light rainfall forecasted.
- **Temperature:** Maximum and minimum temperature likely to be in range of 36-37°C and 27-27°C, respectively.
- **Relative Humidity:** The RH-I and RH-II is likely to be in the range of 77-81% & 59-68, respectively.
- **Wind Direction and Speed:** Direction and speed of wind is likely to be W, SW and 17-28 km/h, respectively.
- **Complete sowing of kharif crops in time where sufficient rainfall received.**
- **Carry out Planking or harrowing after sowing to conserve moisture for better germination, where sowing is over.**
- **Make provision to drain excess water from field, harvest and store it in reservoir.**
- **Farmer are also advised to adopt cotton-sesame, cotton-groundnut and cotton-**

maize inter cropping system as risk management.			
Crops	Stage	Insect/Pest/ Variety	Advisory
Kharif groundnut	Germination/On set of monsoon	Semi spreading, GG-20, GG-22 Bunch, GG- 2,GJG-31, GJG- 32, GG-5, TAG-24, TG-37 A, TPG-41, TLG-45,	Carry out sowing in time where sufficient rainfall received. Apply 156 kg/ha SSP, 62 kg/ha Ammonium sulphate and 500 kg/ha Castor cake as basal dose. Seed treatment with Dithane M-45/Captan/vitavax @ 2 grams per kg of seeds before sowing is advised. As preventive measure for white grub infestation in groundnut crop treat the seed with Chlorpyrifos 20 % EC OR Quinalfos 25 % EC @ 25 ml/kg seed at the time of sowing.
Cotton	Germination/ sowing	Early: Bg-II GCH- 8, GTHH-49, ATM, Jay, First Class, 7576, V- 333, Neo, Beejdhan-2, Sarju, Atom, Bhakti,Bhadra Medium Late: Bg- II GCH-6, RCH- 659, Agritop-777, Agritop-444, Solar-76, Swapna, Mallika, Ajit-155	Procure good quality of seeds for kharif . Carry out sowing in time where sufficient rainfall received. Farmers are kindly advised to sow non-BT cotton seeds (Refuse) given with cotton BT-II seeds to prevent resistance in ball worms against BT-II technology. Apply 20 kg nitrogen/ha as basal for rainfed cotton. Adopt inter crop like sesame, groundnut, soybean, pulses and maize in cotton crop for risk management.
Sesame	Germination/ sowing	GT-2 or GT-3 variety.	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pearl millet	Germination/ sowing	GHB- 558,577,538,719, 732,744	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pulses	Germination/ sowing	Farmers growing pulses are advised to select variety GM-4, K-851 & GAM-5for green gram, Guj. Adud-1 for black gram, Gujarat cowpea-1,2,4, PusaFalguni for cowpea , BDN-2, GT-100, GTH-1 for pigeon pea and Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.	

Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
	Sowing	Farmers are advised for sowing of Cucurbits in rainy season. It is most suitable time to sow recommended hybrids/ varieties. Some popular hybrids are Bottle Gourd-Junagadh long tender , Pusa Naveen, PusaSamridhi Bitter Gourd- PusaVisesh, Pusa-2 mosumi, Pumpkin-PusaVisvash , PusaVikash Ridge Gourd-G.J.H-1 , PusaChikni etc. Farmers are also advised for Sowing of ladyfinger, cluster bean, kidney bean, cowpea, tur, amranthus.	
Lady finger	Flowering/ fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised. After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.
Ridge gourd, Bottle gourd, Sponge guard, Bitter guard & cucurbitaceous crops	Flowering & fruiting	Thrips & white flies/ Mites Downy mildew	Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation as dry conditions may lead to poor pollination and thus drop in yield of the crop. In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites. For the control of downy mildew spray mencozeb 40g/pump.
Cluster bean	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval

Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  Mites	<p>In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear.</p> <p>Constant monitoring of the chilli crop against attack of mites is advised If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water</p>
Lemon	Fruit development	Mite  Psylla and thrips  Citrus canker Leaf minor	<p>Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water.</p> <p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water</p> <p>Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.</p>
Livestock & poultry 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of light rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-27 for the Devbhumi Dwarka district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 03 to 07-July - 2019)

Weather forecast until 08-30 hrs of 07/07/2019

(4) DEVBHUMI DWARKA DISTRICT:







	03/07/19	04/07/19	05/07/19	06/07/19	07/07/19
Rainfall (mm)	3	3	2	0	3
Max. Temp.(°C)	33	32	32	32	32
Min. Temp.(°C)	29	29	29	28	28
Total CC (octa)	5	6	7	7	7
RH-I (%)	78	77	80	80	78
RH-II (%)	62	56	62	61	63
Wind Speed (kmph)	14	21	25	28	29
Wind Direction(deg.)	262	255	242	237	236

Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none">• Rain fall: hot, humid and generally cloudy weather forecasted for next five days. During 03 to 05, and 07-July light rainfall forecasted.• Temperature: Maximum and minimum temperature likely to be in range of 32-33°C and 28-29°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 77-80% & 56-63, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 14-29 km/h, respectively.
-----------------------	--

	<ul style="list-style-type: none"> • Complete sowing of kharif crops in time where sufficient rainfall received. • Carry out Planking or harrowing after sowing to conserve moisture for better germination, where sowing is over. • Make provision to drain excess water from field, harvest and store it in reservoir. • Farmer are also advised to adopt cotton-sesame, cotton-groundnut and cotton-maize inter cropping system as risk management. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Kharif groundnut	Germination/On set of monsoon	Semi spreading, GG-20, GG-22 Bunch, GG-2, GJG-31, GJG-32, GG-5, TAG-24, TG-37 A, TPG-41, TLG-45,	Carry out sowing in time where sufficient rainfall received. Apply 156 kg/ha SSP, 62 kg/ha Ammonium sulphate and 500 kg/ha Castor cake as basal dose. Seed treatment with Dithane M-45/Captan/vitavax @ 2 grams per kg of seeds before sowing is advised. As preventive measure for white grub infestation in groundnut crop treat the seed with Chlorpyrifos 20 % EC OR Quinalfos 25 % EC @ 25 ml/kg seed at the time of sowing.
Cotton	Germination/sowing	Early: Bg-II GCH-8, GTHH-49, ATM, Jay, First Class, 7576, V-333, Neo, Beejdhan-2, Sarju, Atom, Bhakti, Bhadra Medium Late: Bg-II GCH-6, RCH-659, Agritop-777, Agritop-444, Solar-76, Swapna, Mallika, Ajit-155	Procure good quality of seeds for kharif . Carry out sowing in time where sufficient rainfall received. Farmers are kindly advised to sow non-BT cotton seeds (Refuse) given with cotton BT-II seeds to prevent resistance in boll worms against BT-II technology. Apply 20 kg nitrogen/ha as basal for rainfed cotton. Adopt inter crop like sesame, groundnut, soybean, pulses and maize in cotton crop for risk management.
Sesame	Germination/sowing	GT-2 or GT-3 variety.	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pearl millet	Germination/sowing	GHB-558,577,538,719, 732,744	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pulses	Germination/	Farmers growing pulses are advised to select variety GM-4, K-851 &	

	sowing	GAM-5 for green gram, Guj. Adud-1 for black gram, Gujarat cowpea-1,2,4, PusaFalguni for cowpea, BDN-2, GT-100, GTH-1 for pigeon pea and Procure good quality of seeds for kharif sowing & Carry out sowing in time where sufficient rainfall received.	
Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
	Sowing	Farmers are advised for sowing of Cucurbits in rainy season. It is most suitable time to sow recommended hybrids/ varieties. Some popular hybrids are Bottle Gourd-Junagadh long tender , Pusa Naveen, PusaSamridhi Bitter Gourd- PusaVisesh, Pusa-2 mosumi, Pumpkin- PusaVisvash, PusaVikash Ridge Gourd-G.J.H-1 , PusaChikni etc. Farmers are also advised for Sowing of ladyfinger, cluster bean, kidney bean, cowpea, tur, amaranthus.	
Lady finger	Flowering/ fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised. After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.
Ridge gourd, Bottle gourd, Sponge guard, Bitter guard & cucurbitaceous crops	Flowering & fruiting	Thrips & white flies/ Mites Downy mildew	Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation as dry conditions may lead to poor pollination and thus drop in yield of the crop. In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites. For the control of downy mildew spray mencozeb 40g/pump.
Cluster	Growing/fruiti	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20

bean	ng		% SP@ 5.0 gm / 10 lit of water at 15 days interval
Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  Mites	In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear. Constant monitoring of the chilli crop against attack of mites is advised If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water
Lemon	Fruit development	Mite  Psylla and thrips  Citrus canker Leaf minor	Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water. To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water. Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of light rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-27 for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 03 to 07-July - 2019)

Weather forecast until 08-30 hrs of 07/07/2019

(5) AMRELI DISTRICT:







	03/07/19	04/07/19	05/07/19	06/07/19	07/07/19
Rainfall (mm)	70	10	4	7	9
Max.Temp.(°C)	36	35	35	35	36
Min.Temp.(°C)	28	28	28	28	28
Total CC (octa)	7	8	8	6	6
RH-I (%)	76	75	81	79	78
RH-II (%)	52	50	68	55	55
Wind Speed (kmph)	15	22	17	20	24
Wind Direction(deg.)	261	252	240	238	238

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: hot, humid and generally cloudy to cloudy weather forecasted for next five days. During 03-July very heavy rainfall & 04 to 07-July light rainfall forecasted.• Temperature: Maximum and minimum temperature likely to be in range of 35-36°C and 28-28°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 75-81% & 50-68, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and
-----------------------	--

	<p>15-24 km/h, respectively.</p> <ul style="list-style-type: none"> • Complete sowing of kharif crops in time where sufficient rainfall received. • Carry out Planking or harrowing after sowing to conserve moisture for better germination, where sowing is over. • Make provision to drain excess water from field, harvest and store it in reservoir. • Farmer are also advised to adopt cotton-sesame, cotton-groundnut and cotton-maize inter cropping system as risk management. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Kharif groundnut	Germination/On set of monsoon	Semi spreading, GG-20, GG-22 Bunch, GG-2, GJG-31, GJG-32, GG-5, TAG-24, TG-37 A, TPG-41, TLG-45,	Carry out sowing in time where sufficient rainfall received. Apply 156 kg/ha SSP, 62 kg/ha Ammonium sulphate and 500 kg/ha Castor cake as basal dose. Seed treatment with Dithane M-45/Captan/vitavax @ 2 grams per kg of seeds before sowing is advised. As preventive measure for white grub infestation in groundnut crop treat the seed with Chlorpyrifos 20 % EC OR Quinalfos 25 % EC @ 25 ml/kg seed at the time of sowing.
Cotton	Germination/sowing	Early: Bg-II GCH-8, GTHH-49, ATM, Jay, First Class, 7576, V-333, Neo, Beejdhan-2, Sarju, Atom, Bhakti, Bhadra Medium Late: Bg-II GCH-6, RCH-659, Agritop-777, Agritop-444, Solar-76, Swapna, Mallika, Ajit-155	Procure good quality of seeds for kharif . Carry out sowing in time where sufficient rainfall received. Farmers are kindly advised to sow non-BT cotton seeds (Refuse) given with cotton BT-II seeds to prevent resistance in ball worms against BT-II technology. Apply 20 kg nitrogen/ha as basal for rainfed cotton. Adopt inter crop like sesame, groundnut, soybean, pulses and maize in cotton crop for risk management.
Sesame	Germination/sowing	GT-2 or GT-3 variety.	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pearl millet	Germination/sowing	GHB-558,577,538,719, 732,744	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.

Pulses	Germination/ sowing	Farmers growing pulses are advised to select variety GM-4, K-851 & GAM-5 for green gram, Guj. Adud-1 for black gram, Gujarat cowpea-1,2,4, PusaFalguni for cowpea, BDN-2, GT-100, GTH-1 for pigeon pea and Procure good quality of seeds for kharif sowing & Carry out sowing in time where sufficient rainfall received.	
Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
	Sowing	Farmers are advised for sowing of Cucurbits in rainy season. It is most suitable time to sow recommended hybrids/ varieties. Some popular hybrids are Bottle Gourd-Junagadh long tender , Pusa Naveen, Pusa Samridhi Bitter Gourd- Pusa Visesh, Pusa-2 mosumi, Pumpkin-Pusa Visvash , Pusa Vikash Ridge Gourd-G.J.H-1 , Pusa Chikni etc. Farmers are also advised for Sowing of ladyfinger, cluster bean, kidney bean, cowpea, tur, amaranthus.	
Lady finger	Flowering/ fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised. After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.
Ridge gourd, Bottle gourd, Sponge guard, Bitter guard & cucurbitaceous crops	Flowering & fruiting	Thrips & white flies/ Mites Downy mildew	Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation as dry conditions may lead to poor pollination and thus drop in yield of the crop. In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites. For the control of downy mildew spray mencozeb 40g/pump.

Cluster bean	Growing/fruitletting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
Brinjal & Tomato	Flowering/fruitletting	Sucking pest → & Fruit borer 	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
Chili	Flowering/fruitletting	Thrips →  Leaf curling Virus →  Mites	In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10 litre of water is advised for control the vector, when sky remains clear. Constant monitoring of the chilli crop against attack of mites is advised If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water
Lemon	Fruit development	Mite  Psylla and thrips  Citrus canker Leaf minor	Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water. To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water. Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of light rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Agro met Advisory Service bulletin No.-27 for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 03 to 07-July - 2019)

Weather forecast until 08-30 hrs of 07/07/2019

(6) SURENDRANAGAR DISTRICT:







	03/07/19	04/07/19	05/07/19	06/07/19	07/07/19
Rainfall (mm)	5	7	5	0	0
Max.Temp.(°C)	38	37	36	36	36
Min.Temp.(°C)	27	27	27	27	27
Total CC (octa)	3	6	8	7	6
RH-I (%)	69	68	75	70	70
RH-II (%)	41	40	57	44	48
Wind Speed (kmph)	19	31	19	22	23
Wind Direction(deg.)	268	250	232	231	230

Agro meteorological advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: hot, humid and generally cloudy to cloudy weatherforecasted for next five days. During 03 to 05-July light rainfall forecasted.• Temperature: Maximum and minimum temperature likely to be in range of 36-38°C and 27-27°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 68-75% &40-57, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be W, SW and 19-31 km/h, respectively.
-----------------------	--

	<ul style="list-style-type: none"> • Complete sowing of kharif crops in time where sufficient rainfall received. • Carry out Planking or harrowing after sowing to conserve moisture for better germination, where sowing is over. • Make provision to drain excess water from field, harvest and store it in reservoir. • Farmer are also advised to adopt cotton-sesame, cotton-groundnut and cotton-maize inter cropping system as risk management. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Kharif groundnut	Germination/On set of monsoon	Semi spreading, GG-20, GG-22 Bunch, GG-2, GJG-31, GJG-32, GG-5, TAG-24, TG-37 A, TPG-41, TLG-45,	Carry out sowing in time where sufficient rainfall received. Apply 156 kg/ha SSP, 62 kg/ha Ammonium sulphate and 500 kg/ha Castor cake as basal dose. Seed treatment with Dithane M-45/Captan/vitavax @ 2 grams per kg of seeds before sowing is advised. As preventive measure for white grub infestation in groundnut crop treat the seed with Chlorpyrifos 20 % EC OR Quinalfos 25 % EC @ 25 ml/kg seed at the time of sowing.
Cotton	Germination/sowing	Early: Bg-II GCH-8, GTHH-49, ATM, Jay, First Class, 7576, V-333, Neo, Beejdhan-2, Sarju, Atom, Bhakti, Bhadra Medium Late: Bg-II GCH-6, RCH-659, Agritop-777, Agritop-444, Solar-76, Swapna, Mallika, Ajit-155	Procure good quality of seeds for kharif . Carry out sowing in time where sufficient rainfall received. Farmers are kindly advised to sow non-BT cotton seeds (Refuse) given with cotton BT-II seeds to prevent resistance in boll worms against BT-II technology. Apply 20 kg nitrogen/ha as basal for rainfed cotton. Adopt inter crop like sesame, groundnut, soybean, pulses and maize in cotton crop for risk management.
Sesame	Germination/sowing	GT-2 or GT-3 variety.	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pearl millet	Germination/sowing	GHB-558,577,538,719, 732,744	Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.
Pulses	Germination/	Farmers growing pulses are advised to select variety GM-4, K-851 &	

	sowing	GAM-5for green gram, Guj. Adud-1 for black gram, Gujarat cowpea-1,2,4, PusaFalguni for cowpea , BDN-2, GT-100, GTH-1 for pigeon pea and Procure good quality of seeds for kharif sowing&Carry out sowing in time where sufficient rainfall received.	
Vegetables	Nursery /Transplanting	Farmers are advised for preparation of nursery for tomato, chilli, brinjal and early cauliflower (maturing in the month of September). It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. Nursery should be covered by net at the height of 6.5 feet for protection against sunlight. Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
	Sowing	Farmers are advised for sowing of Cucurbits in rainy season. It is most suitable time to sow recommended hybrids/ varieties. Some popular hybrids are Bottle Gourd-Junagadh long tender , Pusa Naveen, PusaSamridhi Bitter Gourd- PusaVisesh, Pusa-2 mosumi, Pumpkin -PusaVisvash, PusaVikash Ridge Gourd-G.J.H-1 , PusaChikni etc. Faremrns are also advised for Sowing of ladyfinger, cluster bean, kidney bean, cowpea, tur, amranthus.	
Lady finger	Flowering/ fruiting	White flies & mites	Spray 500 ml of 5 % neem seed extract orneem seed oil 50 ml with withacetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. Due to prevailing high temperature, light irrigation at short interval is advised. After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.
Ridge gourd, Bottle gourd, Sponge guard, Bitter guard& cucurbitaceous crops	Flowering & fruiting	Thrips& white flies/ Mites Downy mildew	Optimum moisture level should be maintained in cucurbitaceous crops by light and frequent irrigation as dry conditions may lead to poor pollination and thus drop in yield of the crop. In all vegetable crops thrips infestation observed above ETL. Spray 500 ml of 5 % neem seed extract orneem seed oil 50 ml withacetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 3 ml/10 lit of water for whiteflies control. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites. For the control of downy mildew spray mencozeb 40g/pump.
Cluster	Growing/fruitti	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20

bean	ng		% SP@ 5.0 gm / 10 lit of water at 15 days interval
Brinjal& Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
Chili	Flowering/ fruiting	Thrips →  Leaf curling Virus →  Mites	In present weather conditions attack of mites has been reported. To control mite infestation spray propargite 57 % EC 10 ml/10 litre of water. In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear. Constant monitoring of the chilli crop against attack of mites is advised If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water
Lemon	Fruit development	Mite  Psylla and thrips  Citrus canker Leaf minor	Due to high temperature mites infestation in lemon orchard observed. To control mite infestation spray propargite 57 % EC 10/10 liter of water. To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water. Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litre of water Spray Dichlorvosh 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of light rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.

