



ગ્રામીણ કૃષિ મૌસમ સેવા
હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ
તા.૦૮-૦૧-૨૦૨૦ થી તા.૧૨-૦૧-૨૦૨૦



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કૃ.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિ હવામાન બુલેટીન નં. ૮૧/૨૦૨૦

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી)	0.0
મહત્તમ તાપમાન (સે.ગ્રે)	23.8-31.1
લઘુત્તમ તાપમાન (સે.ગ્રે)	7.9-15.0
ભેજ(%) સવાર	64-90
ભેજ(%) બપોરબાદ	33-47
પવનની ગતિ (કિમી/કલાક)	3.0-6.3

હવામાન આગાહી: પત્રક-ક

ઉત્તરસૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.૦૮-૦૧-૨૦૨૦ થી ૧૨-૦૧-૨૦૨૦

કૃષિ સલાહ:

સામાન્ય	ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના					
	રાજકોટ: જીલ્લામાં આગામી તા.૦૮-૦૧-૨૦૨૦ થી તા.૧૨-૦૧-૨૦૨૦ દરમિયાન હવામાન સૂકું, ઠંડુ અને યોખ્ખું રહેશે. પવનની દિશા નૈઋત્ય, ઈશાન અને અગ્નિની રહેવાની અને પવનની ઝડપ ૭ થી ૧૯ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૨૮-૩૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૧૨-૧૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.					
	તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
	પેરામીટર	૦૮-૦૧-૨૦	૦૯-૦૧-૨૦	૧૦-૦૧-૨૦	૧૧-૦૧-૨૦	૧૨-૦૧-૨૦
	વરસાદ(મી.મી)	0	0	0	0	0
	મહત્તમતાપમાન (સે.ગ્રે)	31	29	28	29	30
	લઘુત્તમતાપમાન (સે.ગ્રે)	15	14	12	12	13
	વાદળનીસ્થિતિ(ઓક્ટા)	0	0	0	0	0
	મહત્તમભેજ(%) સવાર	59	63	47	39	58

લઘુતમભેજ (%) બપોરબાદ	37	36	21	22	28
પવનનીગતિ(કિમી/કલાક)	14	19	18	16	7
પવનનીદિશા(ડીગ્રી)	232	28	50	57	153

મોરબી : જીલ્લામાં આગામી તા.૦૮-૦૧-૨૦૨૦ થી તા.૧૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને મોટાભાગે ચોખ્ખું રહશે. પવનની દિશા અગ્નિ, ઈશાન અને પૂર્વની રહેવાની અને પવનની ઝડપ ૪ થી ૧૬ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૮-૩૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૧૨-૧૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૮-૦૧-૨૦	૦૯-૦૧-૨૦	૧૦-૦૧-૨૦	૧૧-૦૧-૨૦	૧૨-૦૧-૨૦
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	31	29	28	29	30
લઘુત્તમતાપમાન (સે.ગ્રે)	15	14	12	12	13
વાદળનીસ્થિતિ(ઓકટા)	1	0	0	0	0
મહત્તમભેજ(%) સવાર	66	60	61	44	62
લઘુત્તમભેજ (%) બપોરબાદ	37	31	21	23	33
પવનનીગતિ(કિમી/કલાક)	4	13	16	15	9
પવનનીદિશા(ડીગ્રી)	133	26	60	79	146

જામનગર: જીલ્લામાં આગામી તા.૦૮-૦૧-૨૦૨૦ થી તા.૧૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને ચોખ્ખું રહશે. પવનની દિશા ઉત્તર, ઈશાન અને પૂર્વની રહેવાની અને પવનની ઝડપ ૯ થી ૨૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૮-૩૦ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૧૪-૧૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૮-૦૧-૨૦	૦૯-૦૧-૨૦	૧૦-૦૧-૨૦	૧૧-૦૧-૨૦	૧૨-૦૧-૨૦
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	30	28	28	28	29
લઘુત્તમતાપમાન (સે.ગ્રે)	14	14	14	14	15
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	0	0
મહત્તમભેજ(%) સવાર	64	67	49	44	59
લઘુત્તમભેજ (%) બપોરબાદ	45	41	29	29	34
પવનનીગતિ(કિમી/કલાક)	20	25	24	23	9
પવનનીદિશા(ડીગ્રી)	9	33	47	53	82

દેવભૂમિ દ્વારકા: જીલ્લામાં આગામી તા.૦૮-૦૧-૨૦૧૯ થી તા.૧૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને ચોખ્ખું રહશે. પવનની દિશા ઉત્તર અને ઈશાનની રહેવાની અને પવનની ઝડપ ૭ થી ૨૪

કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **૨૫-૨૭** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન **૧૫-૧૬** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૮-૦૧-૨૦	૦૯-૦૧-૨૦	૧૦-૦૧-૨૦	૧૧-૦૧-૨૦	૧૨-૦૧-૨૦
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	27	25	25	25	26
લઘુત્તમતાપમાન (સે.ગ્રે)	15	15	15	15	16
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	0	0
મહત્તમભેજ(%) સવાર	64	66	49	45	62
લઘુત્તમભેજ (%) બપોરબાદ	45	42	29	26	30
પવનનીગતિ(કિમી/કલાક)	19	24	22	24	7
પવનનીદિશા(ડીગ્રી)	11	35	47	56	44

અમરેલી: જીલ્લામાં આગામી તા.૦૮-૦૧-૨૦૨૦ થી તા.૧૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને મોટાભાગે ચોખ્ખું રહશે. પવનની દિશા વાયવ્ય, ઉતર, ઈશાન અને પશ્ચિમની રહેવાની અને પવનની ઝડપ **૭ થી ૨૪ કીમી/કલાક** રહેની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **૨૮-૩૦** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન **૧૧-૧૪** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૮-૦૧-૨૦	૦૯-૦૧-૨૦	૧૦-૦૧-૨૦	૧૧-૦૧-૨૦	૧૨-૦૧-૨૦
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	30	28	29	28	29
લઘુત્તમતાપમાન (સે.ગ્રે)	14	13	11	11	12
વાદળનીસ્થિતિ(ઓકટા)	1	0	0	0	0
મહત્તમભેજ(%) સવાર	66	54	45	45	68
લઘુત્તમભેજ(%) બપોરબાદ	44	34	27	29	36
પવનનીગતિ(કિમી/કલાક)	12	24	22	17	7
પવનનીદિશા(ડીગ્રી)	298	19	35	42	265




સુરેન્દ્રનગર: જીલ્લામાં આગામી તા.૦૮-૦૧-૨૦૨૦ થી તા.૧૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને મોટાભાગે ચોખ્ખું રહશે. પવનની દિશા પશ્ચિમ, ઉતર, ઈશાન અને અગ્નિની રહેવાની અને પવનની ઝડપ **૭ થી ૨૦ કીમી/કલાક** રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **૨૮-૩૧** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન **૧૨-૧૫** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.






તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૮-૦૧-૨૦	૦૯-૦૧-૨૦	૧૦-૦૧-૨૦	૧૧-૦૧-૨૦	૧૨-૦૧-૨૦



વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	31	29	28	29	30
લઘુત્તમતાપમાન (સે.ગ્રે)	15	14	12	12	13
વાદળનીસ્થિતિ(ઓકટા)	1	0	0	0	0
મહત્તમભેજ(%) સવાર	58	64	51	41	58
લઘુત્તમભેજ(%) બપોરબાદ	39	37	24	25	33
પવનનીગતિ(કિમી/કલાક)	17	20	19	17	7
પવનનીદિશા(ડીગ્રી)	256	20	54	65	114


➤ ૦૮ થી ૧૨-જાન્યુઆરી-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને મોટાભાગે ચોખ્ખું રહશે. પવનની દિશા ઈશાન, અગ્નિ, ઉત્તર અને પશ્ચિમની મોટાભાગે રહેવાની અને પવનની ઝડપ ૪ થી ૨૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૫-૩૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૧૧-૧૬ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.



➤ શિયાળુ પાકોમાં નિદામણ કાર્ય સમયસર કરવું.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટરપાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
લસણ	વૃદ્ધિ	શ્રીપ્સ,  પાનની ટોચનો સુકારો 	<ul style="list-style-type: none"> ➤ લસણના પાકમાં શ્રીપ્સના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને પ્રોફેનોફોસ ૩૦ મિલી અથવા લેમ્બડા સાયહેલોથ્રીન ૧૦ મિલી દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ લસણના પાકમાં પાન પીળા પડતા અટકાવવા મેન્ડોઝેબ ૨૭ ગ્રામ અથવા કાર્બેન્ડાઝીમ ૧૦ ગ્રામ પ્રતિ ૧૦ લીટર પાણીમાં નાખીને છંટકાવ કરવો. ➤ જમીનની પ્રત મુજબ ૧૦-૧૨ દિવસે પિયત આપવું. યુરિઆ ખાતર ૫૪ કી.ગ્રા. પ્રતિ હેક્ટર વાવેતર બાદ ૩૦-૩૫ દિવસે આપવું. ➤ લસણ પીળું પડે નહિ તે માટે વધારે પડતું પિયત આપવું નહિ.
ચણા	વૃદ્ધિ/ફૂલ/પોપટા	લીલી અને લશ્કરી ઈયળ 	<p>ચણાના પાકમાં લીલી અને લશ્કરી ઈયળના નિયંત્રણ માટે પક્ષીઓને બેસવા માટે ' T ' આકારના બેલાખડા મુકવા. જો વધુ ઉપદ્રવ જણાય તો ક્વીનાલફોસ ૨૦ મિલી અથવા એમામેક્ટીન બેન્ઝોએટ ૬.૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો. તેમજ લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.</p> <p>પાનના સુકારાના નિયંત્રણ માટે ડાયથેન એમ ૪૫, ૨૭ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</p>
ધાણા	ફૂલ	શ્રીપ્સ અને મોલો	શ્રીપ્સ અને મોલોના ના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને

		રાખોડીના નિયંત્રણ	થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો. રાખોડીના નિયંત્રણ હેક્ઝાકોનાઝોલ ૧૫ મિલી દવા ૧૦ લીટર પાણીમાં મિશ્ર કરીને ૧૫ દિવસના અન્તરે છંટકાવ કરવો.
જીરું	ફૂલ	શ્રીપ્સ ચરમી અને રાખોડી  સુકારો 	જીરુંના પાકમાં શ્રીપ્સના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો. જીરુંના પાકમાં તેમજ ચરમીનો ઉપદ્રવ ન થાય તે માટે ડાયથેન એમ ૪૫, ૨૭ ગ્રામ અને હેક્ઝાકોનાઝોલ ૧૫ મિલી દવા વારાફરતી ૧૦ લીટર પાણીમાં મિશ્ર કરીને ૧૫ દિવસના અન્તરે છંટકાવ કરવો.  જીરુંના ઉભા પાકમાં સુકારાના નિયંત્રણ માટે પિયત આપવાનું બંધ કરવું. 
ઘઉં	ફૂટ/ગાભ	પહોળા પાન વાળા નિંદામણ 	ઘઉંમાં પહોળા પાન વાળા નિંદામણના નિયંત્રણ માટે વાવેતર બાદ ૩૦-૩૫ દિવસે ૨-૪-ડી ૨૦ મિલી અથવા મેટાસલ્ફ્યુરોન ૦.૮ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી.
રાયડો	ફૂલ	રાય વરુણા, ગુજરાત -૩ અને ૪	શ્રીપ્સ અને મોલોના ના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.
કપાસ	કપાસની વીણી	ગુલાબી ઇયળ ગુલાબી ઇયળનાં નિયંત્રણ માટે તેમજ પિયત આપવું નહિ. બીજો ફાલ લેવો નહિ. છેલ્લી વીણી પછી ઘેટા બકરા ચરાવવા. જેથી ખુલ્લા વગરના ગુલાબી ઇયળના નુકશાન વાળા જીંડવા અને અન્ય કચરો નાશ પામે. ત્યારબાદ કરાંઠી સળગાવવી નહિ પરંતુ રોટાવેટર/મોબાઇલ ચોપરથી નાના ટુકડા કરી જમીનમાં દાંટી દેવા જેથી જમીન સેન્દ્રિય તત્વનો ઉમેરો થવાથી સુંક્ષ્મ જીવોની સંખ્યા વધશે અને જમીનના સ્વાસ્થ્યમાં સુધારો થશે. અથવા કરાંઠી ભેગી કરી શ્રેદરથી નાના ટુકડા કરીને છાણ, યુરિયા, કમ્પોસ્ટ કલ્ચર ઉમેરી દેશી(સેન્દ્રિય) ખાતર બનાવવું.	

			
દિવેલા	વૃદ્ધિ/ફૂલ/ગડ	લીલી પોપટી, સફેદ માખી ગડની ઇયળ	<ul style="list-style-type: none"> ➤ સફેદ માખી અને લીલી પોપટીનો વધુ ઉપદ્રવ જોવા મળે છે. જેના નિયંત્રણ માટે એસીટામાપ્રોડ ૨૦ ટકા ૫ ગ્રામ અને ઈમીડાક્લોપ્રોડ ૩૦.૫ % ૩ મિલી સાથે લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ ગડની ઇયળના નિયંત્રણ માટે ક્લોરાનટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઈસી ૭ મિલી ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
ડુંગળી	રોપણી/વૃદ્ધિ વિકાસ	શ્રીપ્સ, પાનનો સુકારો જાંબલી ઘાબા 	<ul style="list-style-type: none"> ➤ શ્રીપ્સના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને પ્રોફેનોફોસ ૩૦ મિલી અથવા લેમ્બડા સાયહેલોથ્રીન ૧૫ મિલી દવાને ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ ડુંગળીના પાકમાં જાંબલી ઘાબાના રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલર ૨૭ ગ્રામ અથવા ફોઝેટાઇલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા. ➤ નાઈટ્રોજન યુક્ત એમોનિયમ સલ્ફેટ ખાતરનો પુરતી ડોઝ ૩૦ કીલો ગ્રામ પ્રતિ હે. આપવો.
તુવેર	ફૂલ/શીંગ/દાણા ભરાવવા	શીંગ માખી	ક્વીનાલફોસ ૨૦ મિલી અથવા ડીડીવીપી ૫ મિલી ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. તેમજ લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લિટર પાણીમાં કરીને છંટકાવ કરવો.
કોબી અને ફલાવર	વૃદ્ધિ/દડા	હિરા ફૂદુ	કોબી અને ફલાવરમાં હિરાફૂદાના નિયંત્રણ માટે સ્પીનોસાદ ૩ મિલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
રીંગણી અને ટમેટી	ફૂલ/ફળ	તડતડીયા અને સફેદ માખી ડુંખ અને → ફળ કોરી ખાનાર ઇયળ પાન કથીરી	<ul style="list-style-type: none"> ➤ તડતડીયા અને સફેદ માખીના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ડાયફેનથાયુરોન ૫૦ ટકા વે.પા. ૧૬ ગ્રામ અથવા ટ્રાયઝોફોસ ૪૦ ઈસી ૨૫ મિ.લી. દવાને ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ ફળ અને ડોકાની ઇયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને ક્લોરાનટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઈસી મિલી ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ જો પાનકથીરીનો ઉપદ્રવ હોય તો પ્રોપરગાઈટ ૫૭ ઈ.સી. ૧૫ મિલી પ્રતિ પંપ છંટકાવ કરવો.
મરચી	વૃદ્ધિ/ફૂલ/ફળ	શ્રીપ્સ, સફેદ માખી	<ul style="list-style-type: none"> ➤ પ્રોફેનોફોસ ૨૦ મીલી અને લીંબોળીનું તેલ ૫૦ મિલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી ૧૦ લી.

		 <p>કાલવ્રણ →</p>	<p>પાણીમાં મિશ્ર કરી છંટકાવ કરવો. થ્રીપ્સનો વધુ ઉપદ્રવ હોય તો સ્પીનોસાદ અથવા સ્પીનેટોરમ દવા ૫ મિલી. પ્રતિ પંપ છાંટવી.</p> <p>➤ કાલવ્રણ રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.</p>
લીંબુ	ફળનો વિકાસ	<p>પાન કથીરી</p>  <p>સાયલા અને થ્રીપ્સ</p> <p>બદામી ટપકાં</p> <p>પાનકોરીયુ</p>	<p>લીંબુના બગીચામાં પાન કથીરીનો ઉપદ્રવ વધુ હોય તો તેના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>લીંબુમાં સાયલા અને થ્રીપ્સ તેના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૫ મીલી અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. વધુ ઉપદ્રવ હોય તો ટ્રાયઝોફોસ ૪૦ ટકા ૨૦/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>  <p>બદામી ટપકાના નિયંત્રણ માટે કોપર ઓક્ઝીક્લોરાઈડ ૪૦ ગ્રામ અને ૧.૫ ગ્રામ સ્ટ્રેપ્ટોસાયક્લીનના બે છંટકાવ ૧૫ દિવસના ગળે કરવા. પાનકોરીયુના નિયંત્રણ માટે ડીડીવીપી ૧૦ મી.લી. ૧૦ લિટર પાણીમાં મિશ્ર કરીને છાંટવી.</p>
આંબો	ફૂલ	<p>મધીયો અને મગીયાની ઈયળ</p> <p>ભુકીછારો/રાખોડી</p>	<p>આંબામાં મધીયો અને મગીયાની ઈયળના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૫ મિલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>  <p>આંબામાં ભુકીછારો/રાખોડીના નિયંત્રણ માટે હેક્ઝાકોનાઝોલ ૧૫</p>

		<p>મિલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> 
<p>પશુ/ મરઘા</p> 	<p>પ્રજીવથી થતા રોગોની ચકાસણી કરાવવી. ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મી.લી. ૧ લીટર પાણીમાં નાખીને છાંટવી. વિટામીન એ તથા ડી પુરકો આપવા. કરમિયાની દવા આપવી તથા ઉછરતા પશુઓમાં બૃસેલ્વા કાફવુંડનું રસીકરણ કરાવવું. પશુને ૬ થી ૮ કી.ગ્રા. સુકો અને ૧૫ થી ૨૦ કી.ગ્રા. લીલો ચારો આપવો. દુધાળા પશુઓને પ્રતિ લીટર દુધની ઊપજ હોય તેમને નિયમિત ૧ કી.ગ્રા. ફીડ + ૫૦ ગ્રામ ખનીજ તત્ત્વોનું મિશ્રણ આપવું. દુધાળા પશુઓને હુંફાળું ગરમ પાણી પીવા માટે આપવું. તેમજ માખીઓના ઉપદ્રવથી બચવા ફીનાઈલ છાંટવું.</p>	

જુફ્યુ/ટેકનિકલ ઓફિસર.(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૨૦
તરઘડિયા(રાજકોટ), તારીખ: ૦૭/૦૧/૨૦૨૦.

પ્રતિ,
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર
સુકી ખેતીસંશોધન કેન્દ્ર
જુનાગઢ કૃષિ નિવર્સિટી
તરઘડીયા



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94269 38235



Agro met Advisory Service bulletin No.81-for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 08 to 12-January- 2020)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	0.0
Maximum temperature (°C)	23.8-31.1
Minimum temperature (°C)	7.9-15.0
Relative Humidity (%) I	64-90
Relative Humidity (%) II	33-47
Wind speed (Kmph)	3.0-6.3




Weather forecast until 08-30 hrs. of 12/01/2020





(1) RAJKOT DISTRICT:




	08/01/20	09/01/20	10/01/20	11/01/20	12/01/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	31	29	28	29	30
Min.Temp.(°C)	15	14	12	12	13
Total CC (octa)	0	0	0	0	0
RH-I (%)	59	63	47	39	58
RH-II (%)	37	36	21	22	28
Wind Speed (kmph)	14	19	18	16	7
Wind Direction(deg.)	232	28	50	57	153







Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 28-31°C and 12-15°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 39-63% & 21-37, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be SW, NE, SE and 7-19 km/h, respectively. • Carry out weeding as per need in rabi crops. 		
Crops	Stage	Insect/Pest/Variety	Advisory

<p>Garlic</p>	<p>Vegetative growth</p>	<p>Thrips and</p>  <p>leaf blight (yellowing)</p> 	<p>For thrips control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
<p>Chick pea</p>	<p>Flowering/pod formation</p>	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.

<p>Cumin</p>	<p>Growth/Flowering</p>	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
<p>Coriander</p>	<p>Flowering</p>	<p>Thrips and Jassids</p>	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>
<p>Wheat</p>	<p>Flowering</p>	<p>-</p>	<p>Apply irrigation at critical stage without stress.</p>
<p>Cotton</p>	<p>Picking & pulling of cotton stalk</p>	<p>Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time.</p> <p>Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by thedoin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.</p>	

			
Castor	Flowering/ capsule formation	Jassids& white fly→ Pod borer	<p>Spray Acetamiprid 20 % SP@ 5.0 gmORImidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water.</p> <p>For the control of pod borer spray chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p>
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	Thrips Leaf blight→ 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Make alternate spray of Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.</p>
Brinjal& Tomato	Flowering/ fruiting	<p>Sucking pest →</p> <p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infetation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>

<p>Chili</p>	<p>Flowering & fruiting</p>	<p>Thrips → </p> <p>Anthracnose → </p>	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
<p>Lemon</p>	<p>Flowering and fruit formation</p>	<p>Psylla and thrips</p> <p>Citrus canker</p> <p>Leaf minor</p>	<p></p> <p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
<p>Mango</p>	<p>Flowering</p>	<p>Blossom midge and Leaf hopper → </p> <p>Powdery Mildew → </p>	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p> <p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20</p>		

	kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.
--	---

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-81 for the Morbi district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 08 to 12-January- 2020)







Weather forecast until 08-30 hrs. Of 12/01/2020



(2) MORBI DISTRICT:






	08/01/20	09/01/20	10/01/20	11/01/20	12/01/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	31	29	28	29	30
Min. Temp.(°C)	15	14	12	12	13
Total CC (octa)	1	0	0	0	0
RH-I (%)	66	60	61	44	62
RH-II (%)	37	31	21	23	33
Wind Speed (kmph)	4	13	16	15	9
Wind Direction(deg.)	133	26	60	79	146



Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and Mainly clear to clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 28-31°C and 12-15°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 44-66% & 21-37, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be SE, NE, E and 4-16 km/h, respectively. • Carry out weeding as per need in rabi crops. 		
Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Vegetative growth	Thrips and  leaf blight (yellowing)	For thrips control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.

			<p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
Chick pea	Flowering/pod formation	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.
Cumin	Growth/Flowering	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 

Coriander	Flowering	Thrips and Jassids	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water. Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.
Wheat	Flowering	-	Apply irrigation at critical stage without stress.
Cotton	Picking & pulling of cotton stalk	Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time. Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by thedoin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.	
Castor	Flowering/ capsule formation	Jassids& white fly→ Pod borer	Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water. For the control of pod borer spray chlorantraniliprole18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.

<p>Brinjal& Tomato</p>	<p>Flowering/ fruiting</p>	<p>Sucking pest →</p> <p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>
<p>Chili</p>	<p>Flowering & fruiting</p>	<p>Thrips →</p>  <p>Anthracnose →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
<p>Lemon</p>	<p>Flowering and fruit formation</p>	<p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p>	<p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
<p>Mango</p>	<p>Flowering</p>	<p>Blossom midge and Leaf hopper →</p> 	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p>

		<p>Powdery Mildew→</p> 	<p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-81 for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 08 to 12-January-2020)







Weather forecast until 08-30 hrs. of 12/01/2020



(3) JAMNAGAR DISTRICT:






	08/01/20	09/01/20	10/01/20	11/01/20	12/01/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	30	28	28	28	29
Min. Temp.(°C)	14	14	14	14	15
Total CC (octa)	0	0	0	0	0
RH-I (%)	64	67	49	44	59
RH-II (%)	45	41	29	29	34
Wind Speed (kmph)	20	25	24	23	9
Wind Direction(deg.)	9	33	47	53	82



Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 28-30°C and 14-15°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 44-67% & 29-45, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be N, NE, E and 9-25 km/h, respectively. • Carry out weeding as per need in rabi crops. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Garlic	Vegetative growth	Thrips and  leaf blight (yellowing)	For thrips control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.

			<p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
Chick pea	Flowering/pod formation	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.
Cumin	Growth/Flowering	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 

Coriander	Flowering	Thrips and Jassids	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water. Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.
Wheat	Flowering	-	Apply irrigation at critical stage without stress.
Cotton	Picking & pulling of cotton stalk	Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time. Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by thedoin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.	
Castor	Flowering/ capsule formation	Jassids& white fly→ Pod borer	Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water. For the control of pod borer spray chlorantraniliprole18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.

<p>Brinjal& Tomato</p>	<p>Flowering/ fruiting</p>	<p>Sucking pest →</p> <p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>
<p>Chili</p>	<p>Flowering & fruiting</p>	<p>Thrips →</p>  <p>Anthracnose →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
<p>Lemon</p>	<p>Flowering and fruit formation</p>	<p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p>	<p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
<p>Mango</p>	<p>Flowering</p>	<p>Blossom midge and Leaf hopper →</p> 	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p>

		<p>Powdery Mildew →</p> 	<p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-81 for the Devbhumi Dwarka district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 08 to 12-January-2020)







Weather forecast until 08-30 hrs. of 12/01/2020



(4) DEVBHUMI DWARKA DISTRICT:






	08/01/20	09/01/20	10/01/20	11/01/20	12/01/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	27	25	25	25	26
Min. Temp.(°C)	15	15	15	15	16
Total CC (octa)	0	0	0	0	0
RH-I (%)	64	66	49	45	62
RH-II (%)	45	42	29	26	30
Wind Speed (kmph)	19	24	22	24	7
Wind Direction(deg.)	11	35	47	56	44



Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 25-27°C and 15-16°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 45-66% & 26-45, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be N, NE and 7-24 km/h, respectively. • Carry out weeding as per need in rabi crops. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Garlic	Vegetative growth	Thrips and  leaf blight (yellowing)	For thrips control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.

			<p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
Chick pea	Flowering/pod formation	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.
Cumin	Growth/Flowering	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 

Coriander	Flowering	Thrips and Jassids	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water. Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.
Wheat	Flowering	-	Apply irrigation at critical stage without stress.
Cotton	Picking & pulling of cotton stalk	Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time. Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by thedoin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.	
Castor	Flowering/ capsule formation	Jassids& white fly→ Pod borer	Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water. For the control of pod borer spray chlorantraniliprole18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.

<p>Brinjal& Tomato</p>	<p>Flowering/ fruiting</p>	<p>Sucking pest →</p> <p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>
<p>Chili</p>	<p>Flowering & fruiting</p>	<p>Thrips →</p>  <p>Anthracnose →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
<p>Lemon</p>	<p>Flowering and fruit formation</p>	<p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p>	<p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
<p>Mango</p>	<p>Flowering</p>	<p>Blossom midge and Leaf hopper →</p> 	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p>

		<p>Powdery Mildew→</p> 	<p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-81 for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 08 to 12-January-2020)







Weather forecast until 08-30 hrs. of 12/01/2020



(5) AMRELI DISTRICT:






	08/01/20	09/01/20	10/01/20	11/01/20	12/01/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	30	28	29	28	29
Min.Temp.(°C)	14	13	11	11	12
Total CC (octa)	1	0	0	0	0
RH-I (%)	66	54	45	45	68
RH-II (%)	44	34	27	29	36
Wind Speed (kmph)	12	24	22	17	7
Wind Direction(deg.)	298	19	35	42	265



Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and Mainly clear to clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 28-30°C and 11-14°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 45-68% & 27-44, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be NW, N, NE, W and 7-24 km/h, respectively. • Carry out weeding as per need in rabi crops. 		
Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Vegetative growth	Thrips and  leaf blight (yellowing)	For thrips control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.

			<p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
Chick pea	Flowering/pod formation	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.
Cumin	Growth/Flowering	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 

Coriander	Flowering	Thrips and Jassids	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water. Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.
Wheat	Flowering	-	Apply irrigation at critical stage without stress.
Cotton	Picking & pulling of cotton stalk	Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time. Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by thedoin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.	
Castor	Flowering/ capsule formation	Jassids& white fly→ Pod borer	Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water. For the control of pod borer spray chlorantraniliprole18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.

<p>Brinjal& Tomato</p>	<p>Flowering/ fruiting</p>	<p>Sucking pest →</p> <p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>
<p>Chili</p>	<p>Flowering & fruiting</p>	<p>Thrips →</p>  <p>Anthracnose →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
<p>Lemon</p>	<p>Flowering and fruit formation</p>	<p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p>	<p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptomycin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
<p>Mango</p>	<p>Flowering</p>	<p>Blossom midge and Leaf hopper →</p> 	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p>

		<p>Powdery Mildew →</p> 	<p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:9426938235



Agro met Advisory Service bulletin No.-81 for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 08 to 12-January-2020)



Weather forecast until 08-30 hrs of 12/01/2020





(6) SURENDRANAGAR DISTRICT:




	08/01/20	09/01/20	10/01/20	11/01/20	12/01/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	31	29	28	29	30
Min.Temp.(°C)	15	14	12	12	13
Total CC (octa)	1	0	0	0	0
RH-I (%)	58	64	51	41	58
RH-II (%)	39	37	24	25	33
Wind Speed (kmph)	17	20	19	17	7
Wind Direction(deg.)	256	20	54	65	114







Agro meteorological advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and Mainly clear to clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 28-31°C and 12-15°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 41-64% & 24-39, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be W, N, NE, SE and 7-20 km/h, respectively. • Carry out weeding as per need in rabi crops. 		
Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Vegetative growth	Thrips and  leaf blight (yellowing)	For thrips control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.

			<p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
<p>Chick pea</p>	<p>Flowering/pod formation</p>	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.

<p>Cumin</p>	<p>Growth/Flowering</p>	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
<p>Coriander</p>	<p>Flowering</p>	<p>Thrips and Jassids</p>	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>
<p>Wheat</p>	<p>Flowering</p>	<p>-</p>	<p>Apply irrigation at critical stage without stress.</p>
<p>Cotton</p>	<p>Picking & pulling of cotton stalk</p>	<p>Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time.</p> <p>Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by thedoin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.</p>	

			
Castor	Flowering/ capsule formation	Jassids & white fly → Pod borer	<p>Spray Acetamiprid 20 % SP @ 5.0 gm OR Imidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 ml / 10 lit of water ml/10 lit of water.</p> <p>For the control of pod borer spray chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p>
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	Thrips Leaf blight → 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Make alternate spray of Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.</p>
Brinjal & Tomato	Flowering/ fruiting	<p>Sucking pest →</p> <p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>

<p>Chili</p>	<p>Flowering & fruiting</p>	<p>Thrips → </p> <p>Anthracnose → </p>	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
<p>Lemon</p>	<p>Flowering and fruit formation</p>	<p>Psylla and thrips</p> <p>Citrus canker</p> <p>Leaf minor</p>	<p></p> <p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
<p>Mango</p>	<p>Flowering</p>	<p>Blossom midge and Leaf hopper → </p> <p>Powdery Mildew → </p>	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p> <p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20</p>		

	kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.
--	---

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.