



ग्रामीण कृषि मौसम सेवा  
हवामान आगाही आधारित कृषिसलाह बुलेटीन  
उत्तर सौराष्ट्रभेत् आबोहवाकीय विभाग  
ता.१४-०९-२०१९ थी ता.१८-०९-२०१९



(ग्रामीण कृषि मौसम सेवा, सूकी भेती संशोधन केन्द्र, ज.कृ.यु., तरघडिया अने भारत मौसम विभाग द्वारा संपन्न)

कृषिहवामान बुलेटीन नं. ४८/२०१९

पाछला अहवाडीयानुं राजकोटनुं हवामान

हवामान परिबणो	
वरसाह(मी.मी)	52.5
महतम तापमान (से.ग्रे)	28.0-32.5
लघुतम तापमान (से.ग्रे)	22.7-24.1
लेज(%) सवार	85-98
लेज(%) बपोरबाह	79-100
पवननी गति (किमी/कलाक)	2.3-6.0

हवामान आगाही: पत्रक-६

उत्तरसौराष्ट्र भेत् आबोहवाकीय विभाग-हवामान आगाही

ता.१४-०९-२०१९ थी १८-०९-२०१९

कृषि सलाह:

सामान्य	<ul style="list-style-type: none"><li>भारत सरकारश्रीना हवामान भाता तरङ्गथी मणेल हवामान माहिती अनुसार अत्रेना ग्रामीणकृषि मौसम सेवा विभाग,जुनागढ कृषि युनिवर्सिटी, तरघडीया तरङ्गथी जषाववामां आवे छे के, उत्तर सौराष्ट्र भेत् आबोहवाकीय विस्तारना</li></ul>					
	<p>राजकोट: जुल्लामां आगामी ता.१४-०९-२०१९ थी ता.१८-०९-२०१९ दरभ्यान हवामान गरम, लेजवाणुं अने वाहणछायुं रहशे. ता. १४ ना रोजे लारे अने ता. १५ ना रोजे हणवो वरसाह थवानी शक्यता छे. पवननी दिशा पश्चिम अने नैऋत्यनी रहेवानी अने पवननी ञऽप ११ थी १९ कीमी/कलाक रहेवानी शक्यता छे. आ समय गाणामां महतम तापमान दिवस दरभ्यान ३१-३२ डीग्री सेल्सीयस जेटलुं तेमज लघुतम तापमान रात्री दरभ्यान २४-२४ डीग्री सेल्सीयस जेटलुं रहेवानी संभावना छे.</p>					
	तारीभ	दिवस-१	दिवस-२	दिवस-३	दिवस-४	दिवस-५
	पेरामीटर	१४-०९-१९	१५-०९-१९	१६-०९-१९	१७-०९-१९	१८-०९-१९
	वरसाह(मी.मी)	43	6	0	0	0
	महतमतापमान (से.ग्रे)	32	31	32	32	32
	लघुतमतापमान (से.ग्रे)	24	24	24	24	24
	वाहणनीस्थिति(ओकटा)	8	8	8	8	7



મહત્તમભેજ(%) સવાર	89	90	90	88	91
લઘુત્તમભેજ (%) બપોરબાદ	68	75	75	68	67
પવનનીગતિ(કિમી/કલાક)	18	18	18	19	11
પવનનીદિશા(ડીગ્રી)	262	252	248	251	237

**અમરેલી:** જીલ્લામાં આગામી તા.૧૪-૦૯-૨૦૧૯ થી તા.૧૮-૦૯-૨૦૧૯ દરમિયાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૧૪ ના રોજ ભારેથી અતિ ભારે અને તા. ૧૫ થી ૧૭ ના રોજ હળવો થી મધ્યમ વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ ૧૦ થી ૧૭ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૨૯-૩૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૪-૦૯-૧૯	૧૫-૦૯-૧૯	૧૬-૦૯-૧૯	૧૭-૦૯-૧૯	૧૮-૦૯-૧૯
વરસાદ(મી.મી)	70	16	17	10	0
મહત્તમતાપમાન (સે.ગ્રે)	30	29	29	31	31
લઘુત્તમતાપમાન (સે.ગ્રે)	24	24	25	25	25
વાદળનીસ્થિતિ(ઓકટા)	8	8	8	7	7
મહત્તમભેજ(%) સવાર	92	92	92	90	92
લઘુત્તમભેજ(%) બપોરબાદ	76	79	78	72	71
પવનનીગતિ(કિમી/કલાક)	17	12	15	16	10
પવનનીદિશા(ડીગ્રી)	288	262	253	255	254

**સુરેન્દ્રનગર:** જીલ્લામાં આગામી તા.૧૪-૦૯-૨૦૧૯ થી તા.૧૮-૦૯-૨૦૧૯ દરમિયાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૧૪ થી ૧૬ ના રોજ હળવો થી મધ્યમ વરસાદ થવાની શક્યતા છે. પવનની દિશા નૈઋત્ય અને પશ્ચિમની રહેવાની અને પવનની ઝડપ ૧૨ થી ૨૦ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૧-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૪-૨૪ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૪-૦૯-૧૯	૧૫-૦૯-૧૯	૧૬-૦૯-૧૯	૧૭-૦૯-૧૯	૧૮-૦૯-૧૯
વરસાદ(મી.મી)	30	4	3	0	0
મહત્તમતાપમાન (સે.ગ્રે)	32	31	32	32	32
લઘુત્તમતાપમાન (સે.ગ્રે)	24	24	24	24	24
વાદળનીસ્થિતિ(ઓકટા)	8	8	8	8	7
મહત્તમભેજ(%) સવાર	87	86	88	86	87
લઘુત્તમભેજ(%) બપોરબાદ	66	70	70	64	64
પવનનીગતિ(કિમી/કલાક)	20	20	19	19	12
પવનનીદિશા(ડીગ્રી)	258	251	246	251	243

	<ul style="list-style-type: none"> <li>• ૧૪ થી ૧૮-સપ્ટેમ્બર -૨૦૧૯ હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. રાજકોટ જીલ્લામાં ૧૪ ના રોજ ભારે અને તા. ૧૫ ના રોજ હળવો વરસાદ થવાની શક્યતા છે, અમરેલી જીલ્લામાં તા. ૧૪ ના રોજ ભારેથી અતિ ભારે અને તા. ૧૫ થી ૧૭ ના રોજ હળવો થી મધ્યમ વરસાદ થવાની શક્યતા છે અને સુરેન્દ્રનગર જીલ્લામાં તા. ૧૪ થી ૧૬ ના રોજ હળવો થી મધ્યમ વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની મોટાભાગે રહેવાની અને પવનની ઝડપ ૧૦ થી ૨૦ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૯-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૪-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</li> <li>• વધારાના વરસાદી પાણીના નિકાલની વ્યવસ્થા કરવી અને તેનો ચેકડેમ,તળાવ, ખેતતલાવડીમાં સંગ્રહ કરવો</li> <li>• વરાપે જે તે પાકની ભલામણ મુજબ યુરીયા ખાતરનો પુરતી ડોઝ આપવો અને સમયસર નિંદામણ કરવું..</li> </ul>		
મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/ખાત	કૃષિ સલાહ (૧૦ લીટરપાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી )
મગફળી	સુયા બેસવા/ડોડવા બંધાવા	સફેદ ફૂગ  ચીતરી અને ગેરુ 	<ul style="list-style-type: none"> <li>• મગફળી પીળી પડેલ હોય તો હીરાકસી ૧૫૦ ગ્રામ અને લીંબુના ફૂલ ૧૫ ગ્રામ ૧ પમ્પમાં નાખીને છંટકાવ કરવો.</li> <li>• મગફળીમાં સફેદ ફૂગનો ઉપદ્રવ અટકાવવા ટ્રાયકોડર્મા ફૂગ ૨.૫ કી.ગ્રા./હે. મુજબ ગળતીયા દેશી ખાતર અથવા દિવેલાના ખોળ ૫૦૦ કી.ગ્રા./હે. મિશ્ર કરીને જમીનમાં આપવું.</li> </ul> <p>ચીતરી અને ગેરુના પાનના ટપકાના રોગના આગોતરા નિયંત્રણ માટે હેક્ઝાકોનાઝોલ ૧૫ મિલી અથવા ટેબ્યુકોનાઝોલ ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>
કપાસ	ચાપવા/ફૂલ	તડતડીયા અને થ્રીપ્સ  ગુલાબી ઈયળ	<ul style="list-style-type: none"> <li>• યુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૩ મિલી ૧૦ લીટર પાણીમાં નાખીને છંટકાવ કરવો.</li> <li>• કપાસમાં યુરીયા અને એમોનિયમ સલ્ફેટ અને મ્યુરેટ ઓફ પોટાશ દરેક ૩૦ કી.ગ્રા./હે. પુરતી ખાતર તરીકે આપવું.</li> <li>• કપાસમાં ગુલાબી ઈયળના નર કુદાને એકત્રીત કરવા માટે હેક્ટરે ૪૦ ની સંખ્યામાં ફેરોમેન ટ્રેપ ગોઠવવાથી ઈયળોનું શરૂઆતની અવસ્થાથી જ નિયંત્રણ કરી શકાય. કપાસમાં ગુલાબી ઈયળથી નુકશાન પામેલ ફૂલ અને ચાપવા તોડીને</li> </ul>

			<p>જમીનમાં દાટી દેવા. વધુ નુકશાન હોય તો પ્રોફેનોફોસ ૫૦% ઇ.સી. ૨૦ મી.લી. અથવા સાયપરમેથ્રીન ૧૦ મિલી બીવેરીયા બાસીઆના ફૂગનો પાઉડર ૪૦ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <ul style="list-style-type: none"> <li>જમીનમાં ભેજ જળવાય રહે તે માટે વરાપે આંતરખેડ, પાળા ચળાવવા અને નિંદામણ કરવું.</li> </ul>
તલ		<p>તલમાં માથા બાંધનારી</p> <p>ફાયટોફ્થોરા બ્લાઈટ અને ભુકીછારો</p>	<ul style="list-style-type: none"> <li>જમીનમાં ભેજ જળવાય રહે તે માટે વરાપે આંતરખેડ અને નિંદામણ કરવું.</li> <li>વાવેતરના ૩૦ દિવસ પછી વાદળછાયું વાતાવરણ ના હોય ત્યારે પૂર્તિ ખાતર તરીકે નાઈટ્રોજન (યુરીયા ૫૪ કી.ગ્રા./ હેક.) પ્રમાણે આપવું.</li> <li>તલમાં માથા બાંધનારી ઈયળના નિયંત્રણ માટે ક્વીનાલફોસ ૨૫ ટકા ૨૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</li> <li>સતત વરસાદી હવામાનને કારણે તલના પાકમાં સુકારા (ફાયટોફ્થોરા) અને ભુકીછારાના નિયંત્રણ માટે રીડોમિલ ૨૦ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</li> </ul>
બાજરી	દુધીયા દાણા	-	<ul style="list-style-type: none"> <li>જમીનમાં ભેજ જળવાય રહે તે માટે વરાપે આંતરખેડ અને નિંદામણ કરવું.</li> </ul>
કઠોળ પાકો	શીંગ	પીળો પંચરંગીયો	<ul style="list-style-type: none"> <li>સતત ભેજ વાળા હવામાનના કારણે વાયરસ વાળા પીળાછોડ ઉખાડીને દૂર કરવા. આ રોગ સફેદ માખીથી ફેલાતો હોય તેના નિયંત્રણ માટે ટ્રાયઓફોસ ૪૦ ઇસી દવા ૧૦ મિલી /૧૦ લીટર પાણીમાં મિશ્ર કરાવી.</li> <li>પાકેલી શીંગો ઉતારી લેવી</li> </ul>
ચોમાસું ડુંગળી	વૃદ્ધિ	પાનનો સુકારો	<ul style="list-style-type: none"> <li>ચોમાસુ ડુંગળીના પાકમાં પાનનાં સુકારાના નિયંત્રણ માટે મેન્કોઝેબ ૪૦ ગ્રામ અથવા હેક્ઝાકોનાઝોલ ૨૦ મિલી ૧ પમ્પમાં નાખીને ૧૦ દિવસના અંતરે ૩ છંટકાવ કરવા.</li> </ul>
ભીંડા	ફૂલ/ફળ	તડતડીયા,  શીંગ કોરી ખાનાર ઈયળ	<ul style="list-style-type: none"> <li>તડતડીયાના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૪ મિલી અથવા ટ્રાયઓફોસ ૪૦ ઇસી ૨૫ મિ.લી. દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</li> <li>તેના ફળ અને ડોકા કોરી ખાનાર ઈયળનો ઉપદ્રવ વધુ હોય તો સ્પીનોસાડ ૩ મિલી /૧૦ લીટર પાણીમાં ઓગાળી છંટકાવ કરવો.</li> </ul>

		પાન કથીરી	<ul style="list-style-type: none"> <li>જો પાનકથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છંટકાવ કરવો.</li> <li>ભીંડાની વીણી બાદ પ્રતિ એકરે ૫-૭ કી.ગ્રા. યુરિઆ ખાતર આપવું.</li> </ul>
શાકભાજી	ફેરરોપણી	મરચી,ટમેટી, રીંગણ, અને કોબીઝના ધરૂં તૈયાર હોય વરસાદ થયે ફેરરોપણી કરવી.	
તુરીયા, દૂધી, ગલકા, કારેલા, કાકડી	ફૂલ/ફળ	પાન કોરીયું, થ્રીપ્સ, સફેદ માખી,  ફળ માખી  તળછારો  કથીરી	<ul style="list-style-type: none"> <li>વેલાવાળા શાકભાજીમાં યુસીયા જીવાતના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૪ મિલી અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</li> <li>વેલાવાળા શાકભાજીમાં ફળમાખીના નિયંત્રણ માટે વાદળછાયું વાતાવરણ ના હોય ત્યારે રોગર ૧૦ મિલી + ૧૦૦ ગ્રામ ગોળ અથવા ખાંડ ૧૦ લીટર પાણીમાં હેક્ટરે ૫૦ લીટર પાણી જાય એ રીતે છંટકાવ કરવો.</li> <li>જો પાનકથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છંટકાવ કરવો.</li> <li>મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.</li> <li>વેલાવાળા શાકભાજી પાકોમાં હળવા અને ટુંકા અંતરે પિયત આપી યોગ્ય ભેજ જાળવવો જેથી પરાગનયન પ્રક્રિયા સારી રીતે થવાથી ઉત્પાદન સારું મળે.</li> </ul>
રીંગણી અને ટમેટી	ફૂલ/ફળ	સફેદ માખી  ડુંખ અને→ ફળ કોરી ખાનાર ઈયળ  પાન કથીરી	<ul style="list-style-type: none"> <li>લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો દ્રાયઝોફોસ ૪૦ ઈ.સી. ૨૦ મી.લી. અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</li> <li>ફળ અને ડોકાની ઈયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને ક્લોરાન્ટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઈસી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</li> <li>જો પાનકથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છંટકાવ કરવો.</li> </ul>
મરચી	ફૂલ	પાનની કુક્ડ- થ્રીપ્સ→	<ul style="list-style-type: none"> <li>દ્રાયઝોફોસ ૪૦ ઈ.સી. ૨૦ મી.લી. અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અથવા પ્રોફેનોફોસ ૨૦ મી.લી.અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</li> </ul>





ગ્રામીણ કૃષિ મૌસમ સેવા  
હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન  
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ  
તા.૧૪-૦૯-૨૦૧૯ થી તા.૧૮-૦૯-૨૦૧૯



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કૃ.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિહવામાન બુલેટીન નં. ૪૮/૨૦૧૯

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી)	52.5
મહત્તમ તાપમાન (સે.ગ્રે)	28.0-32.5
લઘુત્તમ તાપમાન (સે.ગ્રે)	22.7-24.1
ભેજ(%) સવાર	85-98
ભેજ(%) બપોરબાદ	79-100
પવનની ગતિ (કિમી/કલાક)	2.3-6.0

હવામાન આગાહી: પત્રક-ક

ઉત્તરસૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.૧૪-૦૯-૨૦૧૯ થી ૧૮-૦૯-૨૦૧૯

કૃષિ સલાહ:

સામાન્ય	ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના					
	<b>મોરબી:</b> જીલ્લામાં આગામી તા.૧૪-૦૯-૨૦૧૯ થી તા.૧૮-૦૯-૨૦૧૯ દરમિયાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૧૪ થી ૧૫ ના રોજ મધ્યમ થી ભારે અને તા. ૧૭ ના રોજ હળવો વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૮ થી ૧૭ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૧-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૪-૨૪ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.					
	તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
	પેરામીટર	૧૪-૦૯-૧૯	૧૫-૦૯-૧૯	૧૬-૦૯-૧૯	૧૭-૦૯-૧૯	૧૮-૦૯-૧૯
	વરસાદ(મી.મી)	40	23	0	8	0
	મહત્તમતાપમાન (સે.ગ્રે)	32	31	32	32	32
	લઘુત્તમતાપમાન (સે.ગ્રે)	24	24	24	24	24

વાદળનીસ્થિતિ(ઓકટા)	8	8	8	8	7
મહત્તમભેજ(%) સવાર	96	96	96	92	94
લઘુત્તમભેજ (%) બપોરબાદ	92	93	81	74	74
પવનનીગતિ(કિમી/કલાક)	17	11	9	12	8
પવનનીદિશા(ડીગ્રી)	268	244	244	253	269

**જામનગર:** જીલ્લામાં આગામી તા.૧૪-૦૯-૨૦૧૯ થી તા.૧૮-૦૯-૨૦૧૯ દરમિયાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૧૪ થી ૧૫ ના રોજ હળવો થી મધ્યમ વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની રહેવાની અને પવનની ઝડપ ૧૫ થી ૨૭ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૦-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૬-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.




તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૪-૦૯-૧૯	૧૫-૦૯-૧૯	૧૬-૦૯-૧૯	૧૭-૦૯-૧૯	૧૮-૦૯-૧૯
વરસાદ(મી.મી)	17	6	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	31	30	32	32	32
લઘુત્તમતાપમાન (સે.ગ્રે)	27	26	26	27	26
વાદળનીસ્થિતિ(ઓકટા)	8	8	8	8	7
મહત્તમભેજ(%) સવાર	85	87	87	85	86
લઘુત્તમભેજ (%) બપોરબાદ	74	80	76	73	73
પવનનીગતિ(કિમી/કલાક)	27	20	23	25	15
પવનનીદિશા(ડીગ્રી)	262	256	254	252	247

**દેવભૂમિ દ્વારકા:** જીલ્લામાં આગામી તા.૧૪-૦૯-૨૦૧૯ થી તા.૧૮-૦૯-૨૦૧૯ દરમિયાન હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહેશે. તા. ૧૪ ના રોજ હળવો વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ ૧૩ થી ૨૭ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૨૯-૩૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૬-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.


તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૪-૦૯-૧૯	૧૫-૦૯-૧૯	૧૬-૦૯-૧૯	૧૭-૦૯-૧૯	૧૮-૦૯-૧૯
વરસાદ(મી.મી)	10	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	30	29	30	29	31
લઘુત્તમતાપમાન (સે.ગ્રે)	27	26	26	26	27
વાદળનીસ્થિતિ(ઓકટા)	8	8	8	8	7
મહત્તમભેજ(%) સવાર	84	87	86	84	85
લઘુત્તમભેજ (%) બપોરબાદ	72	79	71	71	70
પવનનીગતિ(કિમી/કલાક)	27	18	22	24	13
પવનનીદિશા(ડીગ્રી)	262	256	255	254	256







- ૧૪ થી ૧૮-સપ્ટેમ્બર -૨૦૧૯ હવામાન ગરમ, ભેજવાળું અને વાદળછાયું રહશે. મોરબી જીલ્લામાં તા. ૧૪ થી ૧૫ ના રોજ મધ્યમ થી ભારે અને તા. ૧૭ ના રોજ હળવો વરસાદ થવાની શક્યતા છે, જામનગર જીલ્લામાં તા. ૧૪ થી ૧૫ ના રોજ હળવો થી મધ્યમ વરસાદ થવાની શક્યતા છે અને દેવભૂમિ દ્વારકા તા. ૧૪ ના રોજ હળવો વરસાદ થવાની શક્યતા છે. પવનની દિશા પશ્ચિમ અને નૈઋત્યની મોટાભાગે રહેવાની અને પવનની ઝડપ ૮ થી ૨૭ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૯-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૪-૨૭ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.
- વધારાના વરસાદી પાણીના નિકાલની વ્યવસ્થા કરવી અને તેનો ચેકડેમ, તળાવ, ખેતતલાવડીમાં સંગ્રહ કરવો
- વરાપે જે તે પાકની ભલામણ મુજબ યુરીયા ખાતરનો પુરતી ડોઝ આપવો અને સમયસર આંતરખેડ અને નિંદામણ કરવું..

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી )
મગફળી	ફૂલ/સુયા	સફેદ ફૂગ  ચીતરી અને ગેરુ 	<ul style="list-style-type: none"> <li>મગફળી પીળી પડેલ હોય તો હીરાકસી ૧૫૦ ગ્રામ અને લીંબુના ફૂલ ૧૫ ગ્રામ ૧ પમ્પમાં નાખીને છંટકાવ કરવો.</li> <li>મગફળીમાં સફેદ ફૂગનો ઉપદ્રવ અટકાવવા ટ્રાયકોડર્મા ફૂગ ૨.૫ કી.ગ્રા./હે. મુજબ ગળતીયા દેશી ખાતર અથવા દિવેલાના ખોળ ૫૦૦ કી.ગ્રા./હે. મિશ્ર કરીને જમીનમાં આપવું.</li> <li>ચીતરી અને ગેરુના પાનના ટપકાના રોગના આગોતરા નિયંત્રણ માટે હેક્ઝાકોનાઝોલ ૧૫ મિલી અથવા ટેબ્યુકોનાઝોલ ૧૦ મિલી ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો.</li> </ul>
કપાસ	ચાપવા	તડતડીયા / થ્રીપ્સ	<ul style="list-style-type: none"> <li>ચુસીયા પ્રકારની જીવાતના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૩ મિલી ૧૦ લીટર પાણીમાં નાખીને છંટકાવ કરવો.</li> <li>સતત ભેજ વાળા હવામાનના કારણે પાનનાં ટપકાના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ / ૧૦ લિટર પાણીમાં નાંખીને ૧૫ દિવસ ના અંતરે છંટકાવ કરવો.</li> <li>૨૫-૩૦ દિવસના કપાસમાં ચુરીયા, એમોનિયમ સલ્ફેટ અને મ્યુરેટ ઓફ પોટાશ દરેક ૨૦ કી.ગ્રા./હે. પુરતી ખાતર તરીકે આપવું.</li> </ul>
તલ	ફૂલ	માથા બાંધનાર ઈયળ  ભુકીછારો અને ફાયટોપ્થોરા બ્લાઈટ	<ul style="list-style-type: none"> <li>કવીનાલફોસ ૨૦ મિલી અથવા કાર્ટેપ હાયડ્રોકલોરાઈડ ૧૫ ગ્રામ ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો.</li> <li>તલમાં ભુકીછારો અને ફાયટોપ્થોરા બ્લાઈટ રોગનો ઉપદ્રવ ના થાય તે માટે રીડોમિલ દવાનો ૨૦ મિલી/૧૦ લિટર પાણીમાં મિશ્ર કરી ૧૦ દિવસે બે છંટકાવ કરવા.</li> <li>તલમાં ચુરીયા ૫૪ કી.ગ્રા./હે. પુરતી ખાતર તરીકે આપવું.</li> <li>વરાપે આંતરખેડ અને નિંદામણ કરવું.</li> </ul>
બાજરી	ફૂટ	-	<ul style="list-style-type: none"> <li>વરાપે આંતરખેડ અને નિંદામણ કરવું.</li> </ul>

કઠોળ પાકો	ફૂલ	પીળો પંચરંગીયો	<ul style="list-style-type: none"> <li>• વરાપે આંતરખેડ અને નિંદામણ કરવું.</li> <li>• સતત ભેજ વાળા હવામાનના કારણે વાયરસ વાળા પીળાછોડ ઉખાડીને દૂર કરવા. આ રોગ સફેદ માખીથી ફેલાતો હોય તેના નિયંત્રણ માટે ટ્રાયએફોસ ૪૦ ઇસી દવા ૧૦ મિલી /૧૦ લીટર પાણીમાં મિશ્ર કરાવી.</li> </ul>
જુવાર/ મકાઈ	વૃદ્ધિ	ડોકા મરડો  મકાઈ:ફોલ આર્મી વર્મ	<p>૨૫-૩૦ દિવસની જુવારમાં યુરીયા ૮૬ કી.ગ્રા./હે આપવું. તેમજ ડોકા મરડો ઉપદ્રવનું સતત નિરીક્ષણ કરતા રહેવું. જરૂર જણાયે વધુ ઉપદ્રવ હોય તો ક્વીનાલફોસ ૨૦ મીલી અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>મકાઈના પાકમાં ફોલ આર્મી વર્મનો ઉપદ્રવ હોય એમામેક્ટીન બેન્ઝોએટ ૭ ગ્રામ પ્રતિ ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</p>
લીંડા	ફૂલ/ફળ	તડતડીયા,  શીગ કોરી ખાનાર ઈયળ  પાન કથીરી	<ul style="list-style-type: none"> <li>• તડતડીયાના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૪ મિલી અથવા ટ્રાયએફોસ ૪૦ ઇસી ૨૫ મિ.લી. દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</li> <li>• તેના ફળ અને ડોકા કોરી ખાનાર ઈયળનો ઉપદ્રવ વધુ હોય તો સ્પીનોસાડ ૩ મિલી /૧૦ લીટર પાણીમા ઓગાળી છંટકાવ કરવો.</li> <li>• જો પાન કથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છંટકાવ કરવો.</li> </ul> <p>લીંડાની વીણી બાદ પ્રતિ એકરે ૫-૭ કી.ગ્રા. યુરિઆ ખાતર આપવું.</p>
શાકભાજી	ધરું ઉછેર/રોપણી વાવેતર	મરચી,ટમેટી, રીંગણ, અને કોબીઝના ધરું તૈયાર હોય વરસાદ થયે ફેરોપણી કરવી.	

<p>તુરીયા, દૂધી, ગલકા, કારેલા, કાકડી</p>	<p>ફૂલ/ફળ</p>	<p>પાન કોરીયું , થ્રીપ્સ, સફેદ માખી,  ફળ માખી  તળછારો  કથીરી</p>	<p>➤ વેલાવાળા શાકભાજીમાં સફેદમાખીના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૪ મિલી અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</p> <p>➤ વેલાવાળા શાકભાજીમાં ફળમાખીના નિયંત્રણ માટે વાદળછાયું વાતાવરણ ના હોય ત્યારે રોગર ૧૦ મિલી + ૧૦૦ ગ્રામ ગોળ અથવા ખાંડ ૧૦ લીટર પાણીમાં હેક્ટરે ૫૦ લીટર પાણી જાય એ રીતે છંટકાવ કરવો.</p> <p>➤ જો પાન કથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છાન્કાવ કરવો.</p> <p>• મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.</p> <p>વેલાવાળા શાકભાજી પાકોમાં હળવા અને ટુંકા અંતરે પિયત આપી યોગ્ય ભેજ જાળવવો જેથી પરાગનયન પ્રક્રિયા સારી રીતે થવાથી ઉત્પાદન સારું મળે.</p>
<p>રીંગણી અને ટમેટી</p>	<p>ફૂલ/ફળ</p>	<p>સફેદ માખી  ડુંખ અને→ ફળ કોરી ખાનાર ઈયળ  કથીરી →</p>	<p>➤ લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો ટ્રાયઝોફોસ ૪૦ ઈ.સી. ૨૦ મી.લી. અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>➤ ફળ અને ડોકાની ઈયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને ક્લોરાનટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઈસી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</p> <p>➤ રીંગણી અને ટમેટીના પાકમાં પાન કથીરીના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</p>
<p>મરચી</p>	<p>ફૂલ/ફળ</p>	<p>પાનની કુક્કડ- થ્રીપ્સ→ </p>	<p>ટ્રાયઝોફોસ ૪૦ ઈ.સી. ૨૦ મી.લી. અથવા એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અથવા પ્રોફેનોફોસ ૨૦ મી.લી.અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>વાયરસના કારણે પીળા થઈ કુક્કાય ગયેલ છોડ ઉપાડી તેનો</p>

		<p>પાનની કુક્કડ-</p>  <p>કથીરી</p>	<p>જમીનમાં દાટી નાશ કરવો. ત્યાં સફેદ માખીથી ફેલાતો હોવાથી તેના નિયંત્રણ માટે હોસ્તાથીયોન દવા ૨૦ મી.લી./૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>જો પાન કથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છાન્કાવ કરવો.</p>
લીંબુ	ફળનો વિકાસ	<p>પાન કથીરી</p>  <p>સાયલા અને થ્રીપ્સ</p>	<p>ઊંચા તાપમાનને કારણે લીંબુના બગીચામાં પાન કથીરીનો ઉપદ્રવ વધુ હોય તો તેના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>તેના નિયંત્રણ માટે ઇમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૫ મીલી અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. વધુ ઉપદ્રવ હોય તો ટ્રાયએફોસ ૪૦ ટકા ૨૦/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>  <p>બદામી ટપકાના નિયંત્રણ માટે કોપર ઓક્સીક્લોરાઈડ ૪૦ ગ્રામ અને ૧.૫ ગ્રામ સ્ટ્રેપ્ટોસાયક્લીનના બે છંટકાવ ૧૫ દિવસના ગળે કરવા.</p> <p>પાનકોરીયુના નિયંત્રણ માટે ડીડીવીપી ૧૦ મી.લી. ૧૦ લિટર પાણીમાં મિશ્ર કરીને છાંટવી.</p>
પશુ/ મરઘા			<p>દુધાળા પશુના આઉને ઝીંક ઓક્સાઈડ અથવા બોરીક પાવડર થી બરાબર સાફ કરવું. તેમજ ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાલ ૨ મી.લી. એક લીટર પાણીમાં નાખીને છાંટવી અને ઉછરતા પશુઓમાં કરમીયાની ગોળી આપવી. પશુને ૬ થી ૮ કી.ગ્રા. સુકો અને ૧૫ થી ૨૦ કી.ગ્રા. લીલો ચારો આપવો. દુધાળા પ્રાણીઓને નિયમિત ૧ કિલ્લો ફીડ + ૫૦ ગ્રામ ખનીજ તત્વોનું મિશ્રણ પ્રતિ ૨ લીટર દુધની ઊપજના પ્રમાણમાં આપવું. તેમજ લીલું ઘાસ + સૂકું ઘાસ + ખનીજ તત્વો ખોરાકમાં આપવા.</p> 

ચોમાસા દરમ્યાન પશુઓના રહેઠાણની જગ્યાની વ્યવસ્થિત સફાઈ રાખવી અને માખીઓના ઉપદ્રવથી બચવા ફિનાઈલ છાંટવું.  
ખુલ્લામાં રાખેલ ઘાસસારો વરસાદથી બગડે નહિ તે માટે જરૂરી પગલા ભરવા.

જુક્યુ/ટેકનિકલ ઓફિસર.(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૧૯  
તરઘડિયા(રાજકોટ), તારીખ: ૧૩/૦૯/૨૦૧૯.

પ્રતિ,  
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,  
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનંતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર  
સુકી ખેતીસંશોધન કેન્દ્ર  
જુનાગઢ કૃષિ નિવર્સિટી  
તરઘડીયા



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :942756433**



**Agro met Advisory Service bulletin No.48-for the Rajkot district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 14to 18-September- 2019)

**Significant past weather for the preceding week of Rajkot**

Rainfall (mm)	52.5
Maximum temperature (°C)	28.0-32.5
Minimum temperature (°C)	22.7-24.1
Relative Humidity (%) I	85-98
Relative Humidity (%) II	79-100
Wind speed (Kmph)	2.3-6.0


**Weather forecast until 08-30 hrs. of 18/09/2019**

**(1) RAJKOT DISTRICT:**

	14/09/19	15/09/19	16/09/19	17/09/19	18/09/19
Rainfall (mm)	43	6	0	0	0
Max.Temp.(°C)	32	31	32	32	32
Min.Temp.(°C)	24	24	24	24	24
Total CC (octa)	8	8	8	8	7
RH-I (%)	89	90	90	88	91
RH-II (%)	68	75	75	68	67
Wind Speed (kmph)	18	18	18	19	11
Wind Direction(deg.)	262	252	248	251	237





**Agro meteorological Advisories:**


<b>General Advice</b>	<p><b>Rain fall: Warm, humid and generally cloudy to cloudy weather &amp; Heavy rainfall during 14-September &amp; Light rainfall during 15-September forecasted.</b></p> <ul style="list-style-type: none"> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 31-32°C and 24-24°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 88-91% &amp; 67-75, respectively.</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W, SW and 11-19 km/h, respectively.</li> <li>• <b>Apply supplementary dose of urea fertilizer as per recommendation of respective</b></li> </ul>
-----------------------	---

	<p><b>crop for optimum growth.</b></p> <ul style="list-style-type: none"> <li><b>Carry out weeding and plant protection measures as per need.</b></li> </ul>		
Crops	Stage	Insect/Pest/ Variety	Advisory
<b>Kharif groundnut</b>	Pegging/ pod formation	<p><b>Sclerotium rolfsii</b></p> <p><b>Tikka &amp; rust</b></p> 	<ul style="list-style-type: none"> <li><b>Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kh/ha as preventive measures against Sclerotiumrolfsii stem rot.</b></li> <li><b>Spray Tebuconazole 10 ml/10 lit of to prevent leaf spot (Tikka &amp; Rust) infestation n groundnut.</b></li> <li><b>Carry out weeding at vapsa condition.</b></li> <li><b>Spray ferrous sulfate 100 gram &amp; citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop.</b></li> </ul>
<b>Cotton</b>	Square& flower initiation	<p><b>Jassid &amp;Thrips</b></p> <p><b>Pink ball worm</b></p>	<ul style="list-style-type: none"> <li><b>Monitor the crops for infestation of sucking pest, if require spray imidacloprid 30.5 % 3 ml / 10 lit of water ml/10 lit of water.</b></li> <li><b>Due to constant moist weather apply supplementary dose of nitrogen through Ammonium sulphate (30 kg/ha) and also apply potash through murate of potash (30 kg/ha) to cotton crop.</b></li> <li><b>Also spray hexaconazole 5 % EC @ 15 ml/10 litre of water for control alternaria&amp;cercospora leaf spot in cotton.</b></li> <li><b>Carry out interculturing &amp; weeding at vapsa condition.</b></li> <li><b>Also earthingup/ridging should be done in time.</b></li> <li><b>For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found <b>pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water.</b></b></li> </ul>



<b>Sesame</b>	Growing/flowering	<b>Leaf webber</b>  <b>Phytophthora blight &amp; Powdery mildew</b>	<ul style="list-style-type: none"> <li>• For the control of leaf webber spray quinalphos 25 % Ec @ 20 ml/ 10 litre of water.</li> <li>• Apply supplementary dose of nitrogen(Urea 54 kg/ha) at 30 DAS under clear sky.</li> <li>• Due to warm &amp; humid weather, for prevention of phytophthora blight &amp; powdery mildew in sesame spray Ridomil 20 gm in 10 liters of water.</li> <li>• Carry out weeding at vapsa condition.</li> </ul>
<b>Pearl millet</b>	Milking	-	<ul style="list-style-type: none"> <li>• Carry out weeding at vapsa condition.</li> </ul>
<b>Pulses</b>	Pod development & picking	<b>Yellow mosaic viral disease</b>	<ul style="list-style-type: none"> <li>• Due to continuous rainfall infestation of yellow mosaic virus observed in pulse crop, for control remove the infected plants and make spray of Triazophos 40 % EC @ 10 ml/10 litre of water.</li> <li>• Carry out interculturing operation for soil moisture conservation &amp; removal of weed at vapsa condition.</li> <li>• Pick mature pod in time.</li> </ul>
<b>Kharif Onion</b>	Growing	<b>Alternaria leaf spot/purple blotch</b>	<ul style="list-style-type: none"> <li>• For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb 40 gm and Hexaconazole at 10 days interval.</li> <li>• Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky.</li> </ul>
<b>Vegetables</b>	Transplanting/growing	Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
<b>Lady finger</b>	Flowering/fruiting	<b>Jassids &amp; white fly</b>  <b>Shoot and fruit borer</b>  <b>Mites</b>	<p>Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassid &amp; white fly control.</p> <p>Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray). If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.</p> <p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear.</p>

			After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.
<b>Ridge gourd, Bottle gourd, Sponge guard, Bitter guard &amp; cucurbitaceous crops</b>	Flowering & fruiting	<b>Fruit fly</b>  <b>Mites</b>	For constant monitoring of cucurbitaceous crops against attack of fruit fly, use of 'Methyl Eugenol trap is recommended. After detecting the symptoms, spraying of Rogor @ 20 ml + 100 g sugar or jaggary/10 litre water at 50 litres/ha. is advised, when sky is clear. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites under clear sky.
<b>Cluster bean</b>	Growing/fruiting	<b>Thrips/whitefly</b>	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
<b>Brinjal &amp; Tomato</b>	Flowering/fruiting	<b>Sucking pest</b> → <b>&amp; Fruit borer</b>   <b>Mites</b>	Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassids & white fly control. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer. If mite population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear.
<b>Chili</b>	Flowering	<b>Thrips</b> →  <b>Leaf curling Virus</b> → 	Under present weather condition, for the control of thrips infestation make alternative spray of profenofos 50 EC and Ethion 50 % EC @ 20 ml /10 litre of water at 10 days interval.  In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear.
<b>Lemon</b>	Fruit development	<b>Psylla and thrips</b> 	To prevent damage of psylla & thrips spray <b>imidacloprid 30.5 % 5 ml</b> / 10 lit of water ml/10 lit of water.

		<p><b>Citrus canker</b></p> <p><b>Leaf minor</b></p>	<p>Spray copper oxychloride 40 gm or streptomycin 1.5 gm in 10 litre of water</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
<p><b>Livestock &amp; poultry</b></p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming testing should be done. Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :94277 28523**



**Agro met Advisory Service bulletin No.-48 for the Morbi district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 14 to 18-September- 2019)


**Weather forecast until 08-30 hrof 18/09/2019**


**(2) MORBI DISTRICT:**





	14/09/19	15/09/19	16/09/19	17/09/19	18/09/19
Rainfall (mm)	40	23	0	8	0
Max. Temp.(°C)	32	31	32	32	32
Min. Temp.(°C)	24	24	24	24	24
Total CC (octa)	8	8	8	8	7
RH-I (%)	96	96	96	92	94
RH-II (%)	92	93	81	74	74
Wind Speed (kmph)	17	11	9	12	8
Wind Direction(deg.)	268	244	244	253	269

**Agro meteorological Advisories:-**

<b>General Advice</b>	<ul style="list-style-type: none"> <li>• <b>Rain fall:</b> Warm, humid and generally cloudy to cloudy weather &amp; Moderate to heavy rainfall during 14 to 15-September &amp; light rainfall during 17-September forecasted.</li> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 31-32°C and 24-24°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 92-96% &amp; 74-93, respectively.</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W, SW and 8-17 km/h, respectively.</li> <li>• <b>Apply supplementary dose of urea fertilizer as per recommendation of respective crop for optimum growth.</b></li> <li>• <b>Drain out excess water from field for better growth of plant and store in reservoirs.</b></li> <li>• <b>Carry out weeding and plant protection measures as per need.</b></li> </ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/Variety</b>	<b>Advisory</b>
<b>Kharif groundnut</b>	Flowering /pegging	Sclerotium rolfsii	• <b>Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kh/ha as</b>

		<p><b>Tikka &amp; rust</b></p> 	<p><b>preventive measures against Sclerotiumrolfsii stem rot.</b></p> <ul style="list-style-type: none"> <li>• <b>Spray Tebuconazole 10 ml/10 lit of to prevent leaf spot (Tikka &amp; Rust) infestation n groundnut.</b></li> <li>• <b>Carry out weeding at vapsa condition.</b></li> <li>• <b>Spray ferrous sulfate 100 gram &amp; citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop.</b></li> </ul>
<b>Cotton</b>	Squaring	<b>Jassid &amp;Thrips</b>	<p>Monitor the crops for infestation of sucking pest, if require spray <b>imidacloprid 30.5 % 3 ml / 10 lit of water ml/10 lit of water.</b></p> <p>Due to continuous humid weather spray Mancozeb 27 gram/10 lit of water at 15 days interval to prevent leaf spot infestation.</p> <p>Apply 20 kg/ha urea, Ammonium sulphate and Murate of Potash each to cotton crop.</p>
<b>Sesame</b>	Flowering	<p><b>Leaf webber</b></p> <p><b>Phytophthora blight&amp; Powdery mildew</b></p>	<ul style="list-style-type: none"> <li>• <b>For the control of leaf webber spray quinalphos 25 % Ec @ 20 ml/ 10 litre of water.</b></li> <li>• <b>Apply supplementary dose of nitrogen(Urea 54 kg/ha) at 30 DASunder clear sky.</b></li> <li>• <b>Due to warm &amp; humid weather, for prevention of phytophthora blight &amp; powdery mildew in seamum spray Ridomil 20 gm in 10 liters of water.</b></li> <li>• <b>Carry out weeding at vapsa condition.</b></li> </ul>
<b>Pearl millet</b>	Tillering	-	<ul style="list-style-type: none"> <li>• <b>Carry out interculturing&amp;weeding at vapsa condition.</b></li> <li>• <b>Apply supplementary dose of nitrogen(Urea 87 kg/ha) at 30 DASunder clear sky.</b></li> </ul>
<b>Pulses</b>	Flowering		<ul style="list-style-type: none"> <li>• <b>Carry out interculturing&amp;weeding at vapsa condition.</b></li> </ul>
<b>Castor</b>	Germination	GAUCH-1,GC-3, GCH-4, GCH-6, GCH-7	
<b>Fodder Sorghum &amp; Maize</b>	Growing	Sorghum-Gundari, GFS-4 & 5 Maize-African tall, Ganga safed-2 & 5	<p><b>Monitor the crop for shoot fly infestation and if require spraycypermethrin 25 EC 5 ml in 10 liters of water. Infestation of fall army worm observed in maize, so for control make spray of Emamectin Benzoate 7 gram/10 litre of water.</b></p>

<b>Kharif Onion</b>	Growing	Alternaria leaf spot/purple blotch	<p><b>For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb 40 gm and Hexaconazole at 10 days interval.</b></p> <ul style="list-style-type: none"> <li>• <b>Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky.</b></li> </ul>
<b>Vegetables</b>	Transplanting	Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
<b>Lady finger</b>	Flowering/ fruiting	<p><b>Jassids &amp; white fly</b></p> <p><b>Shoot and fruit borer</b></p> <p><b>Mites</b></p>	<p>Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassids control.</p> <p>Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray). If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.</p> <p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear.</p> <p>After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.</p>
<b>Ridge gourd, Bottle gourd, Sponge gourd, Bitter guard &amp; cucurbitaceous crops</b>	Flowering & fruiting	<p><b>Fruit fly</b></p> <p><b>Mites</b></p>	<p>For constant monitoring of cucurbitaceous crops against attack of fruit fly, use of 'Methyl Eugenol trap' is recommended. After detecting the symptoms, spraying of Rogor @ 20 ml + 100 g sugar or jaggery/10 litre water at 50 litres/ha is advised, when sky is clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites under clear sky.</p>
<b>Cluster bean</b>	Growing/ fruiting	<b>Thrips/whitefly</b>	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 ml or neem oil 40 ml and <b>acetamiprid</b> 20 % SP @ 5.0 gm / 10 lit of water at 15 days interval
<b>Brinjal &amp; Tomato</b>	Flowering/ fruiting	<p><b>Sucking pest</b> →</p> <p><b>&amp; Fruit borer</b></p>  <p><b>Mites</b></p>	<p>Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassids &amp; white fly control.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p> <p>If mite population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when</p>

			sky remains clear.
<b>Chili</b>	Flowering	<b>Thrips</b> →  <b>Leaf curling Virus</b> → 	<p>Under present weather condition, for the control of thrips infestation make alternative spray of profenofos 50 EC and Ethion 50 % EC @ 20 ml /10 litre of water at 10 days interval.</p> <p>In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advisedfor control the vector, when sky remains clear.</p>
<b>Lemon</b>	Fruit development	<b>Psylla and thrips</b>  <b>Citrus canker</b> <b>Leaf minor</b>	<p>To prevent damage of psylla&amp;thrips spray <b>imidacloprid 30.5 % 5 ml</b> / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.</p>
<b>Livestock &amp; poultry</b> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Dewormingtesting should be done. Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :94277 28523**



**Agro met Advisory Service bulletin No.-48 for the Jamnagar district**

(Issued jointly by DFERS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 14 to 18-September- 2019)

**Weather forecast until 08-30 h of 18/09/2019**


**(3) JAMNAGAR DISTRICT:**


	14/09/19	15/09/19	16/09/19	17/09/19	18/09/19
Rainfall (mm)	17	6	0	0	0
Max. Temp.(°C)	31	30	32	32	32
Min. Temp.(°C)	27	26	26	27	26
Total CC (octa)	8	8	8	8	7
RH-I (%)	85	87	87	85	86
RH-II (%)	74	80	76	73	73
Wind Speed (kmph)	27	20	23	25	15
Wind Direction(deg.)	262	256	254	252	247





**Agro meteorological Advisories:-**

<b>General Advice</b>	<ul style="list-style-type: none"> <li>• <b>Rain fall:</b> Warm, humid and generally cloudy to cloudy weather &amp; Light to moderate rainfall during 14 to 15-September forecasted.</li> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 30-32°C and 26-27°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 85-87% &amp; 73-80, respectively.</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W, SW and 15-27 km/h, respectively.</li> <li>• <b>Apply supplementary dose of urea fertilizer as per recommendation of respective crop for optimum growth.</b></li> <li>• <b>Drain out excess water from field for better growth of plant and store in reservoirs.</b></li> <li>• <b>Carry out weeding and plant protection measures as per need.</b></li> </ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/Variety</b>	<b>Advisory</b>
<b>Kharif groundnut</b>	Flowering /pegging	Sclerotium rolfsii	• <b>Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kg/ha as</b>



		<p><b>Tikka &amp; rust</b></p> 	<p><b>preventive measures against Sclerotiumrolfsii stem rot.</b></p> <ul style="list-style-type: none"> <li>• <b>Spray Tebuconazole 10 ml/10 lit of to prevent leaf spot (Tikka &amp; Rust) infestation n groundnut.</b></li> <li>• <b>Carry out weeding at vapsa condition.</b></li> <li>• <b>Spray ferrous sulfate 100 gram &amp; citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop.</b></li> </ul>
<b>Cotton</b>	Squaring	<b>Jassid &amp;Thrips</b>	<p>Monitor the crops for infestation of sucking pest, if require spray <b>imidacloprid 30.5 % 3 ml / 10 lit of water ml/10 lit of water.</b></p> <p>Due to continuous humid weather spray Mancozeb 27 gram/10 lit of water at 15 days interval to prevent leaf spot infestation.</p> <p>Apply 20 kg/ha urea, Ammonium sulphate and Murate of Potash each to cotton crop.</p>
<b>Sesame</b>	Flowering	<p><b>Leaf webber</b></p> <p><b>Phytophthora blight&amp; Powdery mildew</b></p>	<ul style="list-style-type: none"> <li>• <b>For the control of leaf webber spray quinalphos 25 % Ec @ 20 ml/ 10 litre of water.</b></li> <li>• <b>Apply supplementary dose of nitrogen(Urea 54 kg/ha) at 30 DASunder clear sky.</b></li> <li>• <b>Due to warm &amp; humid weather, for prevention of phytophthora blight &amp; powdery mildew in seamum spray Ridomil 20 gm in 10 liters of water.</b></li> <li>• <b>Carry out weeding at vapsa condition.</b></li> </ul>
<b>Pearl millet</b>	Tillering	-	<ul style="list-style-type: none"> <li>• <b>Carry out interculturing&amp;weeding at vapsa condition.</b></li> <li>• <b>Apply supplementary dose of nitrogen(Urea 87 kg/ha) at 30 DASunder clear sky.</b></li> </ul>
<b>Pulses</b>	Flowering		<ul style="list-style-type: none"> <li>• <b>Carry out interculturing&amp;weeding at vapsa condition.</b></li> </ul>
<b>Castor</b>	Germination	GAUCH-1,GC-3, GCH-4, GCH-6, GCH-7	
<b>Fodder Sorghum &amp; Maize</b>	Growing	Sorghum-Gundari, GFS-4 & 5 Maize-African tall, Ganga safed-2 & 5	<p><b>Monitor the crop for shoot fly infestation and if require spraycypermethrin 25 EC 5 ml in 10 liters of water. Infestation of fall army worm observed in maize, so for control make spray of Emamectin Benzoate 7 gram/10 litre of water.</b></p>

<b>Kharif Onion</b>	Growing	Alternaria leaf spot/purple blotch	<p><b>For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb 40 gm and Hexaconazole at 10 days interval.</b></p> <ul style="list-style-type: none"> <li>• <b>Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky.</b></li> </ul>
<b>Vegetables</b>	Transplanting	Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
<b>Lady finger</b>	Flowering/ fruiting	<p><b>Jassids &amp; white fly</b></p> <p><b>Shoot and fruit borer</b></p> <p><b>Mites</b></p>	<p>Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassids control.</p> <p>Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray). If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.</p> <p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear.</p> <p>After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.</p>
<b>Ridge gourd, Bottle gourd, Sponge gourd, Bitter guard &amp; cucurbitaceous crops</b>	Flowering & fruiting	<p><b>Fruit fly</b></p> <p><b>Mites</b></p>	<p>For constant monitoring of cucurbitaceous crops against attack of fruit fly, use of 'Methyl Eugenol trap' is recommended. After detecting the symptoms, spraying of Rogor @ 20 ml + 100 g sugar or jaggery/10 litre water at 50 litres/ha is advised, when sky is clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites under clear sky.</p>
<b>Cluster bean</b>	Growing/ fruiting	<b>Thrips/whitefly</b>	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 ml or neem oil 40 ml and <b>acetamiprid</b> 20 % SP @ 5.0 gm / 10 lit of water at 15 days interval
<b>Brinjal &amp; Tomato</b>	Flowering/ fruiting	<p><b>Sucking pest</b> →</p> <p><b>&amp; Fruit borer</b></p>  <p><b>Mites</b></p>	<p>Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassids &amp; white fly control.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p> <p>If mite population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when</p>

			sky remains clear.
<b>Chili</b>	Flowering	<b>Thrips</b> →  <b>Leaf curling Virus</b> → 	<p>Under present weather condition, for the control of thrips infestation make alternative spray of profenofos 50 EC and Ethion 50 % EC @ 20 ml /10 litre of water at 10 days interval.</p> <p>In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advisedfor control the vector, when sky remains clear.</p>
<b>Lemon</b>	Fruit development	<b>Psylla and thrips</b>  <b>Citrus canker</b> <b>Leaf minor</b>	<p>To prevent damage of psylla&amp;thrips spray <b>imidacloprid 30.5 % 5 ml</b> / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.</p>
<b>Livestock &amp; poultry</b> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Dewormingtesting should be done. Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :94277 28523**



**Agro met Advisory Service bulletin No.-48 for the Devbhumi Dwarka district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 14 to 18-September- 2019)


**Weather forecast until 08-30 hrof 18/09/2019**


**(4) DEVBHUMI DWARKA DISTRICT:**





	14/09/19	15/09/19	16/09/19	17/09/19	18/09/19
Rainfall (mm)	10	0	0	0	0
Max. Temp.(°C)	30	29	30	29	31
Min. Temp.(°C)	27	26	26	26	27
Total CC (octa)	8	8	8	8	7
RH-I (%)	84	87	86	84	85
RH-II (%)	72	79	71	71	70
Wind Speed (kmph)	27	18	22	24	13
Wind Direction(deg.)	262	256	255	254	256

**Agro meteorological Advisories:-**

<b>General Advice</b>	<ul style="list-style-type: none"> <li>• <b>Rain fall:</b> Warm, humid and generally cloudy to cloudy weather &amp; Light rainfall during 14-September forecasted.</li> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 29-31°C and 26-27°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 84-87% &amp; 70-79, respectively.</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W and 13-27 km/h, respectively.</li> <li>• <b>Apply supplementary dose of urea fertilizer as per recommendation of respective crop for optimum growth.</b></li> <li>• <b>Drain out excess water from field for better growth of plant and store in reservoirs.</b></li> <li>• <b>Carry out weeding and plant protection measures as per need.</b></li> </ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/Variety</b>	<b>Advisory</b>
<b>Kharif groundnut</b>	Flowering /pegging	Sclerotium rolfsii	• <b>Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kh/ha as</b>

		<p><b>Tikka &amp; rust</b></p> 	<p><b>preventive measures against Sclerotiumrolfsii stem rot.</b></p> <ul style="list-style-type: none"> <li>• <b>Spray Tebuconazole 10 ml/10 lit of to prevent leaf spot (Tikka &amp; Rust) infestation n groundnut.</b></li> <li>• <b>Carry out weeding at vapsa condition.</b></li> <li>• <b>Spray ferrous sulfate 100 gram &amp; citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop.</b></li> </ul>
<b>Cotton</b>	Squaring	<b>Jassid &amp;Thrips</b>	<p>Monitor the crops for infestation of sucking pest, if require spray <b>imidacloprid 30.5 % 3 ml / 10 lit of water ml/10 lit of water.</b></p> <p>Due to continuous humid weather spray Mancozeb 27 gram/10 lit of water at 15 days interval to prevent leaf spot infestation.</p> <p>Apply 20 kg/ha urea, Ammonium sulphate and Murate of Potash each to cotton crop.</p>
<b>Sesame</b>	Flowering	<p><b>Leaf webber</b></p> <p><b>Phytophthora blight&amp; Powdery mildew</b></p>	<ul style="list-style-type: none"> <li>• <b>For the control of leaf webber spray quinalphos 25 % Ec @ 20 ml/ 10 litre of water.</b></li> <li>• <b>Apply supplementary dose of nitrogen(Urea 54 kg/ha) at 30 DASunder clear sky.</b></li> <li>• <b>Due to warm &amp; humid weather, for prevention of phytophthora blight &amp; powdery mildew in seamum spray Ridomil 20 gm in 10 liters of water.</b></li> <li>• <b>Carry out weeding at vapsa condition.</b></li> </ul>
<b>Pearl millet</b>	Tillering	-	<ul style="list-style-type: none"> <li>• <b>Carry out interculturing&amp;weeding at vapsa condition.</b></li> <li>• <b>Apply supplementary dose of nitrogen(Urea 87 kg/ha) at 30 DASunder clear sky.</b></li> </ul>
<b>Pulses</b>	Flowering		<ul style="list-style-type: none"> <li>• <b>Carry out interculturing&amp;weeding at vapsa condition.</b></li> </ul>
<b>Castor</b>	Germination	GAUCH-1,GC-3, GCH-4, GCH-6, GCH-7	
<b>Fodder Sorghum &amp; Maize</b>	Growing	Sorghum-Gundari, GFS-4 & 5 Maize-African tall, Ganga safed-2 & 5	<p><b>Monitor the crop for shoot fly infestation and if require spraycypermethrin 25 EC 5 ml in 10 liters of water. Infestation of fall army worm observed in maize, so for control make spray of Emamectin Benzoate 7 gram/10 litre of water.</b></p>

<b>Kharif Onion</b>	Growing	Alternaria leaf spot/purple blotch	<p><b>For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb 40 gm and Hexaconazole at 10 days interval.</b></p> <ul style="list-style-type: none"> <li>• <b>Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky.</b></li> </ul>
<b>Vegetables</b>	Transplanting	Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
<b>Lady finger</b>	Flowering/ fruiting	<p><b>Jassids &amp; white fly</b></p> <p><b>Shoot and fruit borer</b></p> <p><b>Mites</b></p>	<p>Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassids control.</p> <p>Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray). If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.</p> <p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear.</p> <p>After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.</p>
<b>Ridge gourd, Bottle gourd, Sponge gourd, Bitter guard &amp; cucurbitaceous crops</b>	Flowering & fruiting	<p><b>Fruit fly</b></p> <p><b>Mites</b></p>	<p>For constant monitoring of cucurbitaceous crops against attack of fruit fly, use of 'Methyl Eugenol trap' is recommended. After detecting the symptoms, spraying of Rogor @ 20 ml + 100 g sugar or jaggary/10 litre water at 50 litres/ha. is advised, when sky is clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites under clear sky.</p>
<b>Cluster bean</b>	Growing/ fruiting	<b>Thrips/whitefly</b>	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 ml or neem oil 40 ml and <b>acetamiprid</b> 20 % SP @ 5.0 gm / 10 lit of water at 15 days interval
<b>Brinjal &amp; Tomato</b>	Flowering/ fruiting	<p><b>Sucking pest</b> →</p> <p><b>&amp; Fruit borer</b></p>  <p><b>Mites</b></p>	<p>Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassids &amp; white fly control.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p> <p>If mite population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when</p>

			sky remains clear.
<b>Chili</b>	Flowering	<b>Thrips</b> →  <b>Leaf curling Virus</b> → 	<p>Under present weather condition, for the control of thrips infestation make alternative spray of profenofos 50 EC and Ethion 50 % EC @ 20 ml /10 litre of water at 10 days interval.</p> <p>In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advisedfor control the vector, when sky remains clear.</p>
<b>Lemon</b>	Fruit development	<b>Psylla and thrips</b>  <b>Citrus canker</b> <b>Leaf minor</b>	<p>To prevent damage of psylla&amp;thrips spray <b>imidacloprid 30.5 % 5 ml</b> / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leafminor.</p>
<b>Livestock &amp; poultry</b> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Dewormingtesting should be done. Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.</p>		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :94277 28523**



**Agro met Advisory Service bulletin No.-48 for the Amreli district**

(Issued jointly by DFERS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 14 to 18-September- 2019)

**Weather forecast until 08-30 hrs of 18/09/2019**


**(5) AMRELI DISTRICT:**

	14/09/19	15/09/19	16/09/19	17/09/19	18/09/19
Rainfall (mm)	70	16	17	10	0
Max.Temp.(°C)	30	29	29	31	31
Min.Temp.(°C)	24	24	25	25	25
Total CC (octa)	8	8	8	7	7
RH-I (%)	92	92	92	90	92
RH-II (%)	76	79	78	72	71
Wind Speed (kmph)	17	12	15	16	10
Wind Direction(deg.)	288	262	253	255	254






**Agro meteorological Advisories:**

<b>General Advice</b>	<p><b>Rain fall: Warm, humid and generally cloudy to cloudy weather &amp; Very heavy rainfall during 14-September &amp; Light to moderate rainfall during 15 to 17-September forecasted.</b></p> <ul style="list-style-type: none"> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 29-31°C and 24-25°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 90-92% &amp; 71-79, respectively.</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W and 10-17 km/h, respectively.</li> <li>• <b>Apply supplementary dose of urea fertilizer as per recommendation of respective crop for optimum growth.</b></li> <li>• <b>Carry out weeding and plant protection measures as per need.</b></li> </ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/Variety</b>	<b>Advisory</b>
<b>Kharif groundnut</b>	Pegging/ pod formation	Sclerotium rolfsii	<ul style="list-style-type: none"> <li>• <b>Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kg/ha as preventive measures against Sclerotium rolfsii stem rot.</b></li> </ul>



		<p><b>Tikka &amp; rust</b></p> 	<ul style="list-style-type: none"> <li>• Spray Tebuconazole 10 ml/10 lit of to prevent leaf spot (Tikka &amp; Rust) infestation n groundnut.</li> <li>• <b>Carry out weeding at vapsa condition.</b></li> <li>• Spray ferrous sulfate 100 gram &amp; citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop.</li> </ul>
<b>Cotton</b>	Square & flower initiation	<p><b>Jassid &amp; Thrips</b></p> <p><b>Pink ball worm</b></p>	<ul style="list-style-type: none"> <li>• Monitor the crops for infestation of sucking pest, if require spray imidacloprid 30.5 % 3 ml / 10 lit of water ml/10 lit of water.</li> <li>• <b>Due to constant moist weather apply supplementary dose of nitrogen through Ammonium sulphate (30 kg/ha) and also apply potash through murate of potash (30 kg/ha) to cotton crop.</b></li> <li>• Also spray hexaconazole 5 % EC @ 15 ml/10 litre of water for control alternaria &amp; cercospora leaf spot in cotton.</li> <li>• <b>Carry out interculturing &amp; weeding at vapsa condition.</b></li> <li>• <b>Also earthing up/ridging should be done in time.</b></li> <li>• For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found <b>pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water.</b></li> </ul>
<b>Sesame</b>	Growing/flowering	<p><b>Leaf webber</b></p> <p><b>Phytophthora blight &amp; Powdery mildew</b></p>	<ul style="list-style-type: none"> <li>• For the control of leaf webber spray quinalphos 25 % Ec @ 20 ml/ 10 litre of water.</li> <li>• <b>Apply supplementary dose of nitrogen (Urea 54 kg/ha) at 30 DAS under clear sky.</b></li> <li>• <b>Due to warm &amp; humid weather, for prevention of phytophthora blight &amp; powdery mildew in seamum spray Ridomil 20 gm in 10 liters of water.</b></li> <li>• <b>Carry out weeding at vapsa condition.</b></li> </ul>

<b>Pearl millet</b>	Milking	-	<ul style="list-style-type: none"> <li>• <b>Carry out weeding at vapsa condition.</b></li> </ul>
<b>Pulses</b>	Pod development & picking	<b>Yellow mosaic viral disease</b>	<ul style="list-style-type: none"> <li>• Due to continuous rainfall infestation of yellow mosaic virus observed in pulse crop, for control remove the infected plants and make spray of Triazophos 40 % EC @ 10 ml/10 litre of water.</li> <li>• <b>Carry out interculturing operation for soil moisture conservation &amp; removal of weed at vapsa condition.</b></li> <li>• <b>Pick mature pod in time.</b></li> </ul>
<b>Kharif Onion</b>	Growing	<b>Alternaria leaf spot/purple blotch</b>	<p><b>For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb 40 gm and Hexaconazole at 10 days interval.</b></p> <ul style="list-style-type: none"> <li>• <b>Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky.</b></li> </ul>
<b>Vegetables</b>	Transplanting/ growing	Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
<b>Lady finger</b>	Flowering/ fruiting	<b>Jassids &amp; white fly</b>  <b>Shoot and fruit borer</b>  <b>Mites</b>	<p>Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassid &amp; white fly control.</p> <p>Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray). If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.</p> <p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear.</p> <p>After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.</p>
<b>Ridge gourd, Bottle gourd, Sponge gourd, Bitter guard &amp; cucurbitace</b>	Flowering & fruiting	<b>Fruit fly</b>  <b>Mites</b>	<p>For constant monitoring of cucurbitaceous crops against attack of fruit fly, use of 'Methyl Eugenol trap' is recommended. After detecting the symptoms, spraying of Rogor @ 20 ml + 100 g sugar or jaggary/10 litre water at 50 litres/ha. is advised, when sky is clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites under clear sky.</p>

<b>ous crops</b>			
<b>Cluster bean</b>	Growing/fruiting	<b>Thrips/whitefly</b>	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
<b>Brinjal&amp; Tomato</b>	Flowering/fruiting	<b>Sucking pest</b> → <b>&amp; Fruit borer</b>  <b>Mites</b>	Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassids & white fly control. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer. If mite population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear.
<b>Chili</b>	Flowering	<b>Thrips</b> →  <b>Leaf curling Virus</b> → 	Under present weather condition, for the control of thrips infestation make alternative spray of profenofos 50 EC and Ethion 50 % EC @ 20 ml /10 litre of water at 10 days interval.  In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear.
<b>Lemon</b>	Fruit development	<b>Psylla and thrips</b>  <b>Citrus canker</b> <b>Leaf minor</b>	To prevent damage of psylla & thrips spray <b>imidacloprid 30.5 % 5 ml</b> / 10 lit of water ml/10 lit of water. Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.
<b>Livestock &amp; poultry</b> 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming testing should be done. Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay +		

	<p>minerals + dry feed. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.</p>
--	--

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : (0281)2784722, Cell:94277 28523**



**Agro met Advisory Service bulletin No.-48 for the Surendranagar district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 14 to 18-September- 2019)


**Weather forecast until 08-30 hrs of 18/09/2019**

**(6) SURENDRANAGAR DISTRICT:**






	14/09/19	15/09/19	16/09/19	17/09/19	18/09/19
Rainfall (mm)	30	4	3	0	0
Max.Temp.(°C)	32	31	32	32	32
Min.Temp.(°C)	24	24	24	24	24
Total CC (octa)	8	8	8	8	7
RH-I (%)	87	86	88	86	87
RH-II (%)	66	70	70	64	64
Wind Speed (kmph)	20	20	19	19	12
Wind Direction(deg.)	258	251	246	251	243

**Agro meteorological advisories:**

<b>General Advice</b>	<p><b>Rain fall: Warm, humid and generally cloudy to cloudy weather &amp; Light to moderate rainfall during 14 to 16-September forecasted.</b></p> <ul style="list-style-type: none"> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 31-32°C and 24-24°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH-I and RH-II is likely to be in the range of 86-88% &amp; 64-70, respectively.</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W, SW and 12-20 km/h, respectively.</li> <li>• <b>Apply supplementary dose of urea fertilizer as per recommendation of respective crop for optimum growth.</b></li> <li>• <b>Carry out weeding and plant protection measures as per need.</b></li> </ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/ Variety</b>	<b>Advisory</b>
<b>Kharif groundnut</b>	Pegging/ pod formation	<b>Sclerotium rolfsii</b>	<ul style="list-style-type: none"> <li>• <b>Apply 2.5 kg/ha Trichoderma fungus with well decomposed FYM OR castor cake @ 500 kg/ha as preventive measures against Sclerotium rolfsii stem rot.</b></li> </ul>

		<p><b>Tikka &amp; rust</b></p> 	<ul style="list-style-type: none"> <li>• Spray Tebuconazole 10 ml/10 lit of to prevent leaf spot (Tikka &amp; Rust) infestation n groundnut.</li> <li>• <b>Carry out weeding at vapsa condition.</b></li> <li>• Spray ferrous sulfate 100 gram &amp; citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop.</li> </ul>
<b>Cotton</b>	Square & flower initiation	<p><b>Jassid &amp; Thrips</b></p> <p><b>Pink ball worm</b></p>	<ul style="list-style-type: none"> <li>• Monitor the crops for infestation of sucking pest, if require spray imidacloprid 30.5 % 3 ml / 10 lit of water ml/10 lit of water.</li> <li>• <b>Due to constant moist weather apply supplementary dose of nitrogen through Ammonium sulphate (30 kg/ha) and also apply potash through murate of potash (30 kg/ha) to cotton crop.</b></li> <li>• Also spray hexaconazole 5 % EC @ 15 ml/10 litre of water for control alternaria &amp; cercospora leaf spot in cotton.</li> <li>• <b>Carry out interculturing &amp; weeding at vapsa condition.</b></li> <li>• <b>Also earthing up/ridging should be done in time.</b></li> <li>• For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found <b>pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water.</b></li> </ul>
<b>Sesame</b>	Growing/flowering	<p><b>Leaf webber</b></p> <p><b>Phytophthora blight &amp; Powdery mildew</b></p>	<ul style="list-style-type: none"> <li>• For the control of leaf webber spray quinalphos 25 % Ec @ 20 ml/ 10 litre of water.</li> <li>• Apply supplementary dose of nitrogen (Urea 54 kg/ha) at 30 DAS under clear sky.</li> <li>• Due to warm &amp; humid weather, for prevention of phytophthora blight &amp; powdery mildew in seamum spray Ridomil 20 gm in 10 liters of water.</li> <li>• <b>Carry out weeding at vapsa condition.</b></li> </ul>

<b>Pearl millet</b>	Milking	-	<ul style="list-style-type: none"> <li>• <b>Carry out weeding at vapsa condition.</b></li> </ul>
<b>Pulses</b>	Pod development & picking	<b>Yellow mosaic viral disease</b>	<ul style="list-style-type: none"> <li>• Due to continuous rainfall infestation of yellow mosaic virus observed in pulse crop, for control remove the infected plants and make spray of Triazophos 40 % EC @ 10 ml/10 litre of water.</li> <li>• <b>Carry out interculturing operation for soil moisture conservation &amp; removal of weed at vapsa condition.</b></li> <li>• <b>Pick mature pod in time.</b></li> </ul>
<b>Kharif Onion</b>	Growing	<b>Alternaria leaf spot/purple blotch</b>	<p><b>For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb 40 gm and Hexaconazole at 10 days interval.</b></p> <ul style="list-style-type: none"> <li>• <b>Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky.</b></li> </ul>
<b>Vegetables</b>	Transplanting/ growing	Transplanting of brinjal, chilli and early cauliflower are advised for those farmers whose seedlings are ready.	
<b>Lady finger</b>	Flowering/ fruiting	<b>Jassids &amp; white fly</b>  <b>Shoot and fruit borer</b>  <b>Mites</b>	<p>Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassid &amp; white fly control.</p> <p>Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray). If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.</p> <p>Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear.</p> <p>After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done.</p>
<b>Ridge gourd, Bottle gourd, Sponge gourd, Bitter guard &amp; cucurbitace</b>	Flowering & fruiting	<b>Fruit fly</b>  <b>Mites</b>	<p>For constant monitoring of cucurbitaceous crops against attack of fruit fly, use of 'Methyl Eugenol trap' is recommended. After detecting the symptoms, spraying of Rogor @ 20 ml + 100 g sugar or jaggery/10 litre water at 50 litres/ha. is advised, when sky is clear.</p> <p>If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites under clear sky.</p>

<b>ous crops</b>			
<b>Cluster bean</b>	Growing/fruiting	<b>Thrips/whitefly</b>	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
<b>Brinjal&amp; Tomato</b>	Flowering/fruiting	<b>Sucking pest</b> → <b>&amp; Fruit borer</b>  <b>Mites</b>	Spray 500 ml of 5 % <b>neem seed extract</b> or neem seed oil 50 ml with with <b>Trizophos</b> 40 % EC @ 25 ml or <b>imidacloprid 30.5 SC</b> 3 ml/10 lit of water for jassids & white fly control. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer. If mite population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear.
<b>Chili</b>	Flowering	<b>Thrips</b> →  <b>Leaf curling Virus</b> → 	Under present weather condition, for the control of thrips infestation make alternative spray of profenofos 50 EC and Ethion 50 % EC @ 20 ml /10 litre of water at 10 days interval.  In present weather conditions, rouging of viral infested crop should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 5 ml per 10litre of water is advised for control the vector, when sky remains clear.
<b>Lemon</b>	Fruit development	<b>Psylla and thrips</b>  <b>Citrus canker</b> <b>Leaf minor</b>	To prevent damage of psylla & thrips spray <b>imidacloprid 30.5 % 5 ml</b> / 10 lit of water ml/10 lit of water. Spray copper oxychloride 40 gm or streptomycin 1.5 gm in 10 litre of water Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.
<b>Livestock &amp; poultry</b> 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming testing should be done. Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay +		



	<p>minerals + dry feed. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Due to chance of rainfall at isolated places keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall.</p>
--	--

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**