



ग्रामीण कृषि मौसम सेवा
हवामान आगाही आधारित कृषिसलाह बुलेटीन
उत्तर सौराष्ट्रभेत आबोहवाकीय विभाग
ता.१८-०१-२०२० थी ता.२२-०१-२०२०



(ग्रामीण कृषि मौसम सेवा, सूकी भेती संशोधन केन्द्र, जू.कृ.यु.,तरघडिया अने भारत मौसम विभाग द्वारा संपन्न)

कृषि हवामान बुलेटीन नं.८४/२०२०

पाछवा अठवाडीयानुं राजकोटनुं हवामान

हवामान परिबणो	
वरसाह(मी.मी)	0.0
महतम तापमान (से.ग्रे)	20.6-30.9
लघुतम तापमान (से.ग्रे)	5.4-15.9
लेज(%) सवार	63-88
लेज(%) बपोरबाह	30-58
पवननी गति (किमी/कलाक)	3.0-5.6

हवामान आगाही: पत्रक-क

उत्तरसौराष्ट्र भेत आबोहवाकीय विभाग-हवामान आगाही

ता.१८-०१-२०२० थी २२-०१-२०२०

कृषि सलाह:

સામાન્ય

➤ ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ,જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડીયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના

રાજકોટ: જીલ્લામાં આગામી તા.૧૮-૦૧-૨૦૨૦ થી તા.૨૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સૂકું, ઠંડુ અને મુખત્વે વાદળછાયું રહશે. પવનની દિશા વાયવ્ય, ઈશાનની રહેવાની અને પવનની ઝડપ **૧૦ થી ૧૮ કીમી/કલાક** રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **૨૨-૨૬** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન **૯-૧૦** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૮-૦૧-૨૦	૧૯-૦૧-૨૦	૨૦-૦૧-૨૦	૨૧-૦૧-૨૦	૨૨-૦૧-૨૦
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	22	23	24	26	26
લઘુત્તમતાપમાન (સે.ગ્રે)	9	9	9	10	10
વાદળનીસ્થિતિ(ઓકટા)	0	0	1	3	2
મહત્તમભેજ(%) સવાર	52	38	40	42	43
લઘુત્તમભેજ (%) બપોરબાદ	25	17	24	23	16
પવનનીગતિ(કિમી/કલાક)	11	18	13	10	15
પવનનીદિશા(ડીગ્રી)	56	25	29	63	41

મોરબી :જીલ્લામાં આગામી તા.૧૮-૦૧-૨૦૨૦ થી તા.૨૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સૂકું, ઠંડુ અને મુખત્વે વાદળછાયું રહશે. પવનની દિશા અગ્નિ, ઈશાન અને પૂર્વની રહેવાની અને પવનની ઝડપ **૫ થી ૧૪ કીમી/કલાક** રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **૨૨-૨૬** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન **૯-૧૦** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૮-૦૧-૨૦	૧૯-૦૧-૨૦	૨૦-૦૧-૨૦	૨૧-૦૧-૨૦	૨૨-૦૧-૨૦
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	22	23	24	26	26
લઘુત્તમતાપમાન (સે.ગ્રે)	9	9	9	10	10
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	3	4
મહત્તમભેજ(%) સવાર	63	55	54	52	46
લઘુત્તમભેજ (%) બપોરબાદ	32	25	25	25	18
પવનનીગતિ(કિમી/કલાક)	5	14	13	5	7
પવનનીદિશા(ડીગ્રી)	134	41	55	72	92

જામનગર: જીલ્લામાં આગામી તા.૧૮-૦૧-૨૦૨૦ થી તા.૨૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સૂકું, ઠંડુ અને મુખત્વે વાદળછાયું રહશે. પવનની દિશા ઈશાનની રહેવાની અને પવનની ઝડપ **૧૩ થી ૨૪**

કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **૨૩-૨૬** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન **૧૧-૧૪** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૮-૦૧-૨૦	૧૯-૦૧-૨૦	૨૦-૦૧-૨૦	૨૧-૦૧-૨૦	૨૨-૦૧-૨૦
વરસાદ(મી.મી)	૦	૦	૦	૦	૦
મહત્તમતાપમાન (સે.ગ્રે)	૨૩	૨૪	૨૪	૨૫	૨૬
લઘુત્તમતાપમાન (સે.ગ્રે)	૧૧	૧૨	૧૨	૧૩	૧૪
વાદળનીસ્થિતિ(ઓકટા)	૦	૦	૧	૨	૧
મહત્તમભેજ(%) સવાર	૪૫	૪૨	૪૬	૪૮	૪૬
લઘુત્તમભેજ (%) બપોરબાદ	૨૬	૨૬	૩૧	૩૦	૨૭
પવનનીગતિ(કિમી/કલાક)	૧૩	૨૪	૧૯	૧૩	૨૧
પવનનીદિશા(ડીગ્રી)	૩૨	૩૪	૩૭	૨૪	૨૮

દેવભૂમિ દ્વારકા: જીલ્લામાં આગામી તા.૧૮-૦૧-૨૦૧૯ થી તા.૨૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને મુખત્વે વાદળછાયું રહેશે. પવનની દિશા ઈશાનની રહેવાની અને પવનની ઝડપ **૧૨ થી ૨૨ કીમી/કલાક** રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **૨૨-૨૫** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન **૧૨-૧૫** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૮-૦૧-૨૦	૧૯-૦૧-૨૦	૨૦-૦૧-૨૦	૨૧-૦૧-૨૦	૨૨-૦૧-૨૦
વરસાદ(મી.મી)	૦	૦	૦	૦	૦
મહત્તમતાપમાન (સે.ગ્રે)	૨૨	૨૩	૨૩	૨૪	૨૫
લઘુત્તમતાપમાન (સે.ગ્રે)	૧૨	૧૩	૧૩	૧૪	૧૫
વાદળનીસ્થિતિ(ઓકટા)	૦	૦	૧	૨	૧
મહત્તમભેજ(%) સવાર	૪૨	૪૩	૪૯	૫૦	૪૭
લઘુત્તમભેજ (%) બપોરબાદ	૨૫	૨૭	૨૯	૨૮	૨૬
પવનનીગતિ(કિમી/કલાક)	૧૨	૨૨	૧૮	૧૩	૧૯
પવનનીદિશા(ડીગ્રી)	૨૮	૩૭	૪૨	૨૮	૨૮

અમરેલી: જીલ્લામાં આગામી તા.૧૮-૦૧-૨૦૨૦ થી તા.૨૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને મુખત્વે વાદળછાયું રહેશે. પવનની દિશા અગ્નિ, ઈશાન અને પૂર્વની રહેવાની અને પવનની ઝડપ **૧૨ થી ૨૧ કીમી/કલાક** રહેની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **૨૩-૨૭** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન **૧૦-૧૧** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.


તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૮-૦૧-૨૦	૧૯-૦૧-૨૦	૨૦-૦૧-૨૦	૨૧-૦૧-૨૦	૨૨-૦૧-૨૦



વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	23	24	25	27	27
લઘુત્તમતાપમાન (સે.ગ્રે)	10	10	10	11	11
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	1	2
મહત્તમભેજ(%) સવાર	52	39	44	48	44
લઘુત્તમભેજ(%) બપોરબાદ	27	21	29	26	20
પવનનીગતિ(કિમી/કલાક)	15	21	17	12	16
પવનનીદિશા(ડીગ્રી)	115	33	88	112	53








સુરેન્દ્રનગર: જીલ્લામાં આગામી તા.૧૮-૦૧-૨૦૨૦ થી તા.૨૨-૦૧-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને મુખત્વે વાદળછાયું રહશે. પવનની દિશા પૂર્વ, ઈશાન અને અગ્નિની રહેવાની અને વનની ઝડપ ૧૧ થી ૧૮ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૨-૨૬ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૯-૧૦ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.



તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૧૮-૦૧-૨૦	૧૯-૦૧-૨૦	૨૦-૦૧-૨૦	૨૧-૦૧-૨૦	૨૨-૦૧-૨૦
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમતાપમાન (સે.ગ્રે)	22	23	24	26	26
લઘુત્તમતાપમાન (સે.ગ્રે)	9	9	9	10	10
વાદળનીસ્થિતિ(ઓકટા)	0	0	1	3	3
મહત્તમભેજ(%) સવાર	54	41	42	46	43
લઘુત્તમભેજ(%) બપોરબાદ	29	22	28	27	19
પવનનીગતિ(કિમી/કલાક)	13	18	13	11	17
પવનનીદિશા(ડીગ્રી)	88	22	31	126	118








➤ ૧૮ થી ૨૨-જાન્યુઆરી-૨૦૨૦ દરમ્યાન હવામાન સુકું, ઠંડુ અને મુખત્વે વાદળછાયું રહશે. પવનની દિશા ઈશાનની મોટાભાગે રહેવાની અને પવનની ઝડપ ૫ થી ૨૪ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૨-૨૭ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૯-૧૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.



મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટરપાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
વસણ	વૃદ્ધિ	શ્રીપ્સ, 	<ul style="list-style-type: none"> ➤ વસણના પાકમાં શ્રીપ્સના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને પ્રોફેનોફોસ ૩૦ મિલી અથવા લેમ્બડા સાયહેલોથ્રીન ૧૦ મિલી દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. અતિ ઉપદ્રવ હોય તો સ્પીનોસાઇ ૪ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ વસણના પાકમાં પાન પીળા પડતા અટકાવવા મેન્ડોઝેબ ૨૭ ગ્રામ

			<p>અથવા કાર્બેન્ડાઝીમ ૧૦ ગ્રામ પ્રતિ ૧૦ લીટર પાણીમાં નાખીને છંટકાવ કરવો.</p> <ul style="list-style-type: none"> ➤ જમીનની પ્રત મુજબ ૧૦-૧૨ દિવસે પિયત આપવું. યુરિઆ ખાતર ૫૪ કી.ગ્રા. પ્રતિ હેક્ટર વાવેતર બાદ ૩૦-૩૫ દિવસે આપવું. ➤ વસણ પીળું પડે નહિ તે માટે વધારે પડતું પિયત આપવું નહિ.
ચણા	વૃદ્ધિ/ફૂલ/પોપટા	<p>લીલી અને લશ્કરી ઈયળ</p> 	<p>ચણાના પાકમાં લીલી અને લશ્કરી ઈયળના નિયંત્રણ માટે પક્ષીઓને બેસવા માટે ' ' T ' આકારના બેલાખડા મુકવા. જો વધુ ઉપદ્રવ જણાય તો કવીનાલફોસ ૨૦ મિલી અથવા એમામેક્ટીન બેન્ઝોએટ ૬.૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો. તેમજ લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.</p> <p>પાનના સુકારાના નિયંત્રણ માટે ડાયથેન એમ ૪૫, ૨૭ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</p>
ધાણા	ફૂલ	<p>થ્રીપ્સ અને મોલો</p> <p>રાખોડીના નિયંત્રણ</p>	<p>થ્રીપ્સ અને મોલોના ના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.</p> <p>રાખોડીના નિયંત્રણ હેક્ઝાકોનાઝોલ ૧૫ મિલી દવા ૧૦ લીટર પાણીમાં મિશ્ર કરીને ૧૫ દિવસના અંતરે છંટકાવ કરવો.</p>

જીરું	ફૂલ	<p>શ્રીપ્સ & મોલો</p> <p>ચરમી અને રાખોડી</p>  <p>સુકારો</p> 	<p>જીરુંના પાકમાં શ્રીપ્સ અને મોલોના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને એસીટામીપ્રીડ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.</p> <p>જીરુંના પાકમાં કાળીચરમી અને રાખોડીનો ઉપદ્રવ ન થાય તે માટે ડાયથેન એમ ૪૫, ૨૭ ગ્રામ અને હેક્ઝાકોનાઝોલ ૧૫ મિલી દવા વારાફરતી ૧૦ લીટર પાણીમાં મિશ્ર કરીને ૧૫ દિવસના અંતરે છંટકાવ કરવો.</p>  <p>જીરુંના ઉભા પાકમાં સુકારાના નિયંત્રણ માટે પિયત આપવાનું બંધ કરવું.</p> 
ઘઉં	ફૂલ	-	પિયત સમયસર આપવું.
રાયડો	શીંગ	શ્રીપ્સ અને મોલો	શ્રીપ્સ અને મોલોના ના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.
કપાસ	કપાસની વીણી	<p>ગુલાબી ઇયળ</p> <p>ગુલાબી ઇયળનાં નિયંત્રણ માટે તેમજ પિયત આપવું નહિ. બીજો ફાલ લેવો નહિ. છેલ્લી વીણી પછી ઘેટા બકરા ચરાવવા. જેથી ખુલ્યા વગરના ગુલાબી ઇયળના નુકશાન વાળા જીંડવા અને અન્ય કચરો નાશ પામે. ત્યારબાદ કરાંઠી સળગાવવી નહિ પરંતુ રોટાવેટર/મોબાઈલ ચોપરથી નાના ટુકડા કરી જમીનમાં ઢાંટી દેવા જેથી જમીન સેન્દ્રિય તત્વનો ઉમેરો થવાથી સુંક્ષ્મ જીવોની સંખ્યા વધશે અને જમીનના સ્વાસ્થ્યમાં સુધારો થશે. અથવા કરાંઠી ભેગી કરી શ્રેદરથી નાના ટુકડા કરીને છાણ, યુરીયા, કમ્પોસ્ટ કલ્ચર ઉમેરી દેશી(સેન્દ્રિય) ખાતર બનાવવું.</p>	  

દિવેલા	વૃદ્ધિ/ફૂલ/ગડ	લીલી પોપટી, સફેદ માખી ગડની ઇયળ	<ul style="list-style-type: none"> ➤ સફેદ માખી અને લીલી પોપટીનો વધુ ઉપદ્રવ જોવા મળે છે. જેના નિયંત્રણ માટે એસીટામાપ્રોડ ૨૦ ટકા ૫ ગ્રામ અને ઈમીડાક્લોપ્રોડ ૩૦.૫ % ૩ મિલી સાથે લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ ગડની ઇયળના નિયંત્રણ માટે ક્લોરાનટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઈસી ૭ મિલી ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
ડુંગળી	રોપણી/વૃદ્ધિ વિકાસ	શ્રીપ્સ, પાનનો સુકારો જાંબલી ધાબા 	<ul style="list-style-type: none"> ➤ શ્રીપ્સના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને પ્રોફેનોફોસ ૩૦ મિલી અથવા લેમ્બડા સાયહેલોથ્રીન ૧૫ મિલી દવાને ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. અતિ ઉપદ્રવ હોય તો સ્પીનોસાદ ૪ મિલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ ડુંગળીના પાકમાં જાંબલી ધાબાના રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા. ➤ નાઈટ્રોજન યુક્ત એમોનિયમ સલ્ફેટ ખાતરનો પુરતી ડોઝ ૩૦ કીલો ગ્રામ પ્રતિ હે. આપવો.
તુવેર	ફૂલ/શીંગ/દાણા ભરાવવા	શીંગ માખી	ક્વીનાલફોસ ૨૦ મિલી અથવા ડીડીવીપી ૫ મિલી ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. તેમજ લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લિટર પાણીમાં કરીને છંટકાવ કરવો.
કોબી અને ફલાવર	વૃદ્ધિ/દડા	હિરા ફૂદુ	કોબી અને ફલાવરમાં હિરાફૂદાના નિયંત્રણ માટે સ્પીનોસાદ ૩ મિલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
રીંગણી અને ટમેટી	ફૂલ/ફળ	તડતડીયા અને સફેદ માખી ડુંખ અને→ ફળ કોરી ખાનાર ઇયળ પાન કથીરી	<ul style="list-style-type: none"> ➤ તડતડીયા અને સફેદ માખીના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ડાયફેન્યાયુરોન ૫૦ ટકા વે.પા. ૧૬ ગ્રામ અથવા ટ્રાયઝોફોસ ૪૦ ઈસી ૨૫ મિ.લી. દવાને ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ ફળ અને ડોકાની ઇયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને ક્લોરાનટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઈસી મિલી ૧૦ લિટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ જો પાનકથીરીનો ઉપદ્રવ હોય તો પ્રોપરગાઈટ ૫૭ ઈ.સી. ૧૫ મિલી પ્રતિ પંપ છંટકાવ કરવો.
મરચી	વૃદ્ધિ/ફૂલ/ફળ	શ્રીપ્સ, સફેદ માખી  કાલવ્રણ →	<ul style="list-style-type: none"> ➤ પ્રોફેનોફોસ ૨૦ મીલી અને લીંબોળીનું તેલ ૫૦ મિલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. શ્રીપ્સનો વધુ ઉપદ્રવ હોય તો સ્પીનોસાદ અથવા સ્પીનેટોરમ દવા ૫ મિલી. પ્રતિ પંપ છાંટવી. ➤ કાલવ્રણ રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.

			
લીબુ	ફળનો વિકાસ	પાન કથીરી   સાયલા અને શ્રીપ્સ બદામી ટપકાં પાનકોરીયુ	લીબુના બગીચામાં પાન કથીરીનો ઉપદ્રવ વધુ હોય તો તેના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. લીબુમાં સાયલા અને શ્રીપ્સ તેના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૫ મીલી અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. વધુ ઉપદ્રવ હોય તો ટ્રાયઝોફોસ ૪૦ ટકા ૨૦/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.    બદામી ટપકાના નિયંત્રણ માટે કોપર ઓક્ઝીક્લોરાઈડ ૪૦ ગ્રામ અને ૧.૫ ગ્રામ સ્ટ્રેપ્ટોસાયક્લીનના બે છંટકાવ ૧૫ દિવસના ગળે કરવા. પાનકોરીયુના નિયંત્રણ માટે ડીડીવીપી ૧૦ મી.લી. ૧૦ લિટર પાણીમાં મિશ્ર કરીને છાંટવી.
આંબો	ફૂલ	મધીયો અને મગીયાની ઈયળ ભુકીછારો/રાખોડી	આંબામાં મધીયો અને મગીયાની ઈયળના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૫ મિલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.  આંબામાં ભુકીછારો/રાખોડીના નિયંત્રણ માટે હેક્ઝાકોનાઝોલ ૧૫ મિલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.

				
<p>પશુ/ મરઘા</p> 	<p>પ્રજીવથી થતા રોગોની ચકાસણી કરાવવી. ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મી.લી. ૧ લીટર પાણીમાં નાખીને છાંટવી. વિટામીન એ તથા ડી પુરકો આપવા. કરમિયાની દવા આપવી તથા ઉછરતા પશુઓમાં બૃસેલ્વા કાફવુંડનું રસીકરણ કરાવવું. પશુને ૬ થી ૮ કી.ગ્રા. સુકો અને ૧૫ થી ૨૦ કી.ગ્રા. લીલો ચારો આપવો. દુધાળા પશુઓને પ્રતિ લીટર દુધની ઊપજ મુજબ નિયમિત ૧ કી.ગ્રા. ફીડ + ૫૦ ગ્રામ ખનીજ તત્ત્વોનું મિશ્રણ આપવું. દુધાળા પશુઓને હુંફાળું ગરમ પાણી પીવા માટે આપવું. તેમજ માખીઓના ઉપદ્રવથી બચવા ફીનાઈલ છાંટવું.</p>			

જુફ્યુ/ટેકનિકલ ઓફિસર.(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૨૦
તરઘડિયા(રાજકોટ), તારીખ: ૧૭/૦૧/૨૦૨૦.

પ્રતિ,
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર
સુકી ખેતીસંશોધન કેન્દ્ર
જુનાગઢ કૃષિ નિવર્સિટી
તરઘડીયા



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94269 38235



Agro met Advisory Service bulletin No.84-for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
(Period 18 to 22-January- 2020)

Significant past weather for the preceding week of Rajkot




Rainfall (mm)	0.0
Maximum temperature (°C)	20.6-30.9
Minimum temperature (°C)	5.4-15.9
Relative Humidity (%) I	63-88
Relative Humidity (%) II	30-58
Wind speed (Kmph)	3.0-5.6





Weather forecast until 08-30 hrs. of 22/01/2020




(1) RAJKOT DISTRICT:







	18/01/20	19/01/20	20/01/20	21/01/20	22/01/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	22	23	24	26	26
Min.Temp.(°C)	9	9	9	10	10
Total CC (octa)	0	0	1	3	2
RH-I (%)	52	38	40	42	43
RH-II (%)	25	17	24	23	16
Wind Speed (kmph)	11	18	13	10	15
Wind Direction(deg.)	56	25	29	63	41

Agro meteorological Advisories:

<p>General Advice</p>	<ul style="list-style-type: none"> • Rain fall: Dry, cold and Mainly clear to Partly cloudy weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 22-26°C and 9-10°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 38-52% & 16-25, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be NE and 10-18 km/h, respectively. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
<p>Garlic</p>	<p>Girthing</p>	<p>Thrips and aphid</p>  <p>leaf blight (yellowing)</p> 	<p>For thrips & aphid control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml with imidacloprid 17.8 % EC 5 ml in 10 liters of water.</p> <p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
<p>Chick pea</p>	<p>Flowering/pod formation</p>	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.

<p>Cumin</p>	<p>Growth/Flowering</p>	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm or imidacloprid 17.8 % 5 ml/10 liters of water.</p> <p>Under present weather condition spray tebuconazole 10 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
<p>Coriander</p>	<p>Flowering</p>	<p>Thrips and Jassids</p>	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>
<p>Wheat</p>	<p>Flowering</p>	<p>-</p>	<p>Apply irrigation at critical stage without stress.</p>
<p>Cotton</p>	<p>Picking & pulling of cotton stalk</p>	<p>Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time.</p> <p>Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by thedoin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.</p>	

			
Castor	Flowering/ capsule formation	Jassids& white fly→ Pod borer	<p>Spray Acetamiprid 20 % SP@ 5.0 gm Imidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water.</p> <p>For the control of pod borer spray chlorantranilprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p>
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	Thrips Leaf blight→ 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Make alternate spray of Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.</p>
Brinjal& Tomato	Flowering/ fruiting	<p>Sucking pest→</p> <p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>Monitor the crop for infestation of jassid& white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantranilprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infetation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>

<p>Chili</p>	<p>Flowering & fruiting</p>	<p>Thrips → </p> <p>Anthracnose → </p>	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
<p>Lemon</p>	<p>Flowering and fruit formation</p>	<p>Psylla and thrips</p> <p>Citrus canker</p> <p>Leaf minor</p>	<p></p> <p>To prevent damage of psylla&thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
<p>Mango</p>	<p>Flowering</p>	<p>Blossom midge and Leaf hopper → </p> <p>Powdery Mildew → </p>	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p> <p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>
<p>Livestock & poultry </p>	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib</p>		

	warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.
--	--

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-84 for the Morbi district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 18 to 22-January- 2020)







Weather forecast until 08-30 hrs. Of 22/01/2020



(2) MORBI DISTRICT:






	18/01/20	19/01/20	20/01/20	21/01/20	22/01/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	22	23	24	26	26
Min. Temp.(°C)	9	9	9	10	10
Total CC (octa)	0	0	0	3	4
RH-I (%)	63	55	54	52	46
RH-II (%)	32	25	25	25	18
Wind Speed (kmph)	5	14	13	5	7
Wind Direction(deg.)	134	41	55	72	92



Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and Partly cloudy to clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 22-26°C and 9-10°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 46-63% & 18-32, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be SE, NE, E and 5-14 km/h, respectively. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Garlic	Girthing	Thrips and aphid  leaf blight (yellowing)	<p>For thrips & aphid control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml with imidacloprid 17.8 % EC 5 ml in 10 liters of water.</p> <p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p>

			Do not over irrigate to prevent leaf yellowing disease.
Chick pea	Flowering/pod formation	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.
Cumin	Growth/Flowering	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm or imidacloprid 17.8 % 5 ml/10 liters of water.</p> <p>Under present weather condition spray tebuconazole 10 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
Coriander	Flowering	Thrips and Jassids	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water.

			Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.
Wheat	Flowering	-	Apply irrigation at critical stage without stress.
Cotton	Picking & pulling of cotton stalk	<p>Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time.</p> <p>Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by the doin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.</p>	
Castor	Flowering/ capsule formation	<p>Jassids& white fly→</p> <p>Pod borer</p>	<p>Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water.</p> <p>For the control of pod borer spray chlorantraniliprole18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p>
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	<p>Thrips</p> <p>Leaf blight→</p> 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Make alternate sprayof Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.</p>
Brinjal& Tomato	Flowering/ fruiting	Sucking pest →	Monitor the crop for infestation of jassid& white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with

		<p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p>  <p>Anthracnose →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
Lemon	Flowering and fruit formation	<p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p>	<p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water. Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
Mango	Flowering	<p>Blossom midge and Leaf hopper →</p>  <p>Powdery Mildew →</p>	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p> <p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>

			
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-84 for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 18 to 22-January-2020)







Weather forecast until 08-30 hrs. of 22/01/2020



(3) JAMNAGAR DISTRICT:






	18/01/20	19/01/20	20/01/20	21/01/20	22/01/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	23	24	24	25	26
Min. Temp.(°C)	11	12	12	13	14
Total CC (octa)	0	0	1	2	1
RH-I (%)	45	42	46	48	46
RH-II (%)	26	26	31	30	27
Wind Speed (kmph)	13	24	19	13	21
Wind Direction(deg.)	32	34	37	24	28



Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and Mainly clear to clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 23-26°C and 11-14°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 42-48% & 26-31, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be NE and 13-24 km/h, respectively. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Garlic	Girthing	Thrips and aphid  leaf blight (yellowing)	<p>For thrips & aphid control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml with imidacloprid 17.8 % EC 5 ml in 10 liters of water.</p> <p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water</p>

			<p>at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
Chick pea	Flowering/pod formation	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.
Cumin	Growth/Flowering	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm or imidacloprid 17.8 % 5 ml/10 liters of water.</p> <p>Under present weather condition spray tebuconazole 10 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
Coriander	Flowering	Thrips and Jassids	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water.</p>

			Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.
Wheat	Flowering	-	Apply irrigation at critical stage without stress.
Cotton	Picking & pulling of cotton stalk	<p>Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time.</p> <p>Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by the doin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.</p>	
Castor	Flowering/ capsule formation	<p>Jassids& white fly→</p> <p>Pod borer</p>	<p>Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water.</p> <p>For the control of pod borer spray chlorantraniliprole18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p>
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	<p>Thrips</p> <p>Leaf blight→</p> 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Make alternate sprayof Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.</p>
Brinjal& Tomato	Flowering/ fruiting	Sucking pest →	Monitor the crop for infestation of jassid& white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with

		<p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p>  <p>Anthracnose →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
Lemon	Flowering and fruit formation	<p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p>	<p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water. Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
Mango	Flowering	<p>Blossom midge and Leaf hopper →</p>  <p>Powdery Mildew →</p>	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p> <p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>

			
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-84 for the Devbhumi Dwarka district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 18 to 22-January-2020)







Weather forecast until 08-30 hrs. of 22/01/2020



(4) DEVBHUMI DWARKA DISTRICT:






	18/01/20	19/01/20	20/01/20	21/01/20	22/01/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	22	23	23	24	25
Min. Temp.(°C)	12	13	13	14	15
Total CC (octa)	0	0	1	2	1
RH-I (%)	42	43	49	50	47
RH-II (%)	25	27	29	28	26
Wind Speed (kmph)	12	22	18	13	19
Wind Direction(deg.)	28	37	42	28	28



Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and Mainly clear to clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 22-25°C and 12-15°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 42-50% & 25-29, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be NE and 12-22 km/h, respectively. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Garlic	Girthing	Thrips and aphid  leaf blight (yellowing)	<p style="color: red;">For thrips & aphid control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml with imidacloprid 17.8 % EC 5 ml in 10 liters of water.</p> <p style="color: green;">To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water</p>

			<p>at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
Chick pea	Flowering/pod formation	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.
Cumin	Growth/Flowering	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm or imidacloprid 17.8 % 5 ml/10 liters of water.</p> <p>Under present weather condition spray tebuconazole 10 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
Coriander	Flowering	Thrips and Jassids	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water.</p>

			Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.
Wheat	Flowering	-	Apply irrigation at critical stage without stress.
Cotton	Picking & pulling of cotton stalk	<p>Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time.</p> <p>Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by thedoin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.</p>	
Castor	Flowering/ capsule formation	<p>Jassids& white fly→</p> <p>Pod borer</p>	<p>Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water.</p> <p>For the control of pod borer spray chlorantraniliprole18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p>
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	<p>Thrips</p> <p>Leaf blight→</p> 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Make alternate sprayof Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.</p>
Brinjal& Tomato	Flowering/ fruiting	Sucking pest →	Monitor the crop for infestation of jassid& white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with

		<p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p>  <p>Anthracnose →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
Lemon	Flowering and fruit formation	<p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p>	<p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water. Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
Mango	Flowering	<p>Blossom midge and Leaf hopper →</p>  <p>Powdery Mildew →</p>	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p> <p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>

			
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-84 for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 18 to 22-January-2020)







Weather forecast until 08-30 hrs. of 22/01/2020



(5) AMRELI DISTRICT:






	18/01/20	19/01/20	20/01/20	21/01/20	22/01/20
Rainfall (mm)		0	0	0	0
Max.Temp.(°C)	23	24	25	27	27
Min.Temp.(°C)	10	10	10	11	11
Total CC (octa)	0	0	0	1	2
RH-I (%)	52	39	44	48	44
RH-II (%)	27	21	29	26	20
Wind Speed (kmph)	15	21	17	12	16
Wind Direction(deg.)	115	33	88	112	53



Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and Mainly clear to clear weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 23-27°C and 10-11°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 39-52% & 20-29, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be SE, NE, E and 12-21 km/h, respectively. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Garlic	Girthing	Thrips and aphid  leaf blight (yellowing)	<p>For thrips & aphid control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml with imidacloprid 17.8 % EC 5 ml in 10 liters of water.</p> <p>To prevent leaf blight (yellowing) disease infestation in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p>

			Do not over irrigate to prevent leaf yellowing disease.
Chick pea	Flowering/pod formation	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.
Cumin	Growth/Flowering	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm or imidacloprid 17.8 % 5 ml/10 liters of water.</p> <p>Under present weather condition spray tebuconazole 10 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
Coriander	Flowering	Thrips and Jassids	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water.

			Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.
Wheat	Flowering	-	Apply irrigation at critical stage without stress.
Cotton	Picking & pulling of cotton stalk	<p>Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time.</p> <p>Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by the doin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.</p>	
Castor	Flowering/ capsule formation	<p>Jassids& white fly→</p> <p>Pod borer</p>	<p>Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water.</p> <p>For the control of pod borer spray chlorantraniliprole18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p>
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	<p>Thrips</p> <p>Leaf blight→</p> 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Make alternate sprayof Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.</p>
Brinjal& Tomato	Flowering/ fruiting	Sucking pest →	Monitor the crop for infestation of jassid& white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with

		<p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p>  <p>Anthracnose →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
Lemon	Flowering and fruit formation	<p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p>	<p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water. Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
Mango	Flowering	<p>Blossom midge and Leaf hopper →</p>  <p>Powdery Mildew →</p>	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p> <p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>

			
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:9426938235



Agro met Advisory Service bulletin No.-84 for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 18 to 22-January-2020)







Weather forecast until 08-30 hrs of 22/01/2020



(6) SURENDRANAGAR DISTRICT:






	18/01/20	19/01/20	20/01/20	21/01/20	22/01/20
Rainfall (mm)		0	0	0	0
Max.Temp.(°C)	22	23	24	26	26
Min.Temp.(°C)	9	9	9	10	10
Total CC (octa)	0	0	1	3	3
RH-I (%)	54	41	42	46	43
RH-II (%)	29	22	28	27	19
Wind Speed (kmph)	13	18	13	11	17
Wind Direction(deg.)	88	22	31	126	118



Agro meteorological advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry, cold and Mainly clear to Partly cloudy weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 22-26°C and 9-10°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 41-54% & 19-29, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be E, NE, SE and 11-18 km/h, respectively. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Garlic	Girthing	Thrips and aphid  leaf blight (yellowing)	<p>For thrips & aphid control spray neem seed extract 5% and make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml with imidacloprid 17.8 % EC 5 ml in 10 liters of water.</p> <p>To prevent leaf blight (yellowing) disease infestation</p>

			<p>in garlic spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure.</p> <p>Do not over irrigate to prevent leaf yellowing disease.</p>
Chick pea	Flowering/pod formation	<p>Monitor the crop regularly for prodenia and heliothes infestation</p> 	<p>Under present weather condition heliothes infestation observed in chickpea. So, Monitor the crop regularly for heliothes and prodenia infestation.</p> <ul style="list-style-type: none"> • “T” shaped bird perches in and around crop field to be installed for natural controlling the insect population. • Spray neem seed extract 5 % or neem seed oil 50 ml in 10 liters of water. • If infestation cross ETL spray 3 ml Spinosad in 10 liters of water. • Also to prevent blight disease infestation spray Mencozeb 27 gm in 10 liters of water at 10 days interval as precautionary measure. • Irrigate gram crop at flowering stage.
Cumin	Growth/Flowering	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm or imidacloprid 17.8 % 5 ml/10 liters of water.</p> <p>Under present weather condition spray tebuconazole 10 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
Coriander	Flowering	Thrips and Jassids	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 litres of water.</p>

			Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.
Wheat	Flowering	-	Apply irrigation at critical stage without stress.
Cotton	Picking & pulling of cotton stalk	<p>Heavy aatack of pink boll worm observed in cotton field. Carry out picking in time.</p> <p>Animals like goats, sheep among others, should be allowed to graze on unopened bolls and plant debris in the cotton fields after the last picking to prevent infestation of pink boll worm in subsequent year. After grazing don't burn cotton stalksbut, recycle the stalk by the doin situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator or make compost from the cotton residues to improve soil biomass and health.</p>	
Castor	Flowering/ capsule formation	<p>Jassids& white fly→</p> <p>Pod borer</p>	<p>Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water.</p> <p>For the control of pod borer spray chlorantraniliprole18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p>
Pigeon pea	Flowering/pod formation/dev elopment	Pod fly	<ul style="list-style-type: none"> • Spray neem seed extract 5 % or neem seed oil 50 mlin 10 litres of water. • If infestation cross ETL spray 20 ml Quinalphos or 5 ml Dichlorvos in 10 litres of water.
Onion	TP/Growth	<p>Thrips</p> <p>Leaf blight→</p> 	<p>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</p> <p>Make alternate sprayof Mencozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.</p>
Brinjal& Tomato	Flowering/ fruiting	Sucking pest →	Monitor the crop for infestation of jassid& white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with

		<p>Shoot & Fruit borer</p>  <p>Mites</p>	<p>diafenthiuron 50 %wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p>  <p>Anthracnose →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray Ethion 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p>
Lemon	Flowering and fruit formation	<p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p>	<p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water. Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p>
Mango	Flowering	<p>Blossom midge and Leaf hopper →</p>  <p>Powdery Mildew →</p>	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p> <p>Spray 15 gm carbendazim in 10 liters of water to prevent powdery mildew disease in mango.</p>

			
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out Brucella and other infectious & reproductive diseases evaluation. Spray phenyl in the animal shed to avoid flies and mosquitoes.</p> <p>Keep young calves indoors during night time to avoid cold stress. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Provide ad lib warm water to lactating animals. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.