



ગ્રામીણ કૃષિ મૌસમ સેવા

હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન
ઉત્તર સૌરાષ્ટ્રખેત આબોહવાકીય વિભાગ



તા.૨૬-૦૧-૨૦૧૬ થી તા.૩૦-૦૧-૨૦૧૬

(ગ્રામીણ કૃષિ મૌસમસેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જુ.કૃ.યુ.,તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિ હવામાન બુલેટીન નં. ૮૭/૨૦૧૬

પાછલા અઠવાડીયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી.)	0.0
મહત્તમ તાપમાન (સે.ગ્રે)	26.8-29.0
લઘુત્તમ તાપમાન (સે.ગ્રે)	8.1-11.0
ભેજ(%) સવાર	47-76
ભેજ(%) બપોરબાદ	30-39
પવનની ગતિ (કિમી/કલાક)	3.58-5.77

હવામાન આગાહી: પત્રક-ક

ઉત્તર સૌરાષ્ટ્રખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.૨૬-૦૧-૨૦૧૬ થી તા.૩૦-૦૧-૨૦૧૬

કૃષિ સલાહ:

સામાન્ય	<p>ભારતસરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા વિભાગ,જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના</p> <p>રાજકોટ: જીલ્લામાં આગામી તા.૨૬-૦૧-૨૦૧૬ થી તા.૩૦-૦૧-૨૦૧૬ દરમ્યાન હવામાન સુકું અને ચોખ્ખું રહેશે. પવનની દિશા નૈઋત્ય, પશ્ચિમ,દક્ષિણ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૧૦ થી ૧૪ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૦-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૧૨-૧૬ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</p>																														
	<table border="1"><thead><tr><th>તારીખ</th><th>દિવસ-૧</th><th>દિવસ-૨</th><th>દિવસ-૩</th><th>દિવસ-૪</th><th>દિવસ-૫</th></tr></thead><tbody><tr><td>પેરામીટર</td><td>૨૬-૦૧-૧૬</td><td>૨૭-૧૨-૧૬</td><td>૨૮-૦૧-૧૬</td><td>૨૯-૦૧-૧૬</td><td>૩૦-૦૧-૧૬</td></tr><tr><td>વરસાદ(મી.મી)</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>મહત્તમ તાપમાન (સે.ગ્રે)</td><td>30</td><td>30</td><td>30</td><td>31</td><td>32</td></tr><tr><td>લઘુત્તમ તાપમાન (સે.ગ્રે)</td><td>12</td><td>13</td><td>14</td><td>16</td><td>16</td></tr></tbody></table>	તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫	પેરામીટર	૨૬-૦૧-૧૬	૨૭-૧૨-૧૬	૨૮-૦૧-૧૬	૨૯-૦૧-૧૬	૩૦-૦૧-૧૬	વરસાદ(મી.મી)	0	0	0	0	0	મહત્તમ તાપમાન (સે.ગ્રે)	30	30	30	31	32	લઘુત્તમ તાપમાન (સે.ગ્રે)	12	13	14	16	16
તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫																										
પેરામીટર	૨૬-૦૧-૧૬	૨૭-૧૨-૧૬	૨૮-૦૧-૧૬	૨૯-૦૧-૧૬	૩૦-૦૧-૧૬																										
વરસાદ(મી.મી)	0	0	0	0	0																										
મહત્તમ તાપમાન (સે.ગ્રે)	30	30	30	31	32																										
લઘુત્તમ તાપમાન (સે.ગ્રે)	12	13	14	16	16																										

વાદળની સ્થિતિ(ઓકટા)	0	0	0	0	0
મહત્તમ ભેજ (%) સવાર	34	71	71	56	85
લઘુત્તમ ભેજ (%) બપોરબાદ	16	20	35	32	32
પવનની ગતિ (કિમી/કલાક)	12	12	13	10	14
પવનની દિશા (ડીગ્રી)	250	240	175	281	311

જામનગર: જુલલામાં આગામી તા.૨૬-૦૧-૨૦૧૬ થી તા.૩૦-૦૧-૨૦૧૬ દરમિયાન હવામાન સુકું અને યોખ્મું રહેશે. પવનની દિશા વાયવ્યન રહેવાની અને પવનની ઝડપ ૧૦ થી ૧૪ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૨૯-૩૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૧૫ - ૧૯ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૨૬-૦૧-૧૬	૨૭-૧૨-૧૬	૨૮-૦૧-૧૬	૨૯-૦૧-૧૬	૩૦-૦૧-૧૬
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમ તાપમાન (સે.ગ્રે)	29	30	30	31	31
લઘુત્તમ તાપમાન (સે.ગ્રે)	15	16	17	18	19
વાદળની સ્થિતિ(ઓકટા)	0	0	0	0	0
મહત્તમ ભેજ (%) સવાર	48	81	80	65	89
લઘુત્તમ ભેજ (%) બપોરબાદ	20	32	45	40	43
પવનની ગતિ (કિમી/કલાક)	12	13	14	10	14
પવનની દિશા (ડીગ્રી)	334	325	313	326	301




અમરેલી: જુલલામાં આગામી તા.૨૬-૦૧-૨૦૧૬ થી તા.૩૦-૦૧-૨૦૧૬ દરમિયાન હવામાન સુકું અને યોખ્મું રહેશે. પવનની દિશા નૈઋત્ય, દક્ષિણ પૂર્વ, અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૧૦ થી ૧૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૨-૩૩ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૧૦-૧૩ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.




તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૨૬-૦૧-૧૬	૨૭-૧૨-૧૬	૨૮-૦૧-૧૬	૨૯-૦૧-૧૬	૩૦-૦૧-૧૬
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમ તાપમાન (સે.ગ્રે)	32	32	33	33	33
લઘુત્તમ તાપમાન (સે.ગ્રે)	10	11	12	13	13
વાદળની સ્થિતિ(ઓકટા)	0	0	0	0	0
મહત્તમ ભેજ (%) સવાર	35	58	66	58	75
લઘુત્તમ ભેજ (%) બપોરબાદ	19	21	33	33	32
પવનની ગતિ (કિમી/કલાક)	12	12	13	10	15
પવનની દિશા (ડીગ્રી)	203	193	71	166	321

સુરેન્દ્રનગર: જીલ્લામાં આગામી તા.૨૬-૦૧-૨૦૧૬ થી તા.૩૦-૦૧-૨૦૧૬ દરમ્યાન હવામાન સુકું અને ચોખ્ખું રહેશે. પવનની દિશા ઉત્તર, નૈઋત્ય, પશ્ચિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૭ થી ૧૩ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૦-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૧૨-૧૬ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.




તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૨૬-૦૧-૧૬	૨૭-૧૨-૧૬	૨૮-૦૧-૧૬	૨૯-૦૧-૧૬	૩૦-૦૧-૧૬
વરસાદ(મી.મી)	૦	૦	૦	૦	૦
મહત્તમ તાપમાન (સે.ગ્રે)	૩૦	૩૦	૩૦	૩૧	૩૨
લઘુત્તમ તાપમાન (સે.ગ્રે)	૧૨	૧૩	૧૪	૧૬	૧૬
વાદળની સ્થિતિ(ઓકટા)	૦	૦	૦	૦	૦
મહત્તમ ભેજ (%) સવાર	૨૭	૨૭	૨૬	૨૬	૨૬
લઘુત્તમ ભેજ (%) બપોરબાદ	૧૩	૧૬	૩૨	૨૬	૨૪
પવનની ગતિ (કિમી/કલાક)	૧૧	૧૧	૧૧	૭	૧૩
પવનની દિશા (ડીગ્રી)	૩૪૨	૩૨૨	૨૭૪	૨૧૧	૨૯૭

- જાન્યુઆરી ૨૭ સુધી સુકું અને ચોખ્ખું હવામાં રહેશે. તેમજ મહત્તમ અને લઘુત્તમ તાપમાન અનુક્રમે ૨૮-૩૧ અને ૧૨-૧૫ ડીગ્રી સે. રહેશે અને તા.૨૭.૧.૧૬ના રોજ આકળ આવવાની શક્યતા હોવાથી આ દિવસે જીરુંના પાકને પિયત આપવું નહિ.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ
લસણ /ડુંગળી	વૃદ્ધિ	શ્રીપ્સ, પાનની ટોચનો સુકારો  પાનની ટોચનો સુકારો અને જાંબલી ધાબાનો રોગ 	(૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી) શ્રીપ્સના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને એમીદાક્લોપ્રીદ ૩૦.૫ ટકા ૪ મિલી અથવા પ્રોફેનોફોસ ૩૦ મિલી અથવા લેમ્બડા સાયહેલોથ્રીન ૧૫ મિલી દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. લસણ/ડુંગળીના પાકમાં પાનની ટોચનો સુકારો અને જાંબલી ધાબાનો રોગના નિયંત્રણ માટે ડાયથેન એમ ૪૫, ૨૭ ગ્રામ અથવા થયોફેનેટ મિથાયલ ૧૨ ગ્રામ અને હેક્ઝાકોનાઝોલ ૧૫ મિલી દવા વારાફરતી ૧૦ લીટર પાણીમાં મિશ્ર કરીને ૧૫ દિવસના અંતરે છંટકાવ કરવો.  પિયત નિયંત્રિત આપવું.

ચણા	ફૂલ-પોપટા બંધાવા	લીલી અને લશ્કરી ઈંચળ	ચણાના પાકમાં લીલી અને લશ્કરી ઈંચળના નિયંત્રણ માટે પક્ષીઓને બેસવા માટે 'T' આકારના બેલાખડા મુકવા. જો વધુ ઉપદ્રવ જણાય તો કવીનાલફોસ ૨૦ મિલી દવા ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
ઘઉં	ફૂલ /દાણા ભરાવા	-	ફૂલ અને દાણા ભરાવાની અવસ્થાએ પાણીની ખેંચ પડવા દેવી નહિ અને સમયસર પિયત આપવું.
જીરું	ફૂલ/દાણા બંધાવા	થ્રીપ્સ ચરમી અને રાખોડી  સુકારો 	થ્રીપ્સના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો. તેમજ ચરમીનો ઉપદ્રવ ન થાય તે માટે ડાયથેન એમ ૪૫, ૨૭ ગ્રામ અને હેક્ઝાકોનાઝોલ ૧૫ મિલી દવા વારાફરતી ૧૦ લીટર પાણીમાં મિશ્ર કરીને ૧૫ દિવસના અન્તરે છંટકાવ કરવો.  ઉભા પાકમાં સુકારાના નિયંત્રણ માટે પિયત આપવાનું બંધ કરવું.
ધાણા	ફૂલ/દાણા બંધાવા	થ્રીપ્સ અને સફેદ માખી	થ્રીપ્સ અને સફેદ માખીના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.

કપાસ	કપાસની વીણી	ગુલાબી ઈયળ 	કપાસની ગુલાબી ઈયળના નિયંત્રણ માટે જે કપાસના પાકમાં ગુલાબી ઈયળનો ઉપદ્રવ હોય ત્યાં બીજો ફાલ લેવો નહિ અને કપાસની વીણી કરીને કપાસ કાઢીને સળગાવી દેવો. 
રીંગણી અને ટમેટી	કુલ/ફળ	સફેદ માખી ડુંખ અને ફળ કોરી ખાનાર ઈયળ → કથીરી →	<ul style="list-style-type: none"> ▶ લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ▶ ફળ અને ડોકાની ઈયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને રીનાક્ષીપાયર દવા ૩ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ▶ રીંગણી અને ટમેટીના પાકમાં પાન કથીરીના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
કોબી અને ફલાવર	વૃદ્ધિ/દડા	હિરાફૂદ	કોબી અને ફલાવરમાં હિરાફૂદાણા નિયંત્રણ માટે સ્પીનોસાઇ ૩ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો
મરચી	કુલ/ફળ	પાનની કુકડ-થ્રીપ્સ →  કાલવ્રણ → 	પ્રોફેનોફોસ ૨૦ મી.લી. અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. કાલવ્રણ રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.

આંબો	ફૂલ	મધીયો અને મગીયાની ઈયળ	<p>આંબામાં મધીયો અને મગીયાની ઈયળના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૫ મિલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>  <p>આંબામાં ભુકીછારો/રાખોડીના નિયંત્રણ માટે હેક્ઝાકોનાઝોલ ૧૫ મિલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> 
<p>પશુ/મરઘા</p> 	<p>નાના વાછરડાઓને રાત્રે ઠંડીથી બચવા પશુ ઘરમાં રખવા. પ્રજીવથી થતા બૃસેલ્લા અને અન્ય રોગોની ચકાસણી કરાવવી. ખરવા મોવાસાના રોગનું રસીકરણ કરાવવું. ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મી.લી. ૧ લીટર પાણીમાં નાખીને છાંટવી. વિટામીન એ તથા ડી પુરકો આપવા. કરમિયાની દવા આપવી તથા ઉછરતા પશુઓમાં બૃસેલ્લા કાફવુંડનું રસીકરણ કરાવવું.</p>		



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-87 for the Rajkot district
 (Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
 (Period 26th to 30th January - 2016)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	0.0
Maximum temperature (°C)	26.8-29.0
Minimum temperature (°C)	8.1-11.0
Relative Humidity (%) I	47-76
Relative Humidity (%) II	30-39
Wind speed (Kmph)	3.58-5.77







Weather forecast until 08-30 hrs of 30/01/2016






(1) RAJKOT DISTRICT:






	26/01/16	27/01/16	28/01/16	29/01/16	30/01/16
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	30	30	30	31	32
Min.Temp.(°C)	12	13	14	16	16
Total CC (octa)	0	0	0	0	0
RH-I (%)	34	71	71	56	85
RH-II (%)	16	20	35	32	32
Wind Speed (kmph)	12	12	13	10	14
Wind Direction(deg.)	250	240	175	281	311

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry and clear weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 30-32°C and 12-16°C, respectively. • Relative Humidity: The RH is likely to be in the range of 16-85 % • Wind Direction and Speed: Direction and speed of wind is likely to be SW,W,S,NW and 10-14 km/h, respectively. <p style="color: red; text-align: center;">The fog would occur on 27-1-2016, so do not irrigate the cumin crop during the sais day.</p>		
Crops	Stage	Insect/Pest/Variety	Advisory

<p>Garlic</p>	<p>Girthing</p>	<p>Thrips</p>  <p>Alternaria blight/Purple Blotch</p> 	<p>In present weather conditions attack of thrips has been reported. Make alternative sprays of imidacloprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval.</p> <p>The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop.</p> <p>Make two spray of 27 gm Mencozeb or thiophanae methyl 12 gm/10 liters of water at 10 days interval.</p>
<p>Chick pea</p>	<p>Pod formation</p>	<p>Heliiothis, prodenia</p>	<p>“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation is cross ETL spray 20 ml Quinalphos in 10 litres of water.</p>
<p>Wheat</p>	<p>Flowering/grain filling</p>	<p>-</p>	<p>Apply irrigation at critical stages without stress.</p>
<p>Cumin</p>	<p>Grain formation/Pod development</p>	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather (fog) condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 

Coriander	Flowering/grain formation	Thrips, White fly	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 liters of water. Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent powdery mildew infestation.
Cotton	Picking/uprooting	Pink boll worm  	Under present weather condition heavy attack of pink boll worm observed in irrigated cotton, therefore farmers are advised not to take second flush of flowering and uproot the crop after picking. 
Onion	Growth	Thrips Purple blotch → 	In present weather conditions attack of thrips has been reported. Make alternative sprays of imidacloprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval. Spray Mencozeb 25 gm and hexaconazole 15 ml in 10 liters of water at 10 days interval.
Brinjal & Tomato	Flowering/fruitletting	Sucking pest → & Fruit borer 	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
Cabbage and cauliflower	Growth & fruitletting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.

<p>Chili</p>	<p>Flowering & fruiting</p>	<p>Thrips, →  Wilting → Anthracnose → </p>	<p>In present weather conditions severe attack of thrips has been reported. Spray 3 ml spinosad in 10 litres of water.</p> <p>Drenching of Trichoderma viridi 4 gm in 1 liter of water.</p> <p>Spray Mencozeb 27 gm in 10 liters of water</p>
<p>Mango</p>	<p>Flowering</p>	<p>Blossom midge and Leaf hopper →  Powdery Mildew→ </p>	<p>To prevent damage spray imidacloprid 5 ml/10 litres of water.</p> <p>Spray 15 ml Hexaconzole in 10 liters of water.</p>
<p>Livestock & poultry </p>	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



**Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523**












Agro met Advisory Service bulletin No.-87 for the Jamnagar district
(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
(Period 26rd to 30th January- 2016)
Weather forecast until 08-30 hrs of 30/01/2016







(2) JAMNAGAR DISTRICT:


	26/01/16	27/01/16	28/01/16	29/01/16	30/01/16
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	29	30	30	31	31
Min.Temp.(°C)	15	16	17	18	19
Total CC (octa)	0	0	0	0	0
RH-I (%)	48	81	80	65	89
RH-II (%)	20	32	45	40	43
Wind Speed (kmph)	12	13	14	10	14
Wind Direction(deg.)	334	325	313	326	301

Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry and clear weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 29-31°C and 15-19 °c, respectively. • Relative Humidity: The RH is likely to be in the range of 20-89 % • Wind Direction and Speed: Direction and speed of wind is likely to be NW and 10-14 km/h, respectively. <p>The fog would occur on 27-1-2016, so do not irrigate the cumin crop during the sais day.</p>		
Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Girthing	Thrips 	In present weather conditions attack of thrips has been reported. Make alternative sprays of imidacloprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval.
		Alternaria blight/Purple Blotch 	The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop. Make two spray of 27 gm Mencozeb or thiophanae methyl 12 gm/10 liters of water at 10 days interval.
Chick pea	Pod formation	Heliiothis, prodenia	“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation is cross ETL spray 20 ml Quinalphos in 10 litres of water.
Wheat	Flowering/grain filling	-	Apply irrigation at critical stages without stress.

<p>Cumin</p>	<p>Grain formation</p>	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather (fog) condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
<p>Coriander</p>	<p>Flowering/grain formation</p>	<p>Thrips, White fly</p>	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent powdery mildew infestation.</p>
<p>Cotton</p>	<p>Picking/uprooting</p>	<p>Pink boll worm</p>  	<p>Under present weather condition heavy attack of pink boll worm observed in irrigated cotton, therefore farmers are advice not to take second flush of flowering and uproot the crop after picking.</p> 
<p>Onion</p>	<p>Growth</p>	<p>Thrips</p> <p>Purple blotch →</p>	<p>In present weather conditions attack of thrips has been reported. Make alternative sprays of imidaclicprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval.</p>

			Spray Mencozeb 25 gm and hexaconazole 15 ml in 10 liters of water at 10 days interval.
Brinjal & Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
Cabbage and cauliflower	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.
Chili	Flowering & fruiting	Thrips, →  Wilting → Anthracnose → 	In present weather conditions severe attack of thrips has been reported. Spray 3 ml spinosad in 10 litres of water. Drenching of Trichoderma viridi 4 gm in 1 liter of water. Spray Mencozeb 27 gm in 10 liters of water
Mango	Flowering	Blossom midge and Leaf hopper →  Powdery Mildew → 	To prevent damage spray imidacloprid 5 ml/10 litres of water. Spray 15 ml Hexaconzole in 10 liters of water.

Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.

	<p>Gramin Krushi Mausam Seva Main Dry Farming Research Station Junagadh Agricultural University Targhadia (Rajkot) - 360 003 Phone: (0281)2784260(O), 2331686(R) Fax : (0281)2784722, Cell :94277 28523</p>	
---	---	---

Agro met Advisory Service bulletin No.87 for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
 (Period 26rd to 30th January - 2016)






Weather forecast until 08-30 hrs of 30/01/2016







(3) AMRELI DISTRICT:






	26/01/16	27/01/16	28/01/16	29/01/16	30/01/16
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	32	32	33	33	33
Min.Temp.(°C)	10	11	12	13	13
Total CC (octa)	0	0	0	0	0
RH-I (%)	35	58	66	58	75
RH-II (%)	19	21	33	33	32
Wind Speed (kmph)	12	12	13	10	15
Wind Direction(deg.)	203	193	71	166	321

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry and clear weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 32-33°C and 10-13°C, respectively. • Relative Humidity: The RH is likely to be in the range of 19-75 % • Wind Direction and Speed: Direction and speed of wind is likely to be,SW,S,E,NW and 10-15 km/h, respectively. • The fog would occur on 27-1-2016, so do not irrigate the cumin crop during the sais day.
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Girthing	<p>Thrips</p>  <p>Alternaria blight/Purple Blotch</p> 	<p>In present weather conditions attack of thrips has been reported. Make alternative sprays of imidacloprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval.</p> <p>The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop. Make two spray of 27 gm Mencozeb or thiophanae methyl 12 gm/10 liters of water at 10 days interval.</p>
Chick pea	Pod formation	Heliiothis, prodenia	<p>“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation is cross ETL spray 20 ml Quinalphos in 10 litres of water.</p>
Wheat	Flowering/grain filling	-	Apply irrigation at critical stages without stress.
Cumin	Grain formation	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather (fog) condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>   <p>Stop irrigation to prevent further infection in standing crop.</p>

		<p>Wilt</p> 	
Coriander	Flowering/grain formation	Thrips, White fly	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent powdery mildew infestation.</p>
Cotton	Picking/uprooting	<p>Pink boll worm</p>  	<p>Under present weather condition heavy attack of pink boll worm observed in irrigated cotton, therefore farmers are advice not to take second flush of flowering and uproot the crop after picking.</p> 
Onion	Growth	<p>Thrips</p> <p>Purple blotch →</p> 	<p>In present weather conditions attack of thrips has been reported. Make alternative sprays of imidacloprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval.</p> <p>Spray Mencozeb 25 gm and hexaconazole 15 ml in 10 liters of water at 10 days interval.</p>
Brinjal & Tomato	Flowering/fruitletting	<p>Sucking pest → & Fruit borer</p> 	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.</p>
Cabbage and cauliflower	Growth & fruitletting	Diamond back moth	<p>If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and</p>

			cabbage.
Chili	Flowering & fruiting	Thrips, →  Wilting → Anthracnose → 	In present weather conditions severe attack of thrips has been reported. Spray 3 ml spinosad in 10 litres of water. Drenching of Trichoderma viridi 4 gm in 1 liter of water. Spray Mencozeb 27 gm in 10 liters of water
Mango	Flowering	Blossom midge and Leaf hopper →  Powdery Mildew → 	To prevent damage spray imidacloprid 5 ml/10 litres of water. Spray 15 ml Hexaconzole in 10 liters of water.
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:94277 28523



Agro met Advisory Service bulletin No.-87 for the Surendranagar district



(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
(Period 26rd to 30th January- 2016)










Weather forecast until 08-30 hrs of 30/01/2016






(4) SURENDRANAGAR DISTRICT:



	26/01/16	27/01/16	28/01/16	29/01/16	30/01/16
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	30	30	30	31	32
Min.Temp.(°C)	12	13	14	16	16
Total CC (octa)	0	0	0	0	0
RH-I (%)	27	57	66	47	78
RH-II (%)	13	16	32	26	24
Wind Speed (kmph)	11	11	11	7	13
Wind Direction(deg.)	342	322	274	211	297

Agro meteorological advisories:

General Advice	<ul style="list-style-type: none"> • Rain fall: Dry and clear weather forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range 30-32 and 12-16°C, respectively. • Relative Humidity: The RH is likely to be in the range of 13-78 % • Wind Direction and Speed: Direction and speed of wind is likely to be N,NW,W,SW, and 7-13 km/h, respectively. • The fog would occur on 27-1-2016, so do not irrigate the cumin crop during the sais day. 		
Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Girthing	Thrips 	In present weather conditions attack of thrips has been reported. Make alternative sprays of imidacloprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval.
		Alternaria blight/Purple Blotch 	The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop. Make two spray of 27 gm Mencozeb or thiophanae methyl 12 gm/10 liters of water at 10 days interval.

Chick pea	Pod formation	Heliiothis, prodenia	“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation is cross ETL spray 20 ml Quinalphos in 10 litres of water.
Wheat	Flowering/grain filling	-	Apply irrigation at critical stages without stress.
Cumin	Grain formation	<p>Aphid and thrips</p> <p>Blight and Powery mildew</p>  <p>Wilt</p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray Acetamiprid 5 gm/10 liters of water.</p> <p>Under present weather (fog) condition spray Hexaconazole 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
Coriander	Flowering/grain formation	Thrips, White fly	<p>To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and 15 gm Acepahte in 10 liters of water.</p> <p>Under present weather condition spray Hexaconazole 15 ml/10 liters of water to prevent powdery mildew infestation.</p>
Cotton	Picking/uprooting	<p>Pink boll worm</p>  	<p>Under present weather condition heavy attack of pink boll worm observed in irrigated cotton, therefore farmers are advice not to take second flush of flowering and uproot the crop after picking.</p>   

Onion	Growth	Thrips Purple blotch → 	In present weather conditions attack of thrips has been reported. Make alternative sprays of imidacloprid 30.5 SC 3 ml and profenofos 20 ml in 10 liters of water at 10 days interval. Spray Mencozeb 25 gm and hexaconazole 15 ml in 10 liters of water at 10 days interval.
Brinjal & Tomato	Flowering/ fruiting	Sucking pest → & Fruit borer 	To control sucking pest spray neem seed extract 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
Cabbage and cauliflower	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.
Chili	Flowering & fruiting	Thrips, →  Wilting → Anthraxnose → 	In present weather conditions severe attack of thrips has been reported. Spray 3 ml spinosad in 10 litres of water. Drenching of Trichoderma viridi 4 gm in 1 liter of water. Spray Mencozeb 27 gm in 10 liters of water
Mango	Flowering	Blossom midge and Leaf hopper →  Powdery	To prevent damage spray imidacloprid 5 ml/10 litres of water. Spray 15 ml Hexaconzole in 10 liters of water.

		<p>Mildew→</p> 	
<p>Livestock & poultry</p> 	<p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.