



ગ્રામીણ કૃષિ મૌસમ સેવા
હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ
તા.૨૭-૧૧-૨૦૧૯ થી તા.૦૧-૧૨-૨૦૧૯



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કૃ.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિ હવામાન બુલેટીન નં. ૬૯/૨૦૧૯

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

| હવામાન પરિબલો | |
|--------------------------|-----------|
| વરસાદ(મી.મી) | 0.0 |
| મહત્તમ તાપમાન (સે.ગ્રે) | 32.3-34.1 |
| લઘુત્તમ તાપમાન (સે.ગ્રે) | 17.0-20.0 |
| ભેજ(%) સવાર | 68-80 |
| ભેજ(%) બપોરબાદ | 40-50 |
| પવનની ગતિ (કિમી/કલાક) | 1.9-3.3 |

હવામાન આગાહી: પત્રક-૬

ઉત્તરસૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.૨૭-૧૧-૨૦૧૯ થી ૦૧-૧૨-૨૦૧૯

કૃષિ સલાહ:

| સામાન્ય | ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના પશ્ચિમ રાજકોટ: જીલ્લામાં આગામી તા.૨૭-૧૧-૨૦૧૯ થી તા.૦૧-૧૨-૨૦૧૯ દરમિયાન હવામાન સૂકું અને ચોખ્ખું રહેશે. પવનની દિશા ઈશાન અને ઉત્તરની રહેવાની અને પવનની ઝડપ ૧૩ થી ૧૭ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૨-૩૩ ડીગ્રી સેલ્શિયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૧૯-૨૦ ડીગ્રી સેલ્શિયસ જેટલું રહેવાની સંભાવના છે. | | | | | |
|---------|---|----------|----------|----------|----------|----------|
| | તારીખ | દિવસ-૧ | દિવસ-૨ | દિવસ-૩ | દિવસ-૪ | દિવસ-૫ |
| | પેરામીટર | ૨૭-૧૧-૧૯ | ૨૮-૧૧-૧૯ | ૨૯-૧૧-૧૯ | ૩૦-૧૧-૧૯ | ૦૧-૧૨-૧૯ |
| | વરસાદ(મી.મી) | 0 | 0 | 0 | 0 | 0 |
| | મહત્તમતાપમાન (સે.ગ્રે) | 33 | 32 | 32 | 32 | 33 |
| | લઘુત્તમતાપમાન (સે.ગ્રે) | 20 | 20 | 19 | 19 | 19 |
| | વાદળનીસ્થિતિ(ઓકટા) | 0 | 0 | 3 | 0 | 0 |
| | મહત્તમભેજ(%) સવાર | 55 | 56 | 55 | 56 | 53 |
| | લઘુત્તમભેજ (%) બપોરબાદ | 31 | 31 | 32 | 31 | 29 |

| | | | | | |
|---------------------|----|----|----|----|----|
| પવનનીગતિ(કિમી/કલાક) | 13 | 17 | 15 | 13 | 13 |
| પવનનીદિશા(ડીગ્રી) | 63 | 16 | 20 | 17 | 30 |

મોરબી : જીલ્લામાં આગામી તા.૨૭-૧૧-૨૦૧૯ થી તા.૦૧-૧૨-૨૦૧૯ દરમ્યાન હવામાન સુકું અને મુખત્વે વાદળછાયું રહશે. પવનની દિશા પૂર્વ અને ઈશાનની રહેવાની અને પવનની ઝડપ ૬ થી ૯ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૨-૩૩ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૧૯-૨૦ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

| તારીખ | દિવસ-૧ | દિવસ-૨ | દિવસ-૩ | દિવસ-૪ | દિવસ-૫ |
|-------------------------|----------|----------|----------|----------|----------|
| પેરામીટર | ૨૭-૧૧-૧૯ | ૨૮-૧૧-૧૯ | ૨૯-૧૧-૧૯ | ૩૦-૧૧-૧૯ | ૦૧-૧૨-૧૯ |
| વરસાદ(મી.મી) | 0 | 0 | 0 | 0 | 0 |
| મહત્તમતાપમાન (સે.ગ્રે) | 33 | 32 | 32 | 32 | 33 |
| લઘુત્તમતાપમાન (સે.ગ્રે) | 20 | 20 | 19 | 19 | 19 |
| વાદળનીસ્થિતિ(ઓકટા) | 0 | 1 | 3 | 1 | 1 |
| મહત્તમભેજ(%) સવાર | 54 | 54 | 54 | 52 | 50 |
| લઘુત્તમભેજ (%) બપોરબાદ | 32 | 35 | 30 | 28 | 27 |
| પવનનીગતિ(કિમી/કલાક) | 6 | 6 | 5 | 9 | 6 |
| પવનનીદિશા(ડીગ્રી) | 105 | 62 | 68 | 45 | 96 |

જામનગર: જીલ્લામાં આગામી તા.૨૭-૧૧-૨૦૧૯ થી તા.૦૧-૧૨-૨૦૧૯ દરમ્યાન હવામાન સુકું અને યોખ્ખું રહશે. પવનની દિશા ઈશાન અને ઉતરની રહેવાની અને પવનની ઝડપ ૧૯ થી ૨૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૧-૩૩ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૦-૨૧ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

| તારીખ | દિવસ-૧ | દિવસ-૨ | દિવસ-૩ | દિવસ-૪ | દિવસ-૫ |
|-------------------------|----------|----------|----------|----------|----------|
| પેરામીટર | ૨૭-૧૧-૧૯ | ૨૮-૧૧-૧૯ | ૨૯-૧૧-૧૯ | ૩૦-૧૧-૧૯ | ૦૧-૧૨-૧૯ |
| વરસાદ(મી.મી) | 0 | 0 | 0 | 0 | 0 |
| મહત્તમતાપમાન (સે.ગ્રે) | 33 | 31 | 31 | 31 | 31 |
| લઘુત્તમતાપમાન (સે.ગ્રે) | 21 | 21 | 21 | 20 | 20 |
| વાદળનીસ્થિતિ(ઓકટા) | 0 | 0 | 3 | 0 | 0 |
| મહત્તમભેજ(%) સવાર | 58 | 57 | 61 | 62 | 62 |
| લઘુત્તમભેજ (%) બપોરબાદ | 41 | 40 | 43 | 42 | 41 |
| પવનનીગતિ(કિમી/કલાક) | 22 | 25 | 24 | 21 | 19 |
| પવનનીદિશા(ડીગ્રી) | 25 | 24 | 29 | 24 | 16 |

દેવભૂમિ દ્વારકા: જીલ્લામાં આગામી તા.૨૭-૧૧-૨૦૧૯ થી તા.૦૧-૧૨-૨૦૧૯ દરમ્યાન હવામાન સુકું અને મુખત્વે વાદળછાયું રહશે. પવનની દિશા ઈશાન અને ઉતરની રહેવાની અને પવનની ઝડપ ૧૯ થી ૨૪ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૨૯-૩૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમ્યાન ૨૧-૨૨ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

| તારીખ | દિવસ-૧ | દિવસ-૨ | દિવસ-૩ | દિવસ-૪ | દિવસ-૫ |
|-------------------------|----------|----------|----------|----------|----------|
| પેરામીટર | ૨૭-૧૧-૧૯ | ૨૮-૧૧-૧૯ | ૨૯-૧૧-૧૯ | ૩૦-૧૧-૧૯ | ૦૧-૧૨-૧૯ |
| વરસાદ(મી.મી) | ૦ | ૦ | ૦ | ૦ | ૦ |
| મહત્તમતાપમાન (સે.ગ્રે) | ૩૧ | ૨૯ | ૨૯ | ૨૯ | ૨૯ |
| લઘુત્તમતાપમાન (સે.ગ્રે) | ૨૨ | ૨૧ | ૨૨ | ૨૨ | ૨૨ |
| વાદળનીસ્થિતિ(ઓકટા) | ૦ | ૧ | ૩ | ૦ | ૦ |
| મહત્તમભેજ(%) સવાર | ૬૧ | ૫૭ | ૬૨ | ૬૪ | ૬૩ |
| લઘુત્તમભેજ (%) બપોરબાદ | ૩૯ | ૩૯ | ૪૨ | ૪૨ | ૪૧ |
| પવનનીગતિ(કિમી/કલાક) | ૨૨ | ૨૪ | ૨૪ | ૨૦ | ૧૯ |
| પવનનીદિશા(ડીગ્રી) | ૨૭ | ૨૬ | ૩૩ | ૨૮ | ૧૮ |

અમરેલી: જીલ્લામાં આગામી તા.૨૭-૧૧-૨૦૧૯ થી તા.૦૧-૧૨-૨૦૧૯ દરમિયાન હવામાન સુકું અને મુખત્વે વાદળછાયું રહેશે. પવનની દિશા દક્ષિણ, પૂર્વ, ઉત્તર અને ઈશાનની રહેવાની અને પવનની ઝડપ ૧૦ થી ૧૫ કીમી/કલાક રહેની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૨-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૦-૨૧ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

| તારીખ | દિવસ-૧ | દિવસ-૨ | દિવસ-૩ | દિવસ-૪ | દિવસ-૫ |
|-------------------------|----------|----------|----------|----------|----------|
| પેરામીટર | ૨૭-૧૧-૧૯ | ૨૮-૧૧-૧૯ | ૨૯-૧૧-૧૯ | ૩૦-૧૧-૧૯ | ૦૧-૧૨-૧૯ |
| વરસાદ(મી.મી) | ૦ | ૦ | ૦ | ૦ | ૦ |
| મહત્તમતાપમાન (સે.ગ્રે) | ૩૨ | ૩૨ | ૩૨ | ૩૨ | ૩૨ |
| લઘુત્તમતાપમાન (સે.ગ્રે) | ૨૧ | ૨૧ | ૨૧ | ૨૧ | ૨૦ |
| વાદળનીસ્થિતિ(ઓકટા) | ૧ | ૦ | ૪ | ૧ | ૧ |
| મહત્તમભેજ(%) સવાર | ૫૮ | ૫૮ | ૫૬ | ૫૪ | ૫૩ |
| લઘુત્તમભેજ(%) બપોરબાદ | ૩૬ | ૩૫ | ૩૫ | ૩૩ | ૩૩ |
| પવનનીગતિ(કિમી/કલાક) | ૧૦ | ૧૫ | ૧૫ | ૧૪ | ૧૪ |
| પવનનીદિશા(ડીગ્રી) | ૨૦૧ | ૭૬ | ૧૫ | ૬૩ | ૯૭ |






સુરેન્દ્રનગર: જીલ્લામાં આગામી તા.૨૭-૧૧-૨૦૧૯ થી તા.૦૧-૧૨-૨૦૧૯ દરમિયાન હવામાન સુકું અને યોખ્ખું રહેશે. પવનની દિશા ઈશાન અને અગ્નિની રહેવાની અને પવનની ઝડપ ૧૨ થી ૧૭ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૨-૩૩ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૧૯-૨૦ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.


| તારીખ | દિવસ-૧ | દિવસ-૨ | દિવસ-૩ | દિવસ-૪ | દિવસ-૫ |
|-------------------------|----------|----------|----------|----------|----------|
| પેરામીટર | ૨૭-૧૧-૧૯ | ૨૮-૧૧-૧૯ | ૨૯-૧૧-૧૯ | ૩૦-૧૧-૧૯ | ૦૧-૧૨-૧૯ |
| વરસાદ(મી.મી) | ૦ | ૦ | ૦ | ૦ | ૦ |
| મહત્તમતાપમાન (સે.ગ્રે) | ૩૩ | ૩૨ | ૩૨ | ૩૨ | ૩૩ |
| લઘુત્તમતાપમાન (સે.ગ્રે) | ૨૦ | ૨૦ | ૧૯ | ૧૯ | ૧૯ |
| વાદળનીસ્થિતિ(ઓકટા) | ૦ | ૦ | ૩ | ૦ | ૦ |

| | | | | | |
|---|--------------|-----------------------------------|--|----|-----|
| મહત્તમભેજ(%) સવાર | 55 | 56 | 55 | 54 | 52 |
| લઘુત્તમભેજ(%) બપોરબાદ | 33 | 35 | 35 | 35 | 34 |
| પવનનીગતિ(કિમી/કલાક) | 14 | 17 | 15 | 12 | 13 |
| પવનનીદિશા(ડીગ્રી) | 62 | 47 | 40 | 33 | 114 |
| <p>➤ ૨૭-નવેમ્બર થી ૦૧-ડિસેમ્બર ૨૦૧૯ હવામાન હુંફાળું, ભેજવાળું અને મોટાભાગે ચોખ્ખું રહશે. પવનની દિશા ઈશાન અને ઉતરની મોટાભાગે રહેવાની અને પવનની ઝડપ ૬ થી ૨૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૨૯-૩૩ ડીગ્રી સેલ્સીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૧૯-૨૧ ડીગ્રી સેલ્સીયસ જેટલું રહેવાની સંભાવના છે.</p> <ul style="list-style-type: none"> હાલ ચોમાસું પાકોની કાપણી થય ગયેલ હોય તો તે જમીનમાં શિયાળુ પાકો જેવા કે ચણા, ઘઉં, જીરું, ધાણાના પાકનું પસંદગી મુજબ વાવેતર કરવું. હાલ પિયતની સગવડવાળા વિસ્તારમાં શિયાળુ પાકોનું વાવેતર ચાલુ હોય ખેડૂતભાઈઓને ખાસ જણાવવાનું કે દરેક પાકના બીજને થાયરમ અથવા મેન્કોઝેબ અથવા કબોન્ડાઝીમ માંથી કોઈ પણ એક દવાનો ૨-૩ ગ્રામ દવા પ્રતિ ૧ કી.ગ્રા. બીજ મુજબ પટ આપવો. તેમજ અસરકારક નિદામણ નિયંત્રણ માટે ચણા, જીરું અને ધાણાના પાકમાં, પાક અને નીંદણના સ્કુરણ પહેલા પેન્ડીમેથાલીન (સ્ટોમ્પ ૩૦ ઈ.સી.) ૬૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી , ઘઉંના પાકમાં પેન્ડીમેથાલીન (સ્ટોમ્પ ૩૦ ઈ.સી.) ૩૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી, લસણ અને ડુંગળીના પાકમાં ઓક્સીફલુઓરફેન (૨૩.૫ ઈ.સી.) ૨૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી અને ઈસબગુલના પાકમાં આઈસોપ્રોટ્યુરોન (કનક ૫૦ ઈ.સી.) ૨૦ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી જેથી નિંદામણનો મજુરી ખર્ચ બચાવી શકાય અને પાક નિંદણ મુક્ત રાખી શકાય . | | | | | |
| મુખ્ય પાક | પાક અવસ્થા | રોગ/જીવાત/જાત | કૃષિ સલાહ (૧૦ લીટરપાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી) | | |
| લસણ | ઉગાવો/વૃદ્ધિ | - | જમીનની પરત મુજબ ૧૦-૧૨ દિવસે પિયત આપવું. | | |
| ચણા | વાવેતર/ઉગાવો | ગુજરાત ચણા-૨, ગુજરાત જુનાગઢ ચણા-૩ | પાયામાં ૨૦ કી.ગ્રા. નાઈટ્રોજન, ૪૦ કી.ગ્રા. ફોસ્ફોરસ, પ્રતિ હેક્ટર મુજબ આપી ચણાની વાવણી કરવી.ચણાના પાક માટે બીયારણનો જથ્થો ૬૦ થી ૭૫ કી.ગ્રા./ હે. રાખવો. વાવેતર સમયે ૨ થી ૨.૫ ગ્રામ /કી.ગ્રા. બીજને કેપ્ટન અથવા થાઈરમનો પટ આપવો અને પાયામા રાસાયણિક ખાતર ૮૭ કી.ગ્રા. ડી.એ.પી. અને ૧૦ કી.ગ્રા. યુરીયા/હે. આપવું. વાવેતર વખતે ચણાના બીજને ૨૫૦ ગ્રામ રાઈઝોબીયમ અને ફોસ્ફેટ કલ્ચરનો પ્રતિ ૧૦ કી.ગ્રા. મુજબ પટ આપવો | | |
| ધાણા | વાવેતર | ગુજરાત ધાણા-૨ | પાયામાં ૨૨ કી.ગ્રા. ડી.એ.પી. અને ૧૩ કી.ગ્રા. યુરીયા પ્રતિ હેક્ટર મુજબ આપી ધાણાની વાવણી કરવી. વાવેતર વખતે ધાણાના બીજને થાયરમ ૨ ગ્રામ પ્રતિ ૧ કી.ગ્રા. મુજબ પટ આપવો. | | |
| જીરું | વાવેતર | ગુજરાત જીરું-૪ | પાયામાં ૩૩ કી.ગ્રા. ડીએપી પ્રતિ હેક્ટર મુજબ આપી જીરૂની | | |

| | | | |
|--------|--------------|---|---|
| | | | વાવણી કરવી. વાવેતર વખતે જીરૂના બીજને થાયરમ ૨ ગ્રામ પ્રતિ ૧ કી.ગ્રા. મુજબ પટ આપવો. |
| ઘઉં | વાવેતર | લોક-૧, જી.ડબ્લ્યુ-૪૯૬ | ઘઉંના પાક માટે બીયારણનો જથ્થો ૧૦૦ થી ૧૨૫ કી.ગ્રા.પ્રતિ હેક્ટર રાખવો.પાયામાં ૧૩૦ કી.ગ્રા. ડીએપી, ૮૦ કી.ગ્રા. યુરિયા અને ૨૫ કી.ગ્રા. ઝીંક સલ્ફેટ પ્રતિ હેક્ટર મુજબ આપી ઘઉંની વાવણી કરવી. જ્યાં ઉધયનો ઉપદ્રવ રહેતો હોય ત્યાં ક્લોરપાયરીફોસ ૨૦ ટકા એ.સી. ૫ લીટર દવા પ્રથમ પિયત પાણી સાથે આપવી. |
| રાયડો | વાવેતર | રાય વરુણા, ગુજરાત -૩ અને ૪ | પાયામાં ૩૦ કી.ગ્રા. નાઈટ્રોજન, ૩૦ કી.ગ્રા. ફોસ્ફોરસ, પ્રતિ હેક્ટર મુજબ આપી રાયડાની વાવણી કરવી. |
| કપાસ | ફૂલ /જીન્ડવા | તડતડીયા અને સફેદ માખી પાનના ટપકા ગુલાબી ઈયળ | <ul style="list-style-type: none"> ➤ કપાસમાં તડતડીયા અને સફેદ માખીના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ડાયકેન્થાયુરોન ૫૦ ટકા વે.પા. ટકા ૧૬ ગ્રામ પ્રતિ ૧૦ લીટર પાણીમાં નાખીને છંટકાવ કરવો. ➤ કપાસમાં યુરીયા અને એમોનિયમ સલ્ફેટ અને મ્યુરેટ ઓફ પોટાશ દરેક ૩૦ કી.ગ્રા./હે. પુરતી ખાતર તરીકે આપવું. ➤ કપાસમાં સતત વરસાદી હવામાને કારણે પાન ટપકા રોગનો ઉપદ્રવ વધે નહિ તે માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા હેક્ઝાકોનાઝોલ ૧૫ મિલી પ્રતિ ૧૦ લીટર પાણીમાં નાખીને છંટકાવ કરવો. ➤ કપાસમાં ગુલાબી ઈયળના નર કુદાને એકત્રીત કરવા માટે હેક્ટરે ૪૦ ની સંખ્યામાં ફેરોમેન ટ્રેપ ગોઠવવાથી ઈયળોનું શરૂઆતની અવસ્થાથી જ નિયંત્રણ કરી શકાય. કપાસમાં ગુલાબી ઈયળથી નુકશાન પામેલ ફૂલ અને યાપવા તોડીને જમીનમાં દાટી દેવા. વધુ નુકશાન હોય તો પ્રોફેનોફોસ ૫૦% ઇ.સી. ૨૦ મી.લી. અથવા સાયપરમેથ્રીન ૧૦ મિલી બીવેરીયા બાસીઆના ફૂગનો પાઉડર ૪૦ ગ્રામ ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો. |
| દિવેલા | વૃદ્ધિ/કુલ | લીલી પોપટી, સફેદ માખી ઘોડિયા અને | <ul style="list-style-type: none"> ➤ સફેદ માખી અને લીલી પોપટીનો વધુ ઉપદ્રવ જોવા મળે છે. જેના નિયંત્રણ માટે એસીટામાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અને ઈમીડાકલોપ્રીડ ૩૦.૫ % ૩ મિલી સાથે લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ ઘોડિયા અને લશ્કરી ઈયળના નિયંત્રણ માટે ક્લોરોપાયરીફોસ |

| | | લશ્કરી | ૨૫ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી. |
|------------------------|---------------|---|--|
| ચોમાસું ડુંગળી | દડાનો વિકાસ | પાનનો સુકારો | <ul style="list-style-type: none"> ➤ ચોમાસુ ડુંગળીના પાકમાં પાનનાં સુકારાના નિયંત્રણ માટે મેન્કોઝેબ ૪૦ ગ્રામ અથવા હેક્ઝાકોનાઝોલ ૨૦ મિલી ૧ પમ્પમાં નાખીને ૧૦ દિવસના અંતરે ૩ છંટકાવ કરવા. ➤ નાઈટ્રોજન યુક્ત એમોનિયમ સલ્ફેટ ખાતરનો પુરતી ડોઝ ૩૦ કીલો ગ્રામ પ્રતિ હે. આપવો. |
| તુવેર | ફૂલ/શીંગ | - | ફૂલ અવસ્થાએ પિયત આપવું. |
| વટાણા ગાજર | વાવેતર | શાકભાજી માટે વટાણાની પુસા પ્રગતિ અને ગાજરની પુસા રુધીરા અને પુસા કેસર જાતો પસંદ કરવી. | |
| કોબી અને ફલાવર | વૃદ્ધિ/દડા | હિરાફૂદુ | કોબી અને ફલાવરમાં હિરાફૂદાના નિયંત્રણ માટે સ્પીનોસાઈ ૩ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. |
| ભીંડા | ફૂલ/ફળ | તડતડીયા અને સફેદ માખી શીંગ કોરી ખાનાર ઈંચળ પાન કથીરી | <ul style="list-style-type: none"> ➤ ભીંડામાં તડતડીયા અને સફેદ માખીના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ડાયકેન્ટાયુરોન ૫૦ ટકા વે.પા. ટકા ૧૬ ગ્રામ અથવા ટ્રાયઝોફોસ ૪૦ ઈસી ૨૫ મિ.લી. દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ તેના ફળ અને ડોકા કોરી ખાનાર ઈંચળનો ઉપદ્રવ વધુ હોય તો સ્પીનોસાઈ ૩ મિલી /૧૦ લીટર પાણીમાં ઓગાળી છંટકાવ કરવો. ➤ જો પાનકથીરીનો ઉપદ્રવ હોય તો ઇથીઓન ૩૦ મિલી પ્રતિ પંપ છંટકાવ કરવો. <p>ભીંડાની વીણી બાદ પ્રતિ એકરે ૫-૭ કી.ગ્રા. યુરિઆ ખાતર આપવું.</p> |
| રીંગણી અને ટમેટી | ફુલ/ફળ | તડતડીયા અને સફેદ માખી ડુંખ અને→ ફળ કોરી ખાનાર ઈંચળ પાન કથીરી | <ul style="list-style-type: none"> ➤ તડતડીયા અને સફેદ માખીના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦ મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ડાયકેન્ટાયુરોન ૫૦ ટકા વે.પા. ૧૬ ગ્રામ અથવા ટ્રાયઝોફોસ ૪૦ ઈસી ૨૫ મિ.લી. દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ ફળ અને ડોકાની ઈંચળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને ક્લોરોનટ્રાનીલીપ્રોલ ૧૮.૫ એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઈસી મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ જો પાનકથીરીનો ઉપદ્રવ હોય તો પ્રોપરગાઈટ ૫૭ ઈ.સી. ૧૫ મિલી પ્રતિ પંપ છંટકાવ કરવો. |
| મરચી | વૃદ્ધિ/ફુલ/ફળ | થ્રીપ્સ, સફેદ | ➤ પ્રોફેનોફોસ ૨૦ મીલી અને લીંબોળીનું તેલ ૫૦ મિલી અથવા |

| | | | |
|-------|------------|---|--|
| | | <p>માખી</p>  <p>કાલવણ →</p>  | <p>લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. થ્રીપ્સનો વધુ ઉપદ્રવ હોય તો સ્પીનોસાઈ અથવા સ્પીનેટોરમ દવા ૫ મિલી. પ્રતિ પંપ છાંટવી.</p> <p>➤ કાલવણ રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.</p> |
| લીંબુ | ફળનો વિકાસ | <p>પાન કથીરી</p>   <p>સાયલા અને થ્રીપ્સ</p> <p>બદામી ટપકાં</p> <p>પાનકોરીયુ હસ્ત બહાર</p> | <p>લીંબુના બગીચામાં પાન કથીરીનો ઉપદ્રવ વધુ હોય તો તેના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>તેના નિયંત્રણ માટે ઈમીડાક્લોપ્રીડ ૩૦.૫ ટકા ૫ મીલી અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. વધુ ઉપદ્રવ હોય તો ટ્રાયઝોફેસ ૪૦ ટકા ૨૦/૧૦ લિટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>  <p>બદામી ટપકાના નિયંત્રણ માટે કોપર ઓક્સીક્લોરાઈડ ૪૦ ગ્રામ અને ૧.૫ ગ્રામ સ્ટ્રેપ્ટોસાયક્લીનના બે છંટકાવ ૧૫ દિવસના ગળે કરવા.</p> <p>પાનકોરીયુના નિયંત્રણ માટે ડીડીવીપી ૧૦ મી.લી. ૧૦ લિટર પાણીમાં મિશ્ર કરીને છાંટવી.</p> <p>લીંબુના પાકમાં ઉનાળામાં વધુ ફળ લેવા માટે વરસાદની ઋતુ પછી વાડીને ૧૦-થી-૧૫ સે.મી. ઊંડાઈ સુધી ગોડ કરી ઝીણી મૂળ આળ કાપી નાખવી અને પાંદડા ખરવા માંડે પછી પુખ્ત ઝાડ દીઠ ૫૦ કી.ગ્રા. છાણીયું ખાતર, યુરિયા ૩૦૦ ગ્રામ, એમોનિયમ સલ્ફેટ ૫૦૦</p> |

| | | | |
|---|---|--|---|
| | | | ગ્રામ, ડી.એ.પી. ૧.૫ કી.ગ્રા. અને મ્યુરેટ ઓફ પોટાશ ૩૦૦ ગ્રામ આપી પિયત આપવું. |
| પશુ/ મરઘા  | દુધાળા પશુના આઉને ઝીંક ઓક્સાઈડ અથવા બોરીક પાવડર થી બરાબર સાફ કરવું. તેમજ ઈંતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાલ ૨ મી.લી. એક લીટર પાણીમાં નાખીને છાંટવી. પશુઓમાં ખરવા મોવાના રોગનું રસીકરણ કરાવવું. ઉછરતા પશુઓમાં કરમીયાની ગોળી આપવી. પશુને ૬ થી ૮ કી.ગ્રા. સુકો અને ૧૫ થી ૨૦ કી.ગ્રા. લીલો ચારો આપવો. ચોમાસા દરમ્યાન પશુઓના રહેઠાણની જગ્યાની વ્યવસ્થિત સફાઈ રાખવી અને માખીઓના ઉપદ્રવથી બચવા ફિનાઈલ છાંટવું. દુધાળા પશુઓને પ્રતિ લીટર દુધની ઊપજ હોય તેમને નિયમિત ૧ કી.ગ્રા. ફીડ + ૫૦ ગ્રામ ખનીજ તત્ત્વોનું મિશ્રણ આપવું. | | |

જુદ્ધયુ/ટેકનિકલ ઓફિસર.(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૧૯
તરઘડિયા(રાજકોટ), તારીખ: ૨૬/૧૧/૨૦૧૯.

પ્રતિ,
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનંતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર
સુકી ખેતીસંશોધન કેન્દ્ર
જુનાગઢ કૃષિ નિવર્સિટી
તરઘડિયા



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :942756433



Agro met Advisory Service bulletin No.69-for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 27-November to 01-December- 2019)

Significant past weather for the preceding week of Rajkot

| | |
|---------------------------|-----------|
| Rainfall (mm) | 0.0 |
| Maximum temperature (°C) | 32.3-34.1 |
| Minimum temperature (°C) | 17.0-20.0 |
| Relative Humidity (%) I | 68-80 |
| Relative Humidity (%) II | 40-50 |
| Wind speed (Kmph) | 1.9-3.3 |

Weather forecast until 08-30 hrs. Of 01/12/2019

(1) RAJKOT DISTRICT:






| | 27/11/19 | 28/11/19 | 29/11/19 | 30/11/19 | 01/12/19 |
|----------------------|----------|----------|----------|----------|----------|
| Rainfall (mm) | 0 | 0 | 0 | 0 | 0 |
| Max.Temp.(°C) | 33 | 32 | 32 | 32 | 33 |
| Min.Temp.(°C) | 20 | 20 | 19 | 19 | 19 |
| Total CC (octa) | 0 | 0 | 3 | 0 | 0 |
| RH-I (%) | 55 | 56 | 55 | 56 | 53 |
| RH-II (%) | 31 | 31 | 32 | 31 | 29 |
| Wind Speed (kmph) | 13 | 17 | 15 | 13 | 13 |
| Wind Direction(deg.) | 63 | 16 | 20 | 17 | 30 |

Agro meteorological Advisories:

| | | | |
|-----------------------|---|----------------------------|-----------------|
| General Advice | <p>Rain fall: Dry and clear sky weather forecasted next five days.</p> <ul style="list-style-type: none"> Temperature: Maximum and minimum temperature likely to be in range of 32-33°C and 19-20°C, respectively. Relative Humidity: The RH-I and RH-II is likely to be in the range of 53-56% & 29-32, respectively. Wind Direction and Speed: Direction and speed of wind is likely to be NE, N and 13-17 km/h, respectively. | | |
| Crops | Stage | Insect/Pest/Variety | Advisory |

| | | | |
|------------------|---------------------------------------|--|--|
| Garlic | Growing | - | Apply supplementary dose of nitrogen as urea @ 54 kg/ha at 30-35 days after sowing Do not over irrigate to prevent leaf blight disease. |
| Chick pea | Sowing/germination | Gujarat chana-2 (Unirrigated), Gujarat Junagadh chana-3 (irrigated) | Carry out sowing of Gram. Seed rate should be 60 to 75 kg/ha & seed should be treated with fungicides captan or thiram 2 gm/kg of seeds and apply 87 kg DAP & 10 kg Urea/ha fertilizer at the time of sowing. Before sowing, seed should be treated with Rhizobium culture and phosphorus solubilizing bacteria |
| Cumin | Sowing | Gujarat cumin-4 | Carry out sowing of cumin, If kharif crop is harvested. Seed must be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. |
| Coriander | Sowing | Gujarat Coriander-1 , Gujarat Coriander-2 | Carry out sowing of coriander & seed should be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. Apply 22 kg DAP & 13 kg Urea/ha at the time of sowing. |
| Wheat | Sowing | Lok-1, GW-496 | Carry out sowing of Wheat. Seed rate should be 100 to 125 kg/ha and apply 130 kg DAP, 80 kg Urea and 25 kg Zinc sulphate fertilizer/ha at the time of sowing. Application of Chlorpyrifos (20 EC) @ 5 litres per hectare with pre-irrigation is recommended before sowing in the fields having perennial problem of termite infestation. |
| Mustard | Sowing/germination | Varuna, Gujarat rai-3, 4 | Carry out sowing of Mustard. Seed rate should be 15 to 20 kg/ha. |
| Cotton | Flowering-ball formation /development | Jassid & whitefly Pink ball worm | <ul style="list-style-type: none"> • Also spray Mencozeb 75 % wp @ 27 gm/10 litre of water for control alternaria & cercospora leaf spot in cotton. • For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and bury the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. • Carry out picking in time. |

| | | | |
|--------------------------------|------------------------------------|--|--|
| Castor | Flowering/ capsule formation | Jassids& white fly→ Prodenia | Spray Acetamiprid 20 % SP@ 5.0 gm ORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water. For the control of prodenia spray Chlorpyriphos 25 ml in 10 liters of water. |
| Pigeon pea | Flowering/po d formation | - | Apply irrigation through drip system |
| Kharif Onion | T.P./Bulb development | Alternaria leaf spot/purple blotch | For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb27 gm and thiophanate methyl 15 gm / 10 liter of water at 10 days interval. Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DASunder clear sky. |
| Lady finger | Flowering/ fruiting | Jassids & white fly Shoot and fruit borer Mites | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid& white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthuron 50 %wp @ 16 gm / 10 lit of water. • For control of Shoot and fruit borer Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray).If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear. • Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. • After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done. |
| Cluster bean | Growing/fruiti ng | Jassid/whitefly | • Monitor the crop for infestation of jassid& white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthuron 50 %wp @ 16 gm / 10 lit of water. |
| Brinjal& Tomato | Flowering/ fruiting | Sucking pest → | Monitor the crop for infestation of jassid& white fly in brinjal, if require spray 500 ml of 5 % |

| | | | |
|--------------------------------|---|---|---|
| | | <p>Shoot & Fruit borer</p>  <p>Mites</p> | <p>neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p> |
| Chili | Flowering & fruiting | <p>Thrips →</p>  <p>Anthracnose →</p>  | <p>Under present weather conditions severe attack of thrips has been reported. Spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p> |
| Lemon | Fruit development | <p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p> <p>Hast Bahar</p> | <p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p> <p>For getting maximum fruit in summer, after last rainfall cultivate lemon orchard up to the depth of 10-15 cm for pruning of tiny small root and keep left until wilting & defoliation start, after that apply FYM @ 50 kg, Urea @ 300 gm, Ammonium sulphate @ 500 gm, DAP @ 1.5 kg & murate of potash 300 gm/tree.</p> |
| Livestock & poultry | <p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flies and mosquitoes. Keep animal shed</p>  | | |

| | |
|--|--|
| | hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed. |
|--|--|

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-69 for the Morbi district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 27-November to 01-December- 2019)

Weather forecast until 08-30 hrs of 01/12/2019


(2) MORBI DISTRICT:





| | 27/11/19 | 28/11/19 | 29/11/19 | 30/11/19 | 01/12/19 |
|----------------------|----------|----------|----------|----------|----------|
| Rainfall (mm) | 0 | 0 | 0 | 0 | 0 |
| Max. Temp.(°C) | 33 | 32 | 32 | 32 | 33 |
| Min. Temp.(°C) | 20 | 20 | 19 | 19 | 19 |
| Total CC (octa) | 0 | 1 | 3 | 1 | 1 |
| RH-I (%) | 54 | 54 | 54 | 52 | 50 |
| RH-II (%) | 32 | 35 | 30 | 28 | 27 |
| Wind Speed (kmph) | 6 | 6 | 5 | 9 | 6 |
| Wind Direction(deg.) | 105 | 62 | 68 | 45 | 96 |

Agro meteorological Advisories:-

| | | | |
|-----------------------|---|--------------------------------|--|
| General Advice | <ul style="list-style-type: none"> • Rain fall: Dry and Mainly clear to partly cloudy weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 32-33°C and 19-20°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 50-54% & 27-35, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be E, NE and 6-9 km/h, respectively. | | |
| Crops | Stage | Insect/Pest/Variety | Advisory |
| Garlic | Growing | - | Apply supplementary dose of nitrogen as urea @ 54 kg/ha at 30-35 days after sowing Do not over irrigate to prevent leaf blight disease. |
| Chick pea | Sowing/germination | Gujarat chana-2 (Unirrigated), | Carry out sowing of Gram. Seed rate should be 60 to 75 kg/ha & seed should be treated with fungicides captanorthiram 2 gm/kg of seeds and apply 87 kg DAP& |

| | | | |
|------------------|---------------------------------------|---|---|
| | | Gujarat Junagadh chana-3 (irrigated) | 10 kg Urea/ha fertilizer at the time of sowing. Before sowing, seed should be treated with Rhizobium culture and phosphorus solubilizing bacteria |
| Cumin | Sowing | Gujarat cumin-4 | Carry out sowing of cumin, If kharif crop is harvested. Seed must be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. |
| Coriander | Sowing | Gujarat Coriander-1 , Gujarat Coriander-2 | Carry out sowing of coriander & seed should be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. Apply 22 kg DAP & 13 kg Urea/ha at the time of sowing. |
| Wheat | Sowing | Lok-1, GW-496 | Carry out sowing of Wheat. Seed rate should be 100 to 125 kg/ha and apply 130 kg DAP, 80 kg Urea and 25 kg Zinc sulphate fertilizer/ha at the time of sowing. Application of Chlorpyrifos (20 EC) @ 5 litres per hectare with pre-irrigation is recommended before sowing in the fields having perennial problem of termite infestation. |
| Mustard | Sowing/germination | Varuna, Gujarat rai-3, 4 | Carry out sowing of Mustard. Seed rate should be 15 to 20 kg/ha. |
| Cotton | Flowering-ball formation /development | Jassid & whitefly Pink ball worm | <ul style="list-style-type: none"> • Also spray Mencozeb 75 % wp @ 27 gm/10 litre of water for control of alternaria & cercospora leaf spot in cotton. • For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. • Carry out picking in time. |
| Castor | Flowering/capsule formation | Jassids & white fly → Prodenia | Spray Acetamiprid 20 % SP @ 5.0 gm OR Imidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 ml / 10 lit of water ml/10 lit of water. For the control of prodenia spray Chlorpyrifos 25 ml in 10 liters of water. |
| Pigeon pea | Flowering/pod formation | - | Apply irrigation through drip system |
| Kharif | T.P./Bulb | Alternaria leaf spot/purple | For the control of purple blotch blight disease in kharif onion make 3 alternate spray of |

| | | | |
|-----------------------------|---------------------|--|---|
| Onion | development | blotch | <p>Mencozeb 27 gm and thiophanate methyl 15 gm / 10 liter of water at 10 days interval.</p> <p>Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky.</p> |
| Lady finger | Flowering/ fruiting | <p>Jassids & white fly</p> <p>Shoot and fruit borer</p> <p>Mites</p> | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. • For control of Shoot and fruit borer Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 spray). If pest population is high, spraying of Spinosad 48 EC @ 1 ml / 4 litres of water is advised when sky remains clear. • Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. • After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done. |
| Cluster bean | Growing/ fruiting | Jassid/whitefly | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. |
| Brinjal & Tomato | Flowering/ fruiting | <p>Sucking pest →</p> <p>Shoot & Fruit borer</p>  <p>Mites</p> | <p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water</p> |

| | | | |
|---|--|---|---|
| | | | for the control of Mites under clear sky. |
| Chili | Flowering & fruiting | Thrips →  Anthracnose →  | <p>Under present weather conditions severe attack of thrips has been reported. Spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p> |
| Lemon | Fruit development | Psylla and thrips  Citrus canker Leaf minor Hast Bahar | <p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p> <p>For getting maximum fruit in summer, after last rainfall cultivate lemon orchard up to the depth of 10-15 cm for pruning of tiny small root and keep left until wilting & defoliation start, after that apply FYM @ 50 kg, Urea @ 300 gm, Ammonium sulphate @ 500 gm, DAP @ 1.5 kg & murate of potash 300 gm/tree.</p> |
| Livestock & poultry  | <p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flies and mosquitoes. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p> | | |

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-69 for the Jamnagar district

(Issued jointly by DFERS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 27-November to 01-December- 2019)

Weather forecast until 08-30 hrs of 01/12/2019


(3) JAMNAGAR DISTRICT:





| | 27/11/19 | 28/11/19 | 29/11/19 | 30/11/19 | 01/12/19 |
|----------------------|----------|----------|----------|----------|----------|
| Rainfall (mm) | 0 | 0 | 0 | 0 | 0 |
| Max. Temp.(°C) | 33 | 31 | 31 | 31 | 31 |
| Min. Temp.(°C) | 21 | 21 | 21 | 20 | 20 |
| Total CC (octa) | 0 | 0 | 3 | 0 | 0 |
| RH-I (%) | 58 | 57 | 61 | 62 | 62 |
| RH-II (%) | 41 | 40 | 43 | 42 | 41 |
| Wind Speed (kmph) | 22 | 25 | 24 | 21 | 19 |
| Wind Direction(deg.) | 25 | 24 | 29 | 24 | 16 |

Agro meteorological Advisories:-

| | | | |
|-----------------------|---|--|--|
| General Advice | <ul style="list-style-type: none"> • Rain fall: Dry and clear sky weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 31-33°C and 20-21°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 57-62% & 40-43, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be NE, N and 19-25 km/h, respectively. | | |
| Crops | Stage | Insect/Pest/Variety | Advisory |
| Garlic | Growing | - | Apply supplementary dose of nitrogen as urea @ 54 kg/ha at 30-35 days after sowing Do not over irrigate to prevent leaf blight disease. |
| Chick pea | Sowing/germination | Gujarat chana-2 (Unirrigated), Gujarat Junagadh chana-3 | Carry out sowing of Gram. Seed rate should be 60 to 75 kg/ha & seed should be treated with fungicides captan or thiram 2 gm/kg of seeds and apply 87 kg DAP & 10 kg Urea/ha fertilizer at the time of sowing. Before sowing, seed should be treated with Rhizobium culture and phosphorus solubilizing bacteria |

| | | | |
|---------------------|---------------------------------------|---|---|
| | | (irrigated) | |
| Cumin | Sowing | Gujarat cumin-4 | Carry out sowing of cumin, If kharif crop is harvested. Seed must be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. |
| Coriander | Sowing | Gujarat Coriander-1 , Gujarat Coriander-2 | Carry out sowing of coriander&seed should be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. Apply 22 kg DAP&13 kg Urea/haat the time of sowing. |
| Wheat | Sowing | Lok-1, GW-496 | Carry out sowing of Wheat. Seed rate should be 100 to 125 kg/ha andapply 130 kg DAP,80 kg Ureaand 25 kg Zinc sulphatefertilizer/ha at the time of sowing.Application of Chlorpyriphos (20 EC) @ 5 litres per hectare with preirrigation is recommended before sowing in the fields having perennial problem of termite infestation. |
| Mustard | Sowing/germination | Varuna, Gujarat rai-3, 4 | Carry out sowing of Mustard.Seed rate should be 15 to 20 kg/ha. |
| Cotton | Flowering-ball formation /development | Jassid & whitefly Pink ball worm | <ul style="list-style-type: none"> • Also spray Mencozeb75 % wp @ 27gm/10 litre of water for control alternaria&cercospora leaf spot in cotton. • For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. • Carry out picking in time. |
| Castor | Flowering/capsule formation | Jassids& white fly→ Prodenia | Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water. For the control of prodenia spray Chlorpyriphos 25 ml in 10 liters of water. |
| Pigeon pea | Flowering/pod formation | - | Apply irrigation through drip system |
| Kharif Onion | T.P./Bulb development | Alternaria leaf spot/purple blotch | For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb27 gm and thiophanate methyl 15 gm / 10 liter of water at 10 days interval. |

| | | | |
|-----------------------------|---------------------|--|--|
| | | | Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky. |
| Lady finger | Flowering/ fruiting | Jassids & white fly Shoot and fruit borer Mites | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. • For control of Shoot and fruit borer Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray).If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear. • Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. • After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done. |
| Cluster bean | Growing/fruiting | Jassid/whitefly | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. |
| Brinjal & Tomato | Flowering/ fruiting | Sucking pest → Shoot & Fruit borer  Mites | <p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p> |

| | | | |
|---|--|--|---|
| <p>Chili</p> | <p>Flowering & fruiting</p> | <p>Thrips → </p> <p>Anthracnose → </p> | <p>Under present weather conditions severe attack of thrips has been reported. Spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p> |
| <p>Lemon</p> | <p>Fruit development</p> | <p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p> <p>Hast Bahar</p> | <p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p> <p>For getting maximum fruit in summer, after last rainfall cultivate lemon orchard up to the depth of 10-15 cm for pruning of tiny small root and keep left until wilting & defoliation start, after that apply FYM @ 50 kg, Urea @ 300 gm, Ammonium sulphate @ 500 gm, DAP @ 1.5 kg & murate of potash 300 gm/tree.</p> |
| <p>Livestock & poultry</p>  | <p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flies and mosquitoes. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p> | | |

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-69 for the Devbhumi Dwarka district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 27-November 01-December- 2019)

Weather forecast until 08-30 hrs of 01/12/2019


(4) DEVBHUMI DWARKA DISTRICT:





| | 27/11/19 | 28/11/19 | 29/11/19 | 30/11/19 | 01/12/19 |
|----------------------|----------|----------|----------|----------|----------|
| Rainfall (mm) | 0 | 0 | 0 | 0 | 0 |
| Max. Temp.(°C) | 31 | 29 | 29 | 29 | 29 |
| Min. Temp.(°C) | 22 | 21 | 22 | 22 | 22 |
| Total CC (octa) | 0 | 1 | 3 | 0 | 0 |
| RH-I (%) | 61 | 57 | 62 | 64 | 63 |
| RH-II (%) | 39 | 39 | 42 | 42 | 41 |
| Wind Speed (kmph) | 22 | 24 | 24 | 20 | 19 |
| Wind Direction(deg.) | 27 | 26 | 33 | 28 | 18 |

Agro meteorological Advisories:-

| | | | |
|-----------------------|---|--|--|
| General Advice | <ul style="list-style-type: none"> • Rain fall: Dry and clear sky weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 29-31°C and 21-22°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 57-64% & 39-42, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be NE, N and 19-24 km/h, respectively. | | |
| Crops | Stage | Insect/Pest/Variety | Advisory |
| Garlic | Growing | - | Apply supplementary dose of nitrogen as urea @ 54 kg/ha at 30-35 days after sowing Do not over irrigate to prevent leaf blight disease. |
| Chick pea | Sowing/germination | Gujarat chana-2 (Unirrigated), Gujarat Junagadh chana-3 | Carry out sowing of Gram. Seed rate should be 60 to 75 kg/ha & seed should be treated with fungicides captan or thiram 2 gm/kg of seeds and apply 87 kg DAP & 10 kg Urea/ha fertilizer at the time of sowing. Before sowing, seed should be treated with Rhizobium culture and phosphorus solubilizing bacteria |

| | | | |
|---------------------|---------------------------------------|---|--|
| | | (irrigated) | |
| Cumin | Sowing | Gujarat cumin-4 | Carry out sowing of cumin, If kharif crop is harvested. Seed must be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. |
| Coriander | Sowing | Gujarat Coriander-1 , Gujarat Coriander-2 | Carry out sowing of coriander&seed should be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. Apply 22 kg DAP&13 kg Urea/haat the time of sowing. |
| Wheat | Sowing | Lok-1, GW-496 | Carry out sowing of Wheat. Seed rate should be 100 to 125 kg/ha andapply 130 kg DAP,80 kg Ureaand 25 kg Zinc sulphatefertilizer/ha at the time of sowing.Application of Chlorpyriphos (20 EC) @ 5 litres per hectare with preirrigation is recommended before sowing in the fields having perennial problem of termite infestation. |
| Mustard | Sowing/germination | Varuna, Gujarat rai-3, 4 | Carry out sowing of Mustard.Seed rate should be 15 to 20 kg/ha. |
| Cotton | Flowering-ball formation /development | Jassid & whitefly Pink ball worm | <ul style="list-style-type: none"> • Also spray Mencozeb75 % wp @ 27gm/10 litre of water for control alternaria&cercospora leaf spot in cotton. • For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. • Carry out picking in time. |
| Castor | Flowering/capsule formation | Jassids& white fly→ Prodenia | Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water. For the control of prodenia spray Chlorpyriphos 25 ml in 10 liters of water. |
| Pigeon pea | Flowering/pod formation | - | Apply irrigation through drip system |
| Kharif Onion | T.P./Bulb development | Alternaria leaf spot/purple blotch | For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb27 gm and thiophanate methyl 15 gm / 10 liter of water at 10 days interval. |

| | | | |
|-----------------------------|---------------------|--|--|
| | | | Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky. |
| Lady finger | Flowering/ fruiting | Jassids & white fly Shoot and fruit borer Mites | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. • For control of Shoot and fruit borer Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray).If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear. • Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. • After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done. |
| Cluster bean | Growing/fruiting | Jassid/whitefly | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. |
| Brinjal & Tomato | Flowering/ fruiting | Sucking pest → Shoot & Fruit borer  Mites | <p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p> |

| | | | |
|--|--|--|---|
| <p>Chili</p> | <p>Flowering & fruiting</p> | <p>Thrips → </p> <p>Anthracnose → </p> | <p>Under present weather conditions severe attack of thrips has been reported. Spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p> |
| <p>Lemon</p> | <p>Fruit development</p> | <p>Psylla and thrips </p> <p>Citrus canker Leaf minor Hast Bahar</p> | <p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water. Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p> <p>For getting maximum fruit in summer, after last rainfall cultivate lemon orchard up to the depth of 10-15 cm for pruning of tiny small root and keep left until wilting & defoliation start, after that apply FYM @ 50 kg, Urea @ 300 gm, Ammonium sulphate @ 500 gm, DAP @ 1.5 kg & murate of potash 300 gm/tree.</p> |
| <p>Livestock & poultry </p> | <p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flies and mosquitoes. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p> | | |

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-69 for the Amreli district

(Issued jointly by DFERS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 27-November to 01-December- 2019)

Weather forecast until 08-30 hrs of 01/12/2019


(5) AMRELI DISTRICT:





| | 27/11/19 | 28/11/19 | 29/11/19 | 30/11/19 | 01/12/19 |
|----------------------|----------|----------|----------|----------|----------|
| Rainfall (mm) | 0 | 0 | 0 | 0 | 0 |
| Max.Temp.(°C) | 32 | 32 | 32 | 32 | 32 |
| Min.Temp.(°C) | 21 | 21 | 21 | 21 | 20 |
| Total CC (octa) | 1 | 0 | 4 | 1 | 1 |
| RH-I (%) | 58 | 58 | 56 | 54 | 53 |
| RH-II (%) | 36 | 35 | 35 | 33 | 33 |
| Wind Speed (kmph) | 10 | 15 | 15 | 14 | 14 |
| Wind Direction(deg.) | 201 | 76 | 15 | 63 | 97 |

Agro meteorological Advisories:

| | | | |
|-----------------------|---|--|---|
| General Advice | <ul style="list-style-type: none"> • Rain fall: Dry and Mainly clear to partly cloudy weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 32-32°C and 20-21°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 53-58% & 33-36, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be S, E, N, NE and 10-15 km/h, respectively. | | |
| Crops | Stage | Insect/Pest/ Variety | Advisory |
| Garlic | Growing | - | Apply supplementary dose of nitrogen as urea @ 54 kg/ha at 30-35 days after sowing Do not over irrigate to prevent leaf blight disease. |
| Chick pea | Sowing/germination | Gujarat chana-2 (Unirrigated), Gujarat Junagadh chana-3 | Carry out sowing of Gram. Seed rate should be 60 to 75 kg/ha & seed should be treated with fungicides captanorthiram 2 gm/kg of seeds and apply 87 kg DAP & 10 kg Urea/ha fertilizer at the time of sowing. Before sowing, seed should be treated with Rhizobium |

| | | | |
|---------------------|---------------------------------------|---|---|
| | | (irrigated) | culture and phosphorus solublesing bacteria |
| Cumin | Sowing | Gujarat cumin-4 | Carry out sowing of cumin, If kharif crop is harvested. Seed must be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. |
| Coriander | Sowing | Gujarat Coriander-1 , Gujarat Coriander-2 | Carry out sowing of coriander&seed should be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. Apply 22 kg DAP&13 kg Urea/haat the time of sowing. |
| Wheat | Sowing | Lok-1, GW-496 | Carry out sowing of Wheat. Seed rate should be 100 to 125 kg/ha andapply 130 kg DAP,80 kg Ureaand 25 kg Zinc sulphatefertilizer/ha at the time of sowing.Application of Chlorpyriphos (20 EC) @ 5 litres per hectare with preirrigation is recommended before sowing in the fields having perennial problem of termite infestation. |
| Mustard | Sowing/germination | Varuna, Gujarat rai-3, 4 | Carry out sowing of Mustard.Seed rate should be 15 to 20 kg/ha. |
| Cotton | Flowering-ball formation /development | Jassid & whitefly Pink ball worm | <ul style="list-style-type: none"> • Also spray Mencozeb75 % wp @ 27gm/10 litre of water for control alternaria&cercospora leaf spot in cotton. • For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. • Carry out picking in time. |
| Castor | Flowering/capsule formation | Jassids& white fly→ Prodenia | Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water. For the control of prodenia spray Chlorpyriphos 25 ml in 10 liters of water. |
| Pigeon pea | Flowering/pod formation | - | Apply irrigation through drip system |
| Kharif Onion | T.P./Bulb development | Alternaria leaf spot/purple blotch | For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb27 gm and thiophanate methyl 15 gm / 10 liter of water at 10 days interval. |

| | | | |
|-----------------------------|---------------------|--|--|
| | | | Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky. |
| Lady finger | Flowering/ fruiting | Jassids & white fly Shoot and fruit borer Mites | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. • For control of Shoot and fruit borer Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray).If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear. • Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. • After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done. |
| Cluster bean | Growing/fruiting | Jassid/whitefly | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. |
| Brinjal & Tomato | Flowering/ fruiting | Sucking pest → Shoot & Fruit borer  Mites | <p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p> |

| | | | |
|---|--|--|---|
| <p>Chili</p> | <p>Flowering & fruiting</p> | <p>Thrips → </p> <p>Anthracnose → </p> | <p>Under present weather conditions severe attack of thrips has been reported. Spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p> |
| <p>Lemon</p> | <p>Fruit development</p> | <p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p> <p>Hast Bahar</p> | <p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p> <p>For getting maximum fruit in summer, after last rainfall cultivate lemon orchard up to the depth of 10-15 cm for pruning of tiny small root and keep left until wilting & defoliation start, after that apply FYM @ 50 kg, Urea @ 300 gm, Ammonium sulphate @ 500 gm, DAP @ 1.5 kg & murate of potash 300 gm/tree.</p> |
| <p>Livestock & poultry</p>  | <p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flies and mosquitoes. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p> | | |

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:94277 28523



Agro met Advisory Service bulletin No.-69 for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 27-November to 01-December- 2019)

Weather forecast until 08-30 hrs of 01/12/2019


(6) SURENDRANAGAR DISTRICT:





| | 27/11/19 | 28/11/19 | 29/11/19 | 30/11/19 | 01/12/19 |
|----------------------|----------|----------|----------|----------|----------|
| Rainfall (mm) | 0 | 0 | 0 | 0 | 0 |
| Max.Temp.(°C) | 33 | 32 | 32 | 32 | 33 |
| Min.Temp.(°C) | 20 | 20 | 19 | 19 | 19 |
| Total CC (octa) | 0 | 0 | 3 | 0 | 0 |
| RH-I (%) | 55 | 56 | 55 | 54 | 52 |
| RH-II (%) | 33 | 35 | 35 | 35 | 34 |
| Wind Speed (kmph) | 14 | 17 | 15 | 12 | 13 |
| Wind Direction(deg.) | 62 | 47 | 40 | 33 | 114 |

Agro meteorological advisories:

| | | | |
|-----------------------|--|--|---|
| General Advice | <ul style="list-style-type: none"> • Rain fall: Dry and clear sky weather forecasted next five days. • Temperature: Maximum and minimum temperature likely to be in range of 32-33°C and 19-20°C, respectively. • Relative Humidity: The RH-I and RH-II is likely to be in the range of 52-56% & 33-35, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be NE, SE and 12-17 km/h, respectively. | | |
| Crops | Stage | Insect/Pest/Variety | Advisory |
| Garlic | Growing | - | Apply supplementary dose of nitrogen as urea @ 54 kg/ha at 30-35 days after sowing Do not over irrigate to prevent leaf blight disease. |
| Chick pea | Sowing/germination | Gujarat chana-2 (Unirrigated), Gujarat Junagadh chana-3 | Carry out sowing of Gram. Seed rate should be 60 to 75 kg/ha & seed should be treated with fungicides captan or thiram 2 gm/kg of seeds and apply 87 kg DAP & 10 kg Urea/ha fertilizer at the time of sowing. Before sowing, seed should be treated with Rhizobium |

| | | | |
|---------------------|---------------------------------------|---|---|
| | | (irrigated) | culture and phosphorus solublesing bacteria |
| Cumin | Sowing | Gujarat cumin-4 | Carry out sowing of cumin, If kharif crop is harvested. Seed must be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. |
| Coriander | Sowing | Gujarat Coriander-1 , Gujarat Coriander-2 | Carry out sowing of coriander&seed should be treated with fungicides Captan or Thiram or Carbendazim @ 2.0 gm/ kg to prevent wilt. Apply 22 kg DAP&13 kg Urea/haat the time of sowing. |
| Wheat | Sowing | Lok-1, GW-496 | Carry out sowing of Wheat. Seed rate should be 100 to 125 kg/ha andapply 130 kg DAP,80 kg Ureaand 25 kg Zinc sulphatefertilizer/ha at the time of sowing.Application of Chlorpyriphos (20 EC) @ 5 litres per hectare with preirrigation is recommended before sowing in the fields having perennial problem of termite infestation. |
| Mustard | Sowing/germination | Varuna, Gujarat rai-3, 4 | Carry out sowing of Mustard.Seed rate should be 15 to 20 kg/ha. |
| Cotton | Flowering-ball formation /development | Jassid & whitefly Pink ball worm | <ul style="list-style-type: none"> • Also spray Mencozeb75 % wp @ 27gm/10 litre of water for control alternaria&cercospora leaf spot in cotton. • For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha in early sown crop. If infestation found pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water. • Carry out picking in time. |
| Castor | Flowering/capsule formation | Jassids& white fly→ Prodenia | Spray Acetamiprid 20 % SP@ 5.0 gmORimidacloprid 30.5 % 3 ml with neem seed extract 500 ml or neem oil 40 mla / 10 lit of water ml/10 lit of water. For the control of prodenia spray Chlorpyriphos 25 ml in 10 liters of water. |
| Pigeon pea | Flowering/pod formation | - | Apply irrigation through drip system |
| Kharif Onion | T.P./Bulb development | Alternaria leaf spot/purple blotch | For the control of purple blotch blight disease in kharif onion make 3 alternate spray of Mencozeb27 gm and thiophanate methyl 15 gm / 10 liter of water at 10 days interval. |

| | | | |
|-----------------------------|---------------------|--|--|
| | | | Apply supplementary dose of nitrogen (Urea 40 kg/ha) at 30 DAS under clear sky. |
| Lady finger | Flowering/ fruiting | Jassids & white fly Shoot and fruit borer Mites | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. • For control of Shoot and fruit borer Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray).If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear. • Constant monitoring of lady finger crop against attack of mite is advised. If population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water is advised when sky remains clear. • After harvesting of mature okra application of urea @ 5-10 kg/ acre should be done. |
| Cluster bean | Growing/fruiting | Jassid/whitefly | <ul style="list-style-type: none"> • Monitor the crop for infestation of jassid & white fly, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water. |
| Brinjal & Tomato | Flowering/ fruiting | Sucking pest → Shoot & Fruit borer  Mites | <p>Monitor the crop for infestation of jassid & white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 %wp @ 16 gm / 10 lit of water.</p> <p>In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10litre of water for the control of Mites under clear sky.</p> |

| | | | |
|---|--|--|---|
| <p>Chili</p> | <p>Flowering & fruiting</p> | <p>Thrips → </p> <p>Anthracnose → </p> | <p>Under present weather conditions severe attack of thrips has been reported. Spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p> <p>For the control of Anthracnose spray Mencozeb 27gm OR Chlorothalonil 27 gram OR Fosetyl 20 gm in 10 liters of water at 15 days interval.</p> |
| <p>Lemon</p> | <p>Fruit development</p> | <p>Psylla and thrips</p>  <p>Citrus canker</p> <p>Leaf minor</p> <p>Hast Bahar</p> | <p>To prevent damage of psylla & thrips spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.</p> <p>Spray copper oxychloride 40 gm or streptocyclin 1.5 gm in 10 litre of water.</p> <p>Spray Dichlorvos 10 ml or Quinalphos 20 ml in 10 litres of water for control of leaf minor.</p> <p>For getting maximum fruit in summer, after last rainfall cultivate lemon orchard up to the depth of 10-15 cm for pruning of tiny small root and keep left until wilting & defoliation start, after that apply FYM @ 50 kg, Urea @ 300 gm, Ammonium sulphate @ 500 gm, DAP @ 1.5 kg & murate of potash 300 gm/tree.</p> |
| <p>Livestock & poultry</p>  | <p>Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flies and mosquitoes. Keep animal shed hygienic and clean to avoid flies and mosquitoes. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed.</p> | | |

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.